

DIRECTED EVOLUTION OF NOVEL BINDING PROTEINS

This application is a continuation-in-part of Ladner, Guterman, Roberts, and Markland, Ser. No. 07/487,063, filed March 2, 1990, now pending, which is a continuation-in-part of Ladner and Guterman, Ser. No. 07/240,160, filed Sept. 2, 1988, now pending. Ser. No. 07/487,063 claimed priority under 35 U.S.C. 119 from PCT Application No. PCT/US89/03731, filed Sept. 1, 1989. All of the foregoing applications are hereby incorporated by reference.

10 Cross-Reference to Related Applications

The following related and commonly-owned applications are also incorporated by reference:

Robert Charles Ladner, Sonia Kosow Guterman, Rachael Baribault Kent, and Arthur Charles Ley are named as joint 15 inventors on U.S.S.N. 07/293,980, filed January 8, 1989, and entitled GENERATION AND SELECTION OF NOVEL DNA-BINDING PROTEINS AND POLYPEPTIDES. This application has been assigned to Protein Engineering Corporation.

Robert Charles Ladner, Sonia Kosow Guterman, and 20 Bruce Lindsay Roberts are named as a joint inventors on a U.S.S.N. 07/470,651 filed 26 January 1990, entitled "PRODUCTION OF NOVEL SEQUENCE-SPECIFIC DNA-ALTERING ENZYMES", likewise assigned to Protein Engineering Corp.

Ladner, Guterman, Kent, Ley, and Markland, Ser. No. 25 07/558,011 is also assigned to Protein Engineering Corporation.

BACKGROUND OF THE INVENTION

Field of the Invention

30 This invention relates to development of novel binding proteins (including mini-proteins) by an iterative process of mutagenesis, expression, chromatographic

FOIA b 7 - D

selection, and amplification. In this process, a gene encoding a potential binding domain, said gene being obtained by random mutagenesis of a limited number of predetermined codons, is fused to a genetic element which causes the resulting chimeric expression product to be displayed on the outer surface of a virus (especially a filamentous phage) or a cell. Chromatographic selection is then used to identify viruses or cells whose genome includes such a fused gene which coded for the protein which bound to the chromatographic target.

Information Disclosure Statement

A. Protein Structure

The amino acid sequence of a protein determines its three-dimensional (3D) structure, which in turn determines protein function (EPST63, ANFI73). Shortle (SHOR85), Sauer and colleagues (PAKU86, REID88a), and Caruthers and colleagues (EISE85) have shown that some residues on the polypeptide chain are more important than others in determining the 3D structure of a protein. The 3D structure is essentially unaffected by the identity of the amino acids at some loci; at other loci only one or a few types of amino acid is allowed. In most cases, loci where wide variety is allowed have the amino acid side group directed toward the solvent. Loci where limited variety is allowed frequently have the side group directed toward other parts of the protein. Thus substitutions of amino acids that are exposed to solvent are less likely to affect the 3D structure than are substitutions at internal loci. (See also SCHU79, p169-171 and CREI84, p239-245, 314-315).

The secondary structure (helices, sheets, turns, loops) of a protein is determined mostly by local sequence. Certain amino acids have a propensity to appear in certain "secondary structures," they will be found from

FOI 56096860

time to time in other structures, and studies of pentapeptide sequences found in different proteins have shown that their conformation varies considerably from one occurrence to the next (KABS84, ARG087). As a result, a priori design of proteins to have a particular 3D structure is difficult.

Several researchers have designed and synthesized proteins de novo (MOSE83, MOSE87, ERIC86). These designed proteins are small and most have been synthesized in vitro as polypeptides rather than genetically. Hecht et al. (HECH90) have produced a designed protein genetically. Moser, et al. state that design of biologically active proteins is currently impossible.

B. Protein Binding Activity

Many proteins bind non-covalently but very tightly and specifically to some other characteristic molecules (SCHU79, CREI84). In each case the binding results from complementarity of the surfaces that come into contact: bumps fit into holes, unlike charges come together, dipoles align, and hydrophobic atoms contact other hydrophobic atoms. Although bulk water is excluded, individual water molecules are frequently found filling space in intermolecular interfaces; these waters usually form hydrogen bonds to one or more atoms of the protein or to other bound water. Thus proteins found in nature have not attained, nor do they require, perfect complementarity to bind tightly and specifically to their substrates. Only in rare cases is there essentially perfect complementarity; then the binding is extremely tight (as for example, avidin binding to biotin).

C. Protein Engineering

"Protein engineering" is the art of manipulating the sequence of a protein in order to alter its binding

FOIb290" 56096860

characteristics. The factors affecting protein binding are known, (CHOT75, CHOT76, SCHU79, p98-107, and CREI84, Ch8), but designing new complementary surfaces has proved difficult. Although some rules have been developed for substituting side groups (SUTC87b), the side groups of proteins are floppy and it is difficult to predict what conformation a new side group will take. Further, the forces that bind proteins to other molecules are all relatively weak and it is difficult to predict the effects of these forces.

Recently, Quioco and collaborators (QUIO87) elucidated the structures of several periplasmic binding proteins from Gram-negative bacteria. They found that the proteins, despite having low sequence homology and differences in structural detail, have certain important structural similarities. Based on their investigations of these binding proteins, Quioco et al. suggest it is unlikely that, using current protein engineering methods, proteins can be constructed with binding properties superior to those of proteins that occur naturally.

Nonetheless, there have been some isolated successes. Wilkinson et al. (WILK84) reported that a mutant of the tyrosyl tRNA synthetase of Bacillus stearothermophilus with the mutation Thr₅₁→Pro exhibits a 100-fold increase in affinity for ATP. Tan and Kaiser (TANK77) and Tschesche et al. (TSCH87) showed that changing a single amino acid in mini-protein greatly reduces its binding to trypsin, but that some of the mutants retained the parental characteristic of binding to an inhibiting chymotrypsin, while others exhibited new binding to elastase. Caruthers and others (EISE85) have shown that changes of single amino acids on the surface of the lambda Cro repressor greatly reduce its affinity for the natural operator O_R3, but greatly increase the binding of the

100-50-5000000000

mutant protein to a mutant operator. Changing three residues in subtilisin from Bacillus amyloliquefaciens to be the same as the corresponding residues in subtilisin from B. licheniformis produced a protease having nearly the same activity as the latter subtilisin, even though 82 amino acid sequence differences remained (WELL87a). Insertion of DNA encoding 18 amino acids (corresponding to Pro-Glu-Dynorphin-Gly) into the E. coli phoA gene so that the additional amino acids appeared within a loop of the alkaline phosphatase protein resulted in a chimeric protein having both phoA and dynorphin activity (FREI90). Thus, changing the surface of a binding protein may alter its specificity without abolishing binding activity.

D. Techniques Of Mutagenesis

Early techniques of mutating proteins involved manipulations at the amino acid sequence level. In the semi-synthetic method (TSCH87), the protein was cleaved into two fragments, a residue removed from the new end of one fragment, the substitute residue added on in its place, and the modified fragment joined with the other, original fragment. Alternatively, the mutant protein could be synthesized in its entirety (TANK77).

Erickson et al. suggested that mixed amino acid reagents could be used to produce a family of sequence-related proteins which could then be screened by affinity chromatography (ERIC86). They envision successive rounds of mixed synthesis of variant proteins and purification by specific binding. They do not discuss how residues should be chosen for variation. Because proteins cannot be amplified, the researchers must sequence the recovered protein to learn which substitutions improve binding. The researchers must limit the level of diversity so that each variety of protein will be present in sufficient quantity for the isolated fraction to be sequenced.

T06230-5695860

With the development of recombinant DNA techniques, it became possible to obtain a mutant protein by mutating the gene encoding the native protein and then expressing the mutated gene. Several mutagenesis strategies are known. One, "protein surgery" (DILL87), involves the introduction of one or more predetermined mutations within the gene of choice. A single polypeptide of completely predetermined sequence is expressed, and its binding characteristics are evaluated.

10 At the other extreme is random mutagenesis by means of relatively nonspecific mutagens such as radiation and various chemical agents. See Ho et al. (HOCJ85) and Lehtovaara, E.P. Appln. 285,123.

It is possible to randomly vary predetermined
 15 nucleotides using a mixture of bases in the appropriate cycles of a nucleic acid synthesis procedure. The proportion of bases in the mixture, for each position of a codon, will determine the frequency at which each amino acid will occur in the polypeptides expressed from the
 20 degenerate DNA population. Oliphant et al. (OLIP86) and Oliphant and Struhl (OLIP87) have demonstrated ligation and cloning of highly degenerate oligonucleotides, which were used in the mutation of promoters. They suggested that similar methods could be used in the variation of
 25 protein coding regions. They do not say how one should: a) choose protein residues to vary, or b) select or screen mutants with desirable properties. Reidhaar-Olson and Sauer (REID88a) have used synthetic degenerate oligo-nts to vary simultaneously two or three residues through all
 30 twenty amino acids. See also Vershon et al. (VERS86a; VERS86b). Reidhaar-Olson and Sauer do not discuss the limits on how many residues could be varied at once nor do they mention the problem of unequal abundance of DNA encoding different amino acids. They looked for proteins

TO 50950 56096850

that either had wild-type dimerization or that did not dimerize. They did not seek proteins having novel binding properties and did not find any. This approach is likewise limited by the number of colonies that can be
5 examined (ROBE86).

To the extent that this prior work assumes that it is desirable to adjust the level of mutation so that there is one mutation per protein, it should be noted that many desirable protein alterations require multiple amino acid
10 substitutions and thus are not accessible through single base changes or even through all possible amino acid -- substitutions at any one residue.

D. Affinity Chromatography of Cells

Ferenci and coloborators have published a series of
15 papers on the chromatographic isolation of mutants of the maltose-transport protein LamB of E. coli (FERE82a, FERE82b, FERE83, FERE84, CLUN84, HEIN87 and papers cited therein). The mutants were either spontaneous or induced with nonspecific chemical mutagens. Levels of mutagenesis
20 were picked to provide single point mutations or single insertions of two residues. No multiple mutations were sought or found.

While variation was seen in the degree of affinity for the conventional LamB substrates maltose and starch,
25 there was no selection for affinity to a target molecule not bound at all by native LamB, and no multiple mutations were sought or found. FERE84 speculated that the affinity chromatographic selection technique could be adapted to development of similar mutants of other "important
30 bacterial surface-located enzymes", and to selecting for mutations which result in the relocation of an intracellular bacterial protein to the cell surface. Ferenci's mutant surface proteins would not, however, have been chimeras of a bacterial surface protein and an exogenous

FO6290" 55096860

or heterologous binding domain.

Ferenci also taught that there was no need to clone the structural gene, or to know the protein structure, active site, or sequence. The method of the present invention, however, specifically utilizes a cloned structural gene. It is not possible to construct and express a chimeric, outer surface-directed potential binding protein-encoding gene without cloning.

Ferenci did not limit the mutations to particular loci or particular substitutions. In the present invention, knowledge of the protein structure, active site and/or sequence is used as appropriate to predict which residues are most likely to affect binding activity without unduly destabilizing the protein, and the mutagenesis is focused upon those sites. Ferenci does not suggest that surface residues should be preferentially varied. In consequence, Ferenci's selection system is much less efficient than that disclosed herein.

E. Bacterial and Viral Expression of Chimeric Surface Proteins

A number of researchers have directed unmutated foreign antigenic epitopes to the surface of bacteria or phage, fused to a native bacterial or phage surface protein, and demonstrated that the epitopes were recognized by antibodies. Thus, Charbit, et al. (CHAR86) genetically inserted the C3 epitope of the VP1 coat protein of poliovirus into the LamB outer membrane protein of E. coli, and determined immunologically that the C3 epitope was exposed on the bacterial cell surface. Charbit, et al. (CHAR87) likewise produced chimeras of LamB and the A (or B) epitopes of the preS2 region of hepatitis B virus.

A chimeric LacZ/OmpB protein has been expressed in E.

T06290" 55095B50

Dulbecco (DULB86) suggests a procedure for incorporating a foreign antigenic epitope into a viral surface protein so that the expressed chimeric protein is displayed on the surface of the virus in a manner such that the foreign epitope is accessible to antibody. In 1985 Smith (SMIT85) reported inserting a nonfunctional segment of the EcoRI endonuclease gene into gene III of bacteriophage f1, "in phase". The gene III protein is a minor coat protein necessary for infectivity. Smith demonstrated that the recombinant phage were adsorbed by immobilized antibody raised against the EcoRI endonuclease, and could be eluted with acid. De la Cruz et al. (DELA88) have expressed a fragment of the repeat region of the circumsporozoite protein from Plasmodium falciparum on the surface of M13 as an insert in the gene III protein. They showed that the recombinant phage were both antigenic and immunogenic in rabbits, and that such recombinant phage could be used for B epitope mapping. The researchers suggest that similar recombinant phage could be used for T epitope mapping and for vaccine development.

None of these researchers suggested mutagenesis of the inserted material, nor is the inserted material a complete binding domain conferring on the chimeric protein the ability to bind specifically to a receptor other than

the antigen combining site of an antibody.

McCafferty et al. (MCCA90) expressed a fusion of an Fv fragment of an antibody to the N-terminal of the pIII protein. The Fv fragment was not mutated.

5 F. Epitope Libraries on Fusion Phage

Parmley and Smith (PARM88) suggested that an epitope library that exhibits all possible hexapeptides could be constructed and used to isolate epitopes that bind to antibodies. In discussing the epitope library, the authors did not suggest that it was desirable to balance the representation of different amino acids. Nor did they teach that the insert should encode a complete domain of the exogenous protein. Epitopes are considered to be unstructured peptides as opposed to structured proteins.

After the filing of the parent application whose benefit is claimed herein under 35 U.S.C. 120, certain groups reported the construction of "epitope libraries." Scott and Smith (SCOT90) and Cwirla et al. (CWIR90) prepared "epitope libraries" in which potential hexapeptide epitopes for a target antibody were randomly mutated by fusing degenerate oligonucleotides, encoding the epitopes, with gene III of fd phage, and expressing the fused gene in phage-infected cells. The cells manufactured fusion phage which displayed the epitopes on their surface; the phage which bound to immobilized antibody were eluted with acid and studied. In both cases, the fused gene featured a segment encoding a spacer region to separate the variable region from the wild type pIII sequence so that the varied amino acids would not be constrained by the nearby pIII sequence. Devlin et al. (DEVL90) similarly screened, using M13 phage, for random residue epitopes recognized by streptavidin. Again, a spacer was used to move the random peptides away from the rest of the chimeric phage protein. These references

FD-290" 56095850

therefore taught away from constraining the conformational repertoire of the mutated residues.

Another problem with the Scott and Smith, Cwirla et al., and Devlin et al., libraries was that they provided a highly biased sampling of the possible amino acids at each position. Their primary concern in designing the degenerate oligonucleotide encoding their variable region was to ensure that all twenty amino acids were encodable at each position; a secondary consideration was minimizing the frequency of occurrence of stop signals. Consequently, Scott and Smith and Cwirla et al. employed NNK (N=equal mixture of G, A, T, C; K=equal mixture of G and T) while Devlin et al. used NNS (S=equal mixture of G and C). There was no attempt to minimize the frequency ratio of most favored-to-least favored amino acid, or to equalize the rate of occurrence of acidic and basic amino acids.

Devlin et al. characterized several affinity-selected streptavidin-binding peptides, but did not measure the affinity constants for these peptides. Cwirla et al. did determine the affinity constant for his peptides, but were disappointed to find that his best hexapeptides had affinities (350-300nM), "orders of magnitude" weaker than that of the native Met-enkephalin epitope (7nM) recognized by the target antibody. Cwirla et al. speculated that phage bearing peptides with higher affinities remained bound under acidic elution, possibly because of multivalent interactions between phage (carrying about 4 copies of pIII) and the divalent target IgG. Scott and Smith were able to find peptides whose affinity for the target antibody (A2) was comparable to that of the reference myohemerythrin epitope (50nM). However, Scott and Smith likewise expressed concern that some high-affinity peptides were lost, possibly through irreversible binding of fusion phage to target.

FOIb290-56096860

G. Non-Commonly Owned Patents and Applications Naming Robert Ladner as an Inventor

Ladner, US Patent No. 4,704,692, "Computer Based System and Method for Determining and Displaying Possible
5 Chemical Structures for Converting Double- or Multiple-Chain Polypeptides to Single-Chain Polypeptides" describes a design method for converting proteins composed of two or more chains into proteins of fewer polypeptide chains, but with essentially the same 3D structure. There is no
10 mention of variegated DNA and no genetic selection.

Ladner and Bird, WO88/01649 (Publ. March 10, 1988) disclose the specific application of computerized design of linker peptides to the preparation of single chain antibodies.

15 Ladner, Glick, and Bird, WO88/06630 (publ. 7 Sept.
1988 and having priority from US application 07/021,046,
assigned to Genex Corp.) (LGB) speculate that diverse
single chain antibody domains (SCAD) may be screened for
binding to a particular antigen by varying the DNA
20 encoding the combining determining regions of a single
chain antibody, subcloning the SCAD gene into the gpV gene
of phage lambda so that a SCAD/gpV chimera is displayed on
the outer surface of phage lambda, and selecting phage
which bind to the antigen through affinity chromatography.
25 The only antigen mentioned is bovine growth hormone. No
other binding molecules, targets, carrier organisms, or
outer surface proteins are discussed. Nor is there any
mention of the method or degree of mutagenesis. Further-
more, there is no teaching as to the exact structure of
30 the fusion nor of how to identify a successful fusion or
how to proceed if the SCAD is not displayed.

Ladner and Bird, WO88/06601 (publ. 7 September 1988) suggest that single chain "pseudodimeric" repressors (DNA-binding proteins) may be prepared by mutating a

putative linker peptide followed by in vivo selection that mutation and selection may be used to create a dictionary of recognition elements for use in the design of asymmetric repressors. The repressors are not displayed on
5 the outer surface of an organism.

Methods of identifying residues in protein which can be replaced with a cysteine in order to promote the formation of a protein-stabilizing disulfide bond are given in Pantoliano and Ladner, U.S. Patent No. 4,903,773
10 (PANT90), Pantoliano and Ladner (PANT87), Pabo and Suchenek (PAB086), MATS89, and SAUE86.

No admission is made that any cited reference is prior art or pertinent prior art, and the dates given are
15 those appearing on the reference and may not be identical to the actual publication date. All references cited in this specification are hereby incorporated by reference.

0935095.062001
T06250" 55055560

SUMMARY OF THE INVENTION

The present invention is intended to overcome the deficiencies discussed above. It relates to the construction, expression, and selection of mutated genes that specify novel proteins with desirable binding properties, as well as these proteins themselves. The substances bound by these proteins, hereinafter referred to as "targets", may be, but need not be, proteins. Targets may include other biological or synthetic macromolecules as well as other organic and inorganic substances.

The fundamental principle of the invention is one of forced evolution. In nature, evolution results from the combination of genetic variation, selection for advantageous traits, and reproduction of the selected individuals, thereby enriching the population for the trait. The present invention achieves genetic variation through controlled random mutagenesis ("variegation") of DNA, yielding a mixture of DNA molecules encoding different but related potential binding proteins. It selects for mutated genes that specify novel proteins with desirable binding properties by 1) arranging that the product of each mutated gene be displayed on the outer surface of a replicable genetic package (GP) (a cell, spore or virus) that contains the gene, and 2) using affinity selection-- selection for binding to the target material -- to enrich the population of packages for those packages containing genes specifying proteins with improved binding to that target material. Finally, enrichment is achieved by allowing only the genetic packages which, by virtue of the displayed protein, bound to the target, to reproduce. The evolution is "forced" in that selection is for the target material provided.

The display strategy is first perfected by modifying

a genetic package to display a stable, structured domain (the "initial potential binding domain", IPBD) for which an affinity molecule (which may be an antibody) is obtainable. The success of the modifications is readily
5 measured by, e.g., determining whether the modified genetic package binds to the affinity molecule.

The IPBD is chosen with a view to its tolerance for extensive mutagenesis. Once it is known that the IPBD can be displayed on a surface of a package and subjected to
10 affinity selection, the gene encoding the IPBD is subjected to a special pattern of multiple mutagenesis, here termed "variegation", which after appropriate cloning and amplification steps leads to the production of a population of genetic packages each of which displays a single
15 potential binding domain (a mutant of the IPBD), but which collectively display a multitude of different though structurally related potential binding domains (PBDs). Each genetic package carries the version of the pbd gene that encodes the PBD displayed on the surface of that particular package. Affinity selection is then used to
20 identify the genetic packages bearing the PBDs with the desired binding characteristics, and these genetic packages may then be amplified. After one or more cycles of enrichment by affinity selection and amplification, the
25 DNA encoding the successful binding domains (SBDs) may then be recovered from selected packages.

If need be, the DNA from the SBD-bearing packages may then be further "variegated", using an SBD of the last round of variegation as the "parental potential binding
30 domain" (PPBD) to the next generation of PBDs, and the process continued until the worker in the art is satisfied with the result. At that point, the SBD may be produced by any conventional means, including chemical synthesis.

When the number of different amino acid sequences

obtainable by mutation of the domain is large when compared to the number of different domains which are displayable in detectable amounts, the efficiency of the forced evolution is greatly enhanced by careful choice of which residues are to be varied. First, residues of a known protein which are likely to affect its binding activity (e.g., surface residues) and not likely to unduly degrade its stability are identified. Then all or some of the codons encoding these residues are varied simultaneously to produce a variegated population of DNA. The variegated population of DNA is used to express a variety of potential binding domains, whose ability to bind the target of interest may then be evaluated.

The method of the present invention is thus further distinguished from other methods in the nature of the highly variegated population that is produced and from which novel binding proteins are selected. We force the displayed potential binding domain to sample the nearby "sequence space" of related amino-acid sequences in an efficient, organized manner. Four goals guide the various variegation plans used herein, preferably: 1) a very large number (e.g. 10^7) of variants is available, 2) a very high percentage of the possible variants actually appears in detectable amounts, 3) the frequency of appearance of the desired variants is relatively uniform, and 4) variation occurs only at a limited number of amino-acid residues, most preferably at residues having side groups directed toward a common region on the surface of the potential binding domain.

This is to be distinguished from the simple use of indiscriminate mutagenic agents such as radiation and hydroxylamine to modify a gene, where there is no (or very oblique) control over the site of mutation. Many of the mutations will affect residues that are not a part of the

binding domain. Moreover, since at a reasonable level of mutagenesis, any modified codon is likely to be characterized by a single base change, only a limited and biased range of possibilities will be explored. Equally remote is the use of site-specific mutagenesis techniques employing mutagenic oligonucleotides of nonrandomized sequence, since these techniques do not lend themselves to the production and testing of a large number of variants. While focused random mutagenesis techniques are known, the importance of controlling the distribution of variation has been largely overlooked.

In order to obtain the display of a multitude of different though related potential binding domains, applicants generate a heterogeneous population of replicable genetic packages each of which comprises a hybrid gene including a first DNA sequence which encodes a potential binding domain for the target of interest and a second DNA sequence which encodes a display means, such as an outer surface protein native to the genetic package but not natively associated with the potential binding domain (or the parental binding domain to which it is related) which causes the genetic package to display the corresponding chimeric protein (or a processed form thereof) on its outer surface.

It should be recognized that by expressing a hybrid protein which comprises an outer surface transport signal not natively associated with the binding domain, the utility of the present invention is greatly extended. The binding domain need not be that of a surface protein of the genetic package (or, in the case of a viral package, of its host cell), since the provided outer surface transport signal is responsible for achieving the desired display. Thus, it is possible to display on the surface of a phage, bacterial cell or bacterial spore a binding

domain related to the binding domain of a normally cytoplasmic binding protein, or the binding domain of eukaryotic protein which is not found on the surface of prokaryotic cells or viruses.

5 Another important aspect of the invention is that each potential binding domain remains physically associated with the particular DNA molecule which encodes it. Thus, once successful binding domains are identified, one may readily recover the gene and either express additional
10 quantities of the novel binding protein or further mutate the gene. The form that this association takes is a "replicable genetic package", a virus, cell or spore which replicates and expresses the binding domain-encoding gene, and transports the binding domain to its outer surface.

15 It is also possible chemically or enzymatically to modify the PBDs before selection. The selection then identifies the best modified amino acid sequence. For example, we could treat the variegated population of genetic packages that display a variegated population of
20 binding domains with a protein tyrosine kinase and then select for binding the target. Any tyrosines on the BD surface will be phosphorylated and this could affect the binding properties. Other chemical or enzymatic modifications are possible.

25 By virtue of the present invention, proteins are obtained which can bind specifically to targets other than the antigen-combining sites of antibodies. A protein is not to be considered a "binding protein" merely because it can be bound by an antibody (see definition of "binding
30 protein" which follows). While almost any amino acid sequence of more than about 6-8 amino acids is likely, when linked to an immunogenic carrier, to elicit an immune response, any given random polypeptide is unlikely to satisfy the stringent definition of "binding protein" with

FOIA b 7 - D

respect to minimum affinity and specificity for its substrate. It is only by testing numerous random polypeptides simultaneously (and, in the usual case, controlling the extent and character of the sequence variation, i.e.,
 5 limiting it to residues of a potential binding domain having a stable structure, the residues being chosen as more likely to affect binding than stability) that this obstacle is overcome.

In one embodiment, the invention relates to:

- 10 a) preparing a variegated population of replicable genetic packages, each package including a nucleic acid construct coding for an outer-surface-displayed potential binding protein other than an antibody, comprising (i) a structural signal directing the
 15 display of the protein (or a processed form thereof) on the outer surface of the package and (ii) a potential binding domain for binding said target, where the population collectively displays a multitude of different potential binding domains having a substantially predetermined range of variation in
 20 sequence,
- b) causing the expression of said protein and the display of said protein on the outer surface of such packages,
- 25 c) contacting the packages with target material, other than an antibody with an exposed antigen-combining site, so that the potential binding domains of the proteins and the target material may interact, and separating packages bearing a potential binding
 30 domain that succeeds in binding the target material from packages that do not so bind,
- d) recovering and replicating at least one package bearing a successful binding domain,
- e) determining the amino acid sequence of the successful

FOIA b 7 - D

binding domain of a genetic package which bound to the target material,

- 5 f) preparing a new variegated population of replicable genetic packages according to step (a), the parental potential binding domain for the potential binding domains of said new packages being a successful binding domain whose sequence was determined in step (e), and repeating steps (b)-(e) with said new population, and, when a package bearing a binding domain of desired binding characteristics is obtained,
- 10 g) abstracting the DNA encoding the desired binding domain from the genetic package and placing it into a suitable expression system. (The binding domain may then be expressed as a unitary protein, or as a domain of a larger protein).
- 15

The invention is not, however, limited to proteins with a single BD since the method may be applied to any or all of the BDs of the protein, sequentially or simultaneously.

20 The invention is not, however, limited to biological synthesis of the binding domains; peptides having an amino-acid sequence determined by the isolated DNA can be chemically synthesized.

The invention further relates to a variegated population of genetic packages. Said population may be used by one user to select for binding to a first target, by a second user to select for binding to a second target, and so on, as the present invention does not require that the initial potential binding domain actually bind to the target of interest, and the variegation is at residues likely to affect binding. The invention also relates to the variegated DNA used in preparing such genetic packages.

25

30

The invention likewise encompasses the procedure by

which the display strategy is verified. The genetic packages are engineered to display a single IPBD sequence. (Variability may be introduced into DNA subsequences adjacent to the ipbd subsequence and within the osp-ipbd gene so that the IPBD will appear on the GP surface.) A molecule, such as an antibody, having high affinity for correctly folded IPBD is used to: a) detect IPBD on the GP surface, b) screen colonies for display of IPBD on the GP surface, or c) select GPs that display IPBD from a population, some members of which might display IPBD on the GP surface. In one preferred embodiment, this verification process (part I) involves:

- 1) choosing a GP such as a bacterial cell, bacterial spore, or phage, having a suitable outer surface protein (OSP),
- 2) choosing a stable IPBD,
- 3) designing an amino acid sequence that: a) includes the IPBD as a subsequence and b) will cause the IPBD to appear on the GP surface,
- 4) engineering a gene, denoted osp-ipbd, that: a) codes for the designed amino acid sequence, b) provides the necessary genetic regulation, and c) introduces convenient sites for genetic manipulation,
- 5) cloning the osp-ipbd gene into the GP, and
- 6) harvesting the transformed GPs and testing them for presence of IPBD on the GP surface; this test is performed with an affinity molecule having high affinity for IPBD, denoted AfM(IPBD).

Once a GP(IPBD) is produced, it can be used many times as the starting point for developing different novel proteins that bind to a variety of different targets. The knowledge of how we engineer the appearance of one IPBD on the surface of a GP can be used to design and produce other GP(IPBD)s that display different IPBDs.

Knowing that a particular genetic package and osp-
ipbd fusion are suitable for the practice of the inven-
tion, we may variegate the genetic packages and select for
binding to a target of interest. Using IPBD as the PPBD
5 to the first cycle of variegation, we prepare a wide
variety of osp-pbd genes that encode a wide variety of
PBDs. We use an affinity separation to enrich the
population of GP(vgPBD)s for GPs that display PBDs with
binding properties relative to the target that are
10 superior to the binding properties of the PPBD. An SBD
selected from one variegation cycle becomes the PPBD to
the next variegation cycle. In a preferred embodiment,
Part II of the process of the present invention involves:

- 15 1) picking a target molecule, and an affinity separation
system which selects for proteins having an affinity
for that target molecule,
- 2) picking a GP(IPBD),
- 20 3) picking a set of several residues in the PPBD to
vary; the principal indicators of which residues to
vary include: a) the 3D structure of the IPBD, b)
sequences of homologous proteins, and c) computer or
theoretical modeling that indicates which residues
can tolerate different amino acids without disrupting
the underlying structure,
- 25 4) picking a subset of the residues picked in Part II.3,
to be varied simultaneously; the principal considera-
tions are the number of different variants and which
variants are within the detection capabilities of the
affinity separation system, and setting the range of
30 variation;
- 5) implementing the variegation by:
 - a) synthesizing the part of the osp-pbd gene that
encodes the residues to be varied using a
specific mixture of nucleotide substrates for
35 some or all of the bases encoding residues

T 06290 5603660

- slated for variation, thereby creating a population of DNA molecules, denoted vgDNA,
- b) ligating this vgDNA, by standard methods, into the operative cloning vector (OCV) (e.g. a plasmid or bacteriophage),
- c) using the ligated DNA to transform cells, thereby producing a population of transformed cells,
- d) culturing (i.e. increasing in number) the population of transformed cells and harvesting the population of GP(PBD)s, said population being denoted as GP(vgPBD),
- e) enriching the population for GPs that bind the target by using affinity separation, with the chosen target molecule as affinity molecule,
- f) repeating steps II.5.d and II.5.e until a GP(SBD) having improved binding to the target is isolated, and
- g) testing the isolated SBD or SBDs for affinity and specificity for the chosen target,
- 6) repeating steps II.3, II.4, and II.5 until the desired degree of binding is obtained.

Part II is repeated for each new target material. Part I need be repeated only if no GP(IPBD) suitable to a chosen target is available.

For each target, there are a large number of SBDs that may be found by the method of the present invention. The process relies on a combination of protein structural considerations, probabilities, and targeted mutations with accumulation of information. To increase the probability that some PBD in the population will bind to the target, we generate as large a population as we can conveniently subject to selection-through-binding in one experiment. Key questions in management of the method are "How many

transformants can we produce?", and "How small a component can we find through selection-through-binding?". The optimum level of variegation is determined by the maximum number of transformants and the selection sensitivity, so
 5 that for any reasonable sensitivity we may use a progressive process to obtain a series of proteins with higher and higher affinity for the chosen target material.

The appended claims are hereby incorporated by reference into this specification as an enumeration of the
 10 preferred embodiments.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows how a phage may be used as a genetic package. At (a) we have a wild-type precoat protein
 15 lodged in the lipid bilayer. The signal peptide is in the periplasmic space. At (b), a chimeric precoat protein, with a potential binding domain interposed between the signal peptide and the mature coat protein sequence, is similarly trapped. At (c) and
 20 (d), the signal peptide has been cleaved off the wild-type and chimeric proteins, respectively, but certain residues of the coat protein sequence interact with the lipid bilayer to prevent the mature protein from passing entirely into the periplasm. At (e) and (f), mature wild-type and
 25 chimeric protein are assembled into the coat of a single stranded DNA phage as it emerges into the periplasmic space. The phage will pass through the outer membrane into the medium where it can be
 30 recovered and chromatographically evaluated.

Figure 2 depicts (a) the optimal stereochemistry of a disulfide bond, based on Creighton, "Disulfide Bonds and Protein Stability" (CREI88) (the two possible torsion angles about the disulfide bond of $+90^\circ$ and

5

10

15

15

20

25

30

35

Figure 11 shows pH profile for the fractionation of the MYMUT Library on cathepsin G beads. The abscissae shows pH of buffer. The ordinants show amount of phage (as fraction of input phage) obtained at given pH. Ordinants scaled by 10^3 .

Figure 12 shows a second fractionation of MYMUT library over cathepsin G.

Figure 13 shows elution profiles on immobilized cathepsin G for phage selected for binding to cathepsin G.

Figure 14 shows the C_{α} s of BPTI and interaction set #2.

Figure 15 shows the main chain of scorpion toxin (Brookhaven Protein Data Bank entry 1SN3) residues 20 through 42. CYS₂₅ and CYS₄₁ are shown forming a disulfide. In the native protein these groups form disulfides to other cysteines, but no main-chain motion is required to bring the gamma sulphurs into acceptable geometry. Residues, other than GLY, are labeled at the β carbon with the one-letter code.

Figure 16 shows profiles of the elution of phage that display EpiNE7 and EpiNE7.23 from HNE beads.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

OVERVIEW

I. DEFINITIONS AND ABBREVIATIONS

II. THE INITIAL POTENTIAL BINDING DOMAIN

A. Generally

B. Influence of Target Size on Choice of IPBD

C. Influence of Target Charge on Choice of IPBD

D. Other Considerations in the Choice of IPBD

E. Bovine Pancreatic Trypsin Inhibitor (BPTI) as an IPBD

F. Mini-Proteins as IPBDs

G. Modified PBDs

III. VARIEGATION STRATEGY - MUTAGENESIS TO OBTAIN POTENTIAL BINDING DOMAINS WITH DESIRED DIVERSITY

- A. Generally
- B. Identification of Residues to be Varied
- 5 C. Determining the Substitution Set for Each Parental Residue
- D. Special Considerations Relating to Variegation of Mini-Proteins with Essential Cysteines
- E. Planning the Second and Later Rounds of Variegation

10

IV. DISPLAY STRATEGY - DISPLAYING FOREIGN BINDING DOMAINS ON THE SURFACE OF A "GENETIC PACKAGE"

- A. General Requirements for Genetic Package
- B. Phages for Use as Genetic Packages
- 15 C. Bacterial Cells as Genetic Packages
- D. Bacterial Spores as Genetic Packages
- E. Artificial Outer Surface Protein
- F. Designing the osp::ipbd Gene Insert
- G. Synthesis of Gene Inserts
- 20 H. Operative Cloning Vector
- I. Transformation of Cells
- J. Verification of Display Strategy
- K. Analysis and Correction of Display Problems

20

V. AFFINITY SELECTION OF TARGET-BINDING MUTANTS

- 25 A. Affinity Separation Technology, Generally
- B. Affinity Chromatography, Generally
- C. Fluorescent-Activated Cell Sorting, Generally
- D. Affinity Electrophoresis, Generally
- E. Target Materials
- 30 F. Immobilization or Labeling of Target Material
- G. Elution of Lower Affinity PBD-Bearing Packages
- H. Optimization of Affinity Separation
- I. Measuring the Sensitivity of Affinity Separation
- J. Measuring the Efficiency of Separation
- 35 K. Reducing Selection due to Non-Specific Binding

30

35

T06290" 56096860

- L. Isolation of Genetic Package PBDs with Binding-to-Target Phenotypes
- M. Recovery of Packages
- N. Amplifying the Enriched Packages
- 5 O. Determining Whether Further Enrichment is Needed
- P. Characterizing the Putative SBDs
- Q. Joint Selections
- R. Selection for Non-Binding
- S. Selection of Potential Binding Domains for
- 10 Retention of Structure
- T. Engineering of Antagonists

VI. EXPLOITATION OF SUCCESSFUL BINDING DOMAINS AND CORRESPONDING DNAS

- A. Generally
- 15 B. Production of Novel Binding Proteins
- C. Mini-Protein Production
- D. Uses of Novel Binding Proteins

VII. EXAMPLES

20 I. DEFINITIONS AND ABBREVIATIONS

Let $K_d(x,y)$ be a dissociation constant,

$$K_d(x,y) = \frac{[x] [y]}{[x:y]}$$

25

For the purposes of the appended claims, a protein P is a binding protein if (1) For one molecular, ionic or atomic species A, other than the variable domain of an antibody, the dissociation constant $K_D(P,A) < 10^{-6}$ moles/liter (preferably, $< 10^{-7}$ moles/liter), and (2) for a different molecular, ionic or atomic species B, $K_D(P,B) > 10^{-4}$ moles/liter (preferably, $> 10^{-1}$ moles/liter). As a result of these two conditions, the protein P exhibits specificity for A over B, and a minimum degree of affinity (or

35 avidity) for A.

T06250" 56096850

The exclusion of "variable domain of an antibody" in (1) above is intended to make clear that for the purposes herein a protein is not to be considered a "binding protein" merely because it is antigenic. However, an antigen may nonetheless qualify as a binding protein because it specifically binds to a substance other than an antibody, e.g., an enzyme for its substrate, or a hormone for its cellular receptor. Additionally, it should be pointed out that "binding protein" may include a protein which binds specifically to the Fc of an antibody, e.g., staphylococcal protein A.

Normally, the binding protein will not be an antibody or a antigen-binding derivative thereof. An antibody is a crosslinked complex of four polypeptides (two heavy and two light chains). The light chains of IgG have a molecular weight of $\approx 23,000$ daltons and the heavy chains of $\approx 53,000$ daltons. A single binding unit is composed of the variable region of a heavy chain (V_H) and the variable region of a light chain (V_L), each about 110 amino-acid residues. The V_H and V_L regions are held in proximity by a disulfide bond between the adjoining C_L and C_{H1} regions; altogether, these total 440 residues and correspond to an Fab fragment. Derivatives of antibodies include Fab fragments and the individual variable light and heavy domains. A special case of antibody derivative is a "single chain antibody." A "single-chain antibody" is a single chain polypeptide comprising at least 200 amino acids, said amino acids forming two antigen-binding regions connected by a peptide linker that allows the two regions to fold together to bind the antigen in a manner akin to that of an Fab fragment. Either the two antigen-binding regions must be variable domains of known antibodies, or they must (1) each fold into a β barrel of nine strands that are spatially related in the same way as are the nine strands of known antibody variable light

FOI 290 5609550

or heavy domains, and (2) fit together in the same way as do the variable domains of said known antibody. Generally speaking, this will require that, with the exception of the amino acids corresponding to the hypervariable region, there is at least 88% homology with the amino acids of the variable domain of a known antibody.

While the present invention may be used to develop novel antibodies through variegation of codons corresponding to the hypervariable region of an antibody's variable domain, its primary utility resides in the development of binding proteins which are not antibodies or even variable domains of antibodies. Novel antibodies can be obtained by immunological techniques; novel enzymes, hormones, etc. cannot.

It will be appreciated that, as a result of evolution, the antigen-binding domains of antibodies have acquired a structure which tolerates great variability of sequence in the hypervariable regions. The remainder of the variable domain is made up of constant regions forming a distinctive structure, a nine strand β barrel, which hold the hypervariable regions (inter-strand loops) in a fixed relationship with each other. Most other binding proteins lack this molecular design which facilitates diversification of binding characteristics. Consequently, the successful development of novel antibodies by modification of sequences encoding known hypervariable regions--which, in nature, vary from antibody to antibody--does not provide any guidance or assurance of success in the development of novel, non-immunoglobulin binding proteins.

It should further be noted that the affinity of antibodies for their target epitopes is typically on the order of 10^6 to 10^{10} liters/mole; many enzymes exhibit much greater affinities (10^9 to 10^{15} liters/mole) for their preferred substrates. Thus, if the goal is to

develop a binding protein with a very high affinity for a target of interest, e.g., greater than 10^{10} , the antibody design may in fact be unduly limiting. Furthermore, the complementarity-determining residues of an antibody comprises many residues, 30 to 50. In most cases, it is not known which of these residues participates directly in binding antigen. Thus, picking an antibody as PPBD does not allow us to focus variegation to a small number of residues.

Most larger proteins fold into distinguishable globules called domains (ROSS81). Protein domains have been defined various ways, but all definitions fall into one of three classes: a) those that define a domain in terms of 3D atomic coordinates, b) those that define a domain as an isolable, stable fragment of a larger protein, and c) those that define a domain based on protein sequence homology plus a method from class a) or b). Frequently, different methods of defining domains applied to a single protein yield identical or very similar domain boundaries. The diversity of definitions for domains stems from the many ways that protein domains are perceived to be important, including the concept of domains in predicting the boundaries of stable fragments, and the relationship of domains to protein folding, function, stability, and evolution. The present invention emphasizes the retention of the structured character of a domain even though its surface residues are mutated. Consequently, definitions of "domain" which emphasize stability -- retention of the overall structure in the face of perturbing forces such as elevated temperatures or chaotropic agents -- are favored, though atomic coordinates and protein sequence homology are not completely ignored.

When a domain of a protein is primarily responsible

106290-55095850

for the protein's ability to specifically bind a chosen target, it is referred to herein as a "binding domain" (BD). A preliminary operation is to engineer the appearance of a stable protein domain, denoted as an "initial potential binding domain" (IPBD), on the surface of a genetic package.

The term "variegated DNA" (vgDNA) refers to a mixture of DNA molecules of the same or similar length which, when aligned, vary at some codons so as to encode at each such codon a plurality of different amino acids, but which encode only a single amino acid at other codon positions. It is further understood that in variegated DNA, the codons which are variable, and the range and frequency of occurrence of the different amino acids which a given variable codon encodes, are determined in advance by the synthesizer of the DNA, even though the synthetic method does not allow one to know, a priori, the sequence of any individual DNA molecule in the mixture. The number of designated variable codons in the variegated DNA is preferably no more than 20 codons, and more preferably no more than 5-10 codons. The mix of amino acids encoded at each variable codon may differ from codon to codon.

A population of genetic packages into which variegated DNA has been introduced is likewise said to be "variegated".

For the purposes of this invention, the term "potential binding protein" refers to a protein encoded by one species of DNA molecule in a population of variegated DNA wherein the region of variation appears in one or more subsequences encoding one or more segments of the polypeptide having the potential of serving as a binding domain for the target substance.

From time to time, it may be helpful to speak of the "parent sequence" of the variegated DNA. When the novel

binding domain sought is an analogue of a known binding domain, the parent sequence is the sequence that encodes the known binding domain. The variegated DNA will be identical with this parent sequence at one or more loci, but will diverge from it at chosen loci. When a potential binding domain is designed from first principles, the parent sequence is a sequence which encodes the amino acid sequence that has been predicted to form the desired binding domain, and the variegated DNA is a population of "daughter DNAs" that are related to that parent by a recognizable sequence similarity.

A "chimeric protein" is a protein composed of a first amino acid sequence substantially corresponding to the sequence of a protein or to a large fragment of a protein (20 or more residues) expressed by the species in which the chimeric protein is expressed and a second amino acid sequence that does not substantially correspond to an amino acid sequence of a protein expressed by the first species but that does substantially correspond to the sequence of a protein expressed by a second and different species of organism. The second sequence is said to be foreign to the first sequence.

One amino acid sequence of the chimeric proteins of the present invention is typically derived from an outer surface protein of a "genetic package" as hereafter defined. The second amino acid sequence is one which, if expressed alone, would have the characteristics of a protein (or a domain thereof) but is incorporated into the chimeric protein as a recognizable domain thereof. It may appear at the amino or carboxy terminal of the first amino acid sequence (with or without an intervening spacer), or it may interrupt the first amino acid sequence. The first amino acid sequence may correspond exactly to a surface protein of the genetic package, or it may be modified,

e.g., to facilitate the display of the binding domain.

In the present invention, the words "select" and "selection" are used in the genetic sense; i.e. a biological process whereby a phenotypic characteristic is used
5 to enrich a population for those organisms displaying the desired phenotype.

One affinity separation is called a "separation cycle"; one pass of variegation followed by as many separation cycles as are needed to isolate an SBD, is
10 called a "variegation cycle". The amino acid sequence of one SBD from one round becomes the PPBD to the next variegation cycle. We perform variegation cycles iteratively until the desired affinity and specificity of binding between an SBD and chosen target are achieved.

15 The following abbreviations will be used throughout the present specification:

Abbreviation	Meaning
GP	Genetic Package, <u>e.g.</u> a bacteriophage
20 wtGP	Wild-type GP
X	Any protein
<u>X</u>	The gene for protein X
BD	Binding Domain
BPTI	Bovine pancreatic trypsin inhibitor, identical to aprotinin (Merck Index, entry 784, p.119)
25 IPBD	Initial Potential Binding Domain, <u>e.g.</u> BPTI
30 PBD	Potential Binding Domain, <u>e.g.</u> a derivative of BPTI
SBD	Successful Binding Domain, <u>e.g.</u> a derivative of BPTI selected for binding to a target

	PPBD	Parental Potential Binding Domain, <u>i.e.</u> an IPBD or an SBD from a previous selection
5	OSP	Outer Surface Protein, <u>e.g.</u> coat protein of a phage or LamB from <u>E. coli</u>
	OSP-PBD	Fusion of an OSP and a PBD, order of fusion not specified
	OSTS	Outer Surface Transport Signal
10	GP(<u>x</u>)	A genetic package containing the <u>x</u> gene
	GP(X)	A genetic package that displays X on its outer surface
	GP(<u>osp-pbd</u>)	GP containing an <u>osp-pbd</u> gene
15	GP(OSP-PBD)	A genetic package that displays PBD on its outside as a fusion to OSP
	GP(<u>pbd</u>)	GP containing a <u>pbd</u> gene, <u>osp</u> implicit
20	GP(PBD)	A genetic package displaying PBD on its outside, OSP unspecified
	{Q}	An affinity matrix supporting "Q", <u>e.g.</u> {T4 lysozyme} is T4 lysozyme attached to an affinity matrix
25	AfM(W)	A molecule having affinity for "W", <u>e.g.</u> trypsin is an AfM(BPTI)
30	AfM(W)*	AfM(W) carrying a label, <u>e.g.</u> ¹²⁵ I
	XINDUCE	A chemical that can induce expression of a gene, <u>e.g.</u> IPTG for the <u>lacUV5</u> promoter
35	OCV	Operative Cloning Vector

706230" 55095850

	K_d	A bimolecular dissociation constant, $K_d = [A][B]/[A:B]$
	K_T	$K_T = [T][SBD]/[T:SBD]$ (T is a target)
5	K_N	$K_N = [N][SBD]/[N:SBD]$ (N is a non-target)
	DoAMoM	Density of AfM(W) on affinity matrix
	mfaa	Most-Favored amino acid
10	lfaa	Least-Favored amino acid
	Abun(x)	Abundance of DNA molecules encoding amino acid x
	OMP	Outer membrane protein
	nt	nucleotide
15	SP-I	Signal-sequence Peptidase I
	Y_{DQ}	Yield of ssDNA up to Q bases long
	M_{ssDNA}	Maximum length of ssDNA that can be synthesized in acceptable yield
20	Y_{pl}	Yield of plasmid DNA per volume of culture
	L_{eff}	DNA ligation efficiency
	M_{ntv}	Maximum number of transformants produced from Y_{D100} DNA of Insert
25	C_{eff}	Efficiency of chromatographic enrichment, enrichment per pass
	C_{sensi}	Sensitivity of chromatographic separation, can find 1 in N,
30	N_{chrom}	Maximum number of enrichment cycles per variegation cycle
	S_{err}	Error level in synthesizing vgDNA
35	::	in-frame genetic fusion or

protein produced from in-frame
fused gene

Single-letter codes for amino acids and nucleotides are
5 given in Table 1.

--- *** ---

II. THE INITIAL POTENTIAL BINDING DOMAIN (IPBD):

II.A. Generally

10 The initial potential binding domain may be: 1) a
domain of a naturally occurring protein, 2) a non-natur-
ally occurring domain which substantially corresponds in
sequence to a naturally occurring domain, but which
differs from it in sequence by one or more substitutions,
15 insertions or deletions, 3) a domain substantially
corresponding in sequence to a hybrid of subsequences of
two or more naturally occurring proteins, or 4) an artifi-
cial domain designed entirely on theoretical grounds
based on knowledge of amino acid geometries and statis-
20 tical evidence of secondary structure preferences of
amino acids. (However, the limitations of a priori
protein design prompted the present invention.) Usually,
the domain will be a known binding domain, or at least a
homologue thereof, but it may be derived from a protein
25 which, while not possessing a known binding activity,
possesses a secondary or higher structure that lends
itself to binding activity (clefts, grooves, etc.). The
protein to which the IPBD is related need not have any
specific affinity for the target material.

30 In determining whether sequences should be deemed to
"substantially correspond", one should consider the
following issues: the degree of sequence similarity when
the sequences are aligned for best fit according to
standard algorithms, the similarity in the connectivity

patterns of any crosslinks (e.g., disulfide bonds), the degree to which the proteins have similar three-dimensional structures, as indicated by, e.g., X-ray diffraction analysis or NMR, and the degree to which the sequenced proteins have similar biological activity. In this context, it should be noted that among the serine protease inhibitors, there are families of proteins recognized to be homologous in which there are pairs of members with as little as 30% sequence homology.

10 A candidate IPBD should meet the following criteria:

- 1) a domain exists that will remain stable under the conditions of its intended use (the domain may comprise the entire protein that will be inserted, e.g. BPTI, α -conotoxin GI, or CMTI-III),
- 15 2) knowledge of the amino acid sequence is obtainable, and
- 3) a molecule is obtainable having specific and high affinity for the IPBD, AfM(IPBD).

Preferably, in order to guide the variegation strategy, 20 knowledge of the identity of the residues on the domain's outer surface, and their spatial relationships, is obtainable; however, this consideration is less important if the binding domain is small, e.g., under 40 residues.

Preferably, the IPBD is no larger than necessary 25 because small SBDs (for example, less than 30 amino acids) can be chemically synthesized and because it is easier to arrange restriction sites in smaller amino-acid sequences. For PBDs smaller than about 40 residues, an added advantage is that the entire variegated pbd gene can be 30 synthesized in one piece. In that case, we need arrange only suitable restriction sites in the osp gene. A smaller protein minimizes the metabolic strain on the GP or the host of the GP. The IPBD is preferably smaller than about 200 residues. The IPBD must also be large

T06230" 56056857

enough to have acceptable binding affinity and specificity. For an IPBD lacking covalent crosslinks, such as disulfide bonds, the IPBD is preferably at least 40 residues; it may be as small as six residues if it
 5 contains a crosslink. These small, crosslinked IPBDs, known as "mini-proteins", are discussed in more detail later in this section.

Some candidate IPBDs, which meet the conditions set forth above, will be more suitable than others. Informa-
 10 tion about candidate IPBDs that will be used to judge the suitability of the IPBD includes: 1) a 3D structure (knowledge strongly preferred), 2) one or more sequences homologous to the IPBD (the more homologous sequences known, the better), 3) the pI of the IPBD (knowledge
 15 desirable when target is highly charged), 4) the stability and solubility as a function of temperature, pH and ionic strength (preferably known to be stable over a wide range and soluble in conditions of intended use), 5) ability to bind metal ions such as Ca^{++} or Mg^{++} (knowledge preferred;
 20 binding per se, no preference), 6) enzymatic activities, if any (knowledge preferred, activity per se has uses but may cause problems), 7) binding properties, if any (knowledge preferred, specific binding also preferred), 8) availability of a molecule having specific and strong
 25 affinity ($K_d < 10^{-11}$ M) for the IPBD (preferred), 9) availability of a molecule having specific and medium affinity (10^{-8} M $< K_d < 10^{-6}$ M) for the IPBD (preferred), 10) the sequence of a mutant of IPBD that does not bind to the affinity molecule(s) (preferred), and 11) absorption
 30 spectrum in visible, UV, NMR, etc. (characteristic absorption preferred).

If only one species of molecule having affinity for IPBD (AfM(IPBD)) is available, it will be used to: a) detect the IPBD on the GP surface, b) optimize expression

105-250-56096350

level and density of the affinity molecule on the matrix, and c) determine the efficiency and sensitivity of the affinity separation. As noted above, however, one would prefer to have available two species of AfM(IPBD), one with high and one with moderate affinity for the IPBD. The species with high affinity would be used in initial detection and in determining efficiency and sensitivity, and the species with moderate affinity would be used in optimization.

10 If the IPBD is not itself a binding domain of a known binding protein, or if its native target has not been purified, an antibody raised against the IPBD may be used as the affinity molecule. Use of an antibody for this purpose should not be taken to mean that the antibody is the ultimate target.

There are many candidate IPBDs for which all of the above information is available or is reasonably practical to obtain, for example, bovine pancreatic trypsin inhibitor (BPTI, 58 residues), CMTI-III (29 residues), crambin (46 residues), third domain of ovomucoid (56 residues), heat-stable enterotoxin (ST-Ia of *E. coli*) (18 residues), α -Conotoxin GI (13 residues), μ -Conotoxin GIII (22 residues), Conus King Kong mini-protein (27 residues), T4 lysozyme (164 residues), and azurin (128 residues). Structural information can be obtained from X-ray or neutron diffraction studies, NMR, chemical cross linking or labeling, modeling from known structures of related proteins, or from theoretical calculations. 3D structural information obtained by X-ray diffraction, neutron diffraction or NMR is preferred because these methods allow localization of almost all of the atoms to within defined limits. Table 50 lists several preferred IPBDs. Works related to determination of 3D structure of small proteins via NMR include: CHAZ85, PEAS90, PEAS88, CLOR86,

CLOR87a, HEIT89, LECO87, WAGN79, and PARD89.

In some cases, a protein having some affinity for the target may be a preferred IPBD even though some other criteria are not optimally met. For example, the V1 domain of CD4 is a good choice as IPBD for a protein that binds to gp120 of HIV. It is known that mutations in the region 42 to 55 of V1 greatly affect gp120 binding and that other mutations either have much less effect or completely disrupt the structure of V1. Similarly, tumor necrosis factor (TNF) would be a good initial choice if one wants a TNF-like molecule having higher affinity for the TNF receptor.

Membrane-bound proteins are not preferred IPBDs, though they may serve as a source of outer surface transport signals. One should distinguish between membrane-bound proteins, such as LamB or OmpF, that cross the membrane several times forming a structure that is embedded in the lipid bilayer and in which the exposed regions are the loops that join trans-membrane segments, from non-embedded proteins, such as the soluble domains of CD4, that are simply anchored to the membrane. This is an important distinction because it is quite difficult to create a soluble derivative of a membrane-bound protein. Soluble binding proteins are in general more useful since purification is simpler and they are more tractable and more versatile assay reagents.

Most of the PBDs derived from a PPBD according to the process of the present invention will have been derived by variegation at residues having side groups directed toward the solvent. Reidhaar-Olson and Sauer (REID88a) found that exposed residues can accept a wide range of amino acids, while buried residues are more limited in this regard. Surface mutations typically have only small effects on melting temperature of the PBD, but may reduce

Two general characteristics of the target molecule, size and charge, make certain classes of IPBDs more likely than other classes to yield derivatives that will bind specifically to the target. Because these are very general characteristics, one can divide all targets into six classes: a) large positive, b) large neutral, c) large negative, d) small positive, e) small neutral, and f) small negative. A small collection of IPBDs, one or a few corresponding to each class of target, will contain a preferred candidate IPBD for any chosen target.

II.B. Influence of target size on choice of IPBD:

If the target is a protein or other macromolecule a preferred embodiment of the IPBD is a small protein such as the *Cucurbita maxima* trypsin inhibitor III (29 resi-

dues), BPTI from Bos Taurus (58 residues), crambin from rape seed (46 residues), or the third domain of ovomucoid from Coturnix coturnix Japonica (Japanese quail) (56 residues), because targets from this class have clefts and grooves that can accommodate small proteins in highly specific ways. If the target is a macromolecule lacking a compact structure, such as starch, it should be treated as if it were a small molecule. Extended macromolecules with defined 3D structure, such as collagen, should be treated as large molecules.

If the target is a small molecule, such as a steroid, a preferred embodiment of the IPBD is a protein of about 80-200 residues, such as ribonuclease from Bos taurus (124 residues), ribonuclease from Aspergillus oryzae (104 residues), hen egg white lysozyme from Gallus gallus (129 residues), azurin from Pseudomonas aeruginosa (128 residues), or T4 lysozyme (164 residues), because such proteins have clefts and grooves into which the small target molecules can fit. The Brookhaven Protein Data Bank contains 3D structures for all of the proteins listed. Genes encoding proteins as large as T4 lysozyme can be manipulated by standard techniques for the purposes of this invention.

If the target is a mineral, insoluble in water, one considers the nature of the molecular surface of the mineral. Minerals that have smooth surfaces, such as crystalline silicon, are best addressed with medium to large proteins, such as ribonuclease, as IPBD in order to have sufficient contact area and specificity. Minerals with rough, grooved surfaces, such as zeolites, could be bound either by small proteins, such as BPTI, or larger proteins, such as T4 lysozyme.

II.C. Influence of target charge on choice of IPBD:

Electrostatic repulsion between molecules of like

FOIA b 7 - D

charge can prevent molecules with highly complementary surfaces from binding. Therefore, it is preferred that, under the conditions of intended use, the IPBD and the target molecule either have opposite charge or that one of them is neutral. In some cases it has been observed that protein molecules bind in such a way that like charged groups are juxtaposed by including oppositely charged counter ions in the molecular interface. Thus, inclusion of counter ions can reduce or eliminate electrostatic repulsion and the user may elect to include ions in the eluants used in the affinity separation step. Polyvalent ions are more effective at reducing repulsion than monovalent ions.

II.D. Other considerations in the choice of IPBD:

If the chosen IPBD is an enzyme, it may be necessary to change one or more residues in the active site to inactivate enzyme function. For example, if the IPBD were T4 lysozyme and the GP were E. coli cells or M13, we would need to inactivate the lysozyme because otherwise it would lyse the cells. If, on the other hand, the GP were ϕ X174, then inactivation of lysozyme may not be needed because T4 lysozyme can be overproduced inside E. coli cells without detrimental effects and ϕ X174 forms intracellularly. It is preferred to inactivate enzyme IPBDs that might be harmful to the GP or its host by substituting mutant amino acids at one or more residues of the active site. It is permitted to vary one or more of the residues that were changed to abolish the original enzymatic activity of the IPBD. Those GPs that receive osp-pbd genes encoding an active enzyme may die, but the majority of sequences will not be deleterious.

If the binding protein is intended for therapeutic use in humans or animals, the IPBD may be chosen from proteins native to the designated recipient to minimize

105230"55095567

the possibility of antigenic reactions.

II.E. Bovine Pancreatic Trypsin Inhibitor (BPTI) as an IPBD:

BPTI is an especially preferred IPBD because it meets
 5 or exceeds all the criteria: it is a small, very stable
 protein with a well known 3D structure. Marks et al.
 (MARK86) have shown that a fusion of the phoA signal
 peptide gene fragment and DNA coding for the mature form
 of BPTI caused native BPTI to appear in the periplasm of
 10 E. coli, demonstrating that there is nothing in the
 structure of BPTI to prevent its being secreted.

The structure of BPTI is maintained even when one or
 another of the disulfides is removed, either by chemical
 blocking or by genetic alteration of the amino-acid
 15 sequence. The stabilizing influence of the disulfides in
 BPTI is not equally distributed. Goldenberg (GOLD85)
 reports that blocking CYS14 and CYS38 lowers the T_m of
 BPTI to $\approx 75^\circ\text{C}$ while chemical blocking of either of the
 other disulfides lowers T_m to below 40°C . Chemically
 20 blocking a disulfide may lower T_m more than mutating the
 cysteines to other amino-acid types because the bulky
 blocking groups are more destabilizing than removal of the
 disulfide. Marks et al. (MARK87) replaced both CYS14 and
 CYS38 with either two alanines or two threonines. The
 25 CYS14/CYS38 cystine bridge that Marks et al. removed is
 the one very close to the scissile bond in BPTI; surpris-
 ingly, both mutant molecules functioned as trypsin
 inhibitors. Schnabel et al. (SCHN86) report preparation
 of aprotinin(C14A,C38A) by use of Raney nickel. Eigenbrot
 30 et al. (EIGE90) report the X-ray structure of BPTI(C30A/-
 C51A) which is stable to at least 50°C . The backbone of
 this mutant is as similar to BPTI as are the backbones of
 BPTI molecules that sit in different crystal lattices.
 This indicates that BPTI is redundantly stable and so is

T05290-56096850

likely to fold into approximately the same structure despite numerous surface mutations. Using the knowledge of homologues, vide infra, we can infer which residues should not be varied if the basic BPTI structure is to be maintained.

The 3D structure of BPTI has been determined at high resolution by X-ray diffraction (HUBE77, MARQ83, WLOD84, WLOD87a, WLOD87b), neutron diffraction (WLOD84), and by NMR (WAGN87). In one of the X-ray structures deposited in the Brookhaven Protein Data Bank, entry 6PTI, there was no electron density for A58, indicating that A58 has no uniquely defined conformation. Thus we know that the carboxy group does not make any essential interaction in the folded structure. The amino terminus of BPTI is very near to the carboxy terminus. Goldenberg and Creighton reported on circularized BPTI and circularly permuted BPTI (GOLD83). Some proteins homologous to BPTI have more or fewer residues at either terminus.

BPTI has been called "the hydrogen atom of protein folding" and has been the subject of numerous experimental and theoretical studies (STAT87, SCHW87, GOLD83, CHAZ83, CREI74, CREI77a, CREI77b, CREI80, SIEK87, SINH90, RUEH73, HUBE74, HUBE75, HUBE77 and others).

BPTI has the added advantage that at least 59 homologous proteins are known. Table 13 shows the sequences of 39 homologues. A tally of ionizable groups in 59 homologues is shown in Table 14 and the composite of amino acid types occurring at each residue is shown in Table 15.

BPTI is freely soluble and is not known to bind metal ions. BPTI has no known enzymatic activity. BPTI is not toxic.

All of the conserved residues are buried; of the six

FO6290" 56095850

fully conserved residues only G37 has noticeable exposure. The solvent accessibility of each residue in BPTI is given in Table 16 which was calculated from the entry "6PTI" in the Brookhaven Protein Data Bank with a solvent radius of 1.4 Å, the atomic radii given in Table 7, and the method of Lee and Richards (LEEB71). Each of the 52 non-conserved residues can accommodate two or more kinds of amino acids. By independently substituting at each residue only those amino acids already observed at that residue, we could obtain approximately $1.6 \cdot 10^{43}$ different amino acid sequences, most of which will fold into structures very similar to BPTI.

BPTI will be especially useful as a IPBD for macromolecular targets. BPTI and BPTI homologues bind tightly and with high specificity to a number of enzyme macromolecules.

BPTI is strongly positively charged except at very high pH, thus BPTI is useful as IPBD for targets that are not also strongly positive under the conditions of intended use. There exist homologues of BPTI, however, having quite different charges (viz. SCI-III from Bombyx mori at -7 and the trypsin inhibitor from bovine colostrum at -1). Once a genetic package is found that displays BPTI on its surface, the sequence of the BPTI domain can be replaced by one of the homologous sequences to produce acidic or neutral IPBDs.

BPTI is quite small; if this should cause a pharmacological problem, two or more BPTI-derived domains may be joined as in humans BPTI homologues, one of which has two domains (BALD85, ALBR83b) and another has three (WUNT88).

Another possible pharmacological problem is immunogenicity. BPTI has been used in humans with very few adverse effects. Siekmann et al. (SIEK89) have studied immunological characteristics of BPTI and some homologues.

T06290" 569980

It is an advantage of the method of the present invention that a variety of SBDs can be obtained so that, if one derivative proves to be antigenic, a different SBD may be used. Furthermore, one can reduce the probability of
 5 immune response by starting with a human protein, such as LACI (a BPTI homologue) (WUNT88, GIRA89) or Inter- α -Trypsin Inhibitor (ALBR83a, ALBR83b, DIAR90, ENGH89, TRIB86, GEBH86, GEBH90, KAUM86, ODOM90, SALI90).

Further, a BPTI-derived gene fragment, coding for a
 10 novel binding domain, could be fused in-frame to a gene fragment coding for other proteins, such as serum albumin or the constant parts of IgG.

Tschesche et al. (TSCH87) reported on the binding of several BPTI derivatives to various proteases:

15 Dissociation constants for BPTI derivatives, Molar.

Residue #15	Trypsin (bovine pancreas)	Chymotrypsin (bovine pancreas)	Elastase (porcine pancreas)	Elastase (human leukocytes)
lysine	$6.0 \cdot 10^{-14}$	$9.0 \cdot 10^{-9}$	-	$3.5 \cdot 10^{-6}$
20 glycine	-	-	+	$7.0 \cdot 10^{-9}$
alanine	+	-	$2.8 \cdot 10^{-8}$	$2.5 \cdot 10^{-9}$
valine	-	-	$5.7 \cdot 10^{-8}$	$1.1 \cdot 10^{-10}$
leucine	-	-	$1.9 \cdot 10^{-8}$	$2.9 \cdot 10^{-9}$

25 From the report of Tschesche et al. we infer that molecular pairs marked "+" have $K_d \geq 3.5 \cdot 10^{-6}$ M and that molecular pairs marked "-" have $K_d \gg 3.5 \cdot 10^{-6}$ M. Because of the wealth of data about the binding of BPTI and various mutants to trypsin and other proteases
 30 (TSCH87), we can proceed in various ways in optimizing the affinity separation conditions. (For other PBDs, we can obtain two different monoclonal antibodies, one with a high affinity having K_d of order 10^{-11} M, and one with a moderate affinity having K_d on the order of 10^{-6} M.)

35 Works concerning BPTI and its homologues include:

KIDO88, PONT88, KIDO90, AUER87, AUER90, SCOT87b, AUER88, AUER89, BECK88b, WACH79, WACH80, BECK89a, DUFT85, FIOR88, GIRA89, GOLD84, GOLD88, HOCH84, RITO83, NORR89a, NORR89b, OLTE89, SWAI88, and WAGN79.

5 II.F Mini-Proteins as IPBDs:

A polypeptide is a polymer composed of a single chain of the same or different amino acids joined by peptide bonds. Linear peptides can take up a very large number of different conformations through internal rotations about
10 the main chain single bonds of each α carbon. These rotations are hindered to varying degrees by side groups, with glycine interfering the least, and valine, isoleucine and, especially, proline, the most. A polypeptide of 20 residues may have 10^{20} different conformations which it
15 may assume by various internal rotations.

Proteins are polypeptides which, as a result of stabilizing interactions between amino acids that are not in adjacent positions in the chain, have folded into a well-defined conformation. This folding is usually
20 essential to their biological activity.

For polypeptides of 40-60 residues or longer, noncovalent forces such as hydrogen bonds, salt bridges, and hydrophobic "interactions" are sufficient to stabilize a particular folding or conformation. The polypeptide's
25 constituent segments are held to more or less that conformation unless it is perturbed by a denaturant such as rising temperature or decreasing pH, whereupon the polypeptide unfolds or "melts". The smaller the peptide, the more likely it is that its conformation will be
30 determined by the environment. If a small unconstrained peptide has biological activity, the peptide ligand will be in essence a random coil until it comes into proximity with its receptor. The receptor accepts the peptide only in one or a few conformations because alternative confor-

FOUO 50096860

mations are disfavored by unfavorable van der Waals and other non-covalent interactions.

Small polypeptides have potential advantages over larger polypeptides when used as therapeutic or diagnostic agents, including (but not limited to):

- a) better penetration into tissues,
- b) faster elimination from the circulation (important for imaging agents),
- c) lower antigenicity, and
- 10 d) higher activity per mass.

Moreover, polypeptides of under about 50 residues have the advantage of accessibility via chemical synthesis; polypeptides of under about 30 residues are more easily synthesized than are larger polypeptides. Thus, it would be desirable to be able to employ the combination of variegation and affinity selection to identify small polypeptides which bind a target of choice.

Polypeptides of this size, however, have disadvantages as binding molecules. According to Olivera et al. (OLIV90a): "Peptides in this size range normally equilibrate among many conformations (in order to have a fixed conformation, proteins generally have to be much larger)." Specific binding of a peptide to a target molecule requires the peptide to take up one conformation that is complementary to the binding site. For a decapeptide with three isoenergetic conformations (e.g., β strand, α helix, and reverse turn) at each residue, there are about $6 \cdot 10^4$ possible overall conformations. Assuming these conformations to be equi-probable for the unconstrained decapeptide, if only one of the possible conformations bound to the binding site, then the affinity of the peptide for the target would be expected to be about $6 \cdot 10^4$ higher if it could be constrained to that single effective conformation. Thus, the unconstrained decapeptide, relative to a

decapeptide constrained to the correct conformation, would be expected to exhibit lower affinity. It would also exhibit lower specificity, since one of the other conformations of the unconstrained decapeptide might be one which bound tightly to a material other than the intended target. By way of corollary, it could have less resistance to degradation by proteases, since it would be more likely to provide a binding site for the protease.

In one embodiment, the present invention overcomes these problems, while retaining the advantages of smaller polypeptides, by fostering the biosynthesis of novel mini-proteins having the desired binding characteristics. Mini-Proteins are small polypeptides (usually less than about 60 residues) which, while too small to have a stable conformation as a result of noncovalent forces alone, are covalently crosslinked (e.g., by disulfide bonds) into a stable conformation and hence have biological activities more typical of larger protein molecules than of unconstrained polypeptides of comparable size.

When mini-proteins are variegated, the residues which are covalently crosslinked in the parental molecule are left unchanged, thereby stabilizing the conformation. For example, in the variegation of a disulfide bonded mini-protein, certain cysteines are invariant so that under the conditions of expression and display, covalent crosslinks (e.g., disulfide bonds between one or more pairs of cysteines) form, and substantially constrain the conformation which may be adopted by the hypervariable linearly intermediate amino acids. In other words, a constraining scaffolding is engineered into polypeptides which are otherwise extensively randomized.

Once a mini-protein of desired binding characteristics is characterized, it may be produced, not only by recombinant DNA techniques, but also by nonbiological

synthetic methods.

In vitro, disulfide bridges can form spontaneously in polypeptides as a result of air oxidation. Matters are more complicated in vivo. Very few intracellular proteins
 5 have disulfide bridges, probably because a strong reducing environment is maintained by the glutathione system. Disulfide bridges are common in proteins that travel or operate in intracellular spaces, such as snake venoms and other toxins (e.g., conotoxins, charybdotoxin, bacterial
 10 enterotoxins), peptide hormones, digestive enzymes, complement proteins, immunoglobulins, lysozymes, protease inhibitors (BPTI and its homologues, CMTI-III (Cucurbita maxima trypsin inhibitor III) and its homologues, hirudin, etc.) and milk proteins.

15 Disulfide bonds that close tight intrachain loops have been found in pepsin, thioredoxin, insulin A-chain, silk fibroin, and lipcamide dehydrogenase. The bridged cysteine residues are separated by one to four residues along the polypeptide chain. Model building, X-ray
 20 diffraction analysis, and NMR studies have shown that the α carbon path of such loops is usually flat and rigid.

There are two types of disulfide bridges in immunoglobulins. One is the conserved intrachain bridge, spanning about 60 to 70 amino acid residues and found,
 25 repeatedly, in almost every immunoglobulin domain. Buried deep between the opposing β sheets, these bridges are shielded from solvent and ordinarily can be reduced only in the presence of denaturing agents. The remaining disulfide bridges are mainly interchain bonds and are
 30 located on the surface of the molecule; they are accessible to solvent and relatively easily reduced (STEI85). The disulfide bridges of the mini-proteins of the present invention are intrachain linkages between cysteines having much smaller chain spacings.

FOIA b 7 - D

For the purpose of the appended claims, a mini-protein has between about eight and about sixty residues. However, it will be understood that a chimeric surface protein presenting a mini-protein as a domain will normally have more than sixty residues. Polypeptides containing intrachain disulfide bonds may be characterized as cyclic in nature, since a closed circle of covalently bonded atoms is defined by the two cysteines, the intermediate amino acid residues, their peptidyl bonds, and the disulfide bond. The terms "cycle", "span" and "segment" will be used to define certain structural features of the polypeptides. An intrachain disulfide bridge connecting amino acids 3 and 8 of a 16 residue polypeptide will be said herein to have a cycle of 6 and a span of 4. If amino acids 4 and 12 are also disulfide bonded, then they form a second cycle of 9 with a span of 7. Together, the four cysteines divide the polypeptide into four inter-cysteine segments (1-2, 5-7, 9-11, and 13-16). (Note that there is no segment between Cys3 and Cys4.)

The connectivity pattern of a crosslinked mini-protein is a simple description of the relative location of the termini of the crosslinks. For example, for a mini-protein with two disulfide bonds, the connectivity pattern "1-3, 2-4" means that the first crosslinked cysteine is disulfide bonded to the third crosslinked cysteine (in the primary sequence), and the second to the fourth.

The degree to which the crosslink constrains the conformational freedom of the mini-protein, and the degree to which it stabilizes the mini-protein, may be assessed by a number of means. These include absorption spectroscopy (which can reveal whether an amino acid is buried or exposed), circular dichroism studies (which provides a general picture of the helical content of the protein),

nuclear magnetic resonance imaging (which reveals the number of nuclei in a particular chemical environment as well as the mobility of nuclei), and X-ray or neutron diffraction analysis of protein crystals. The stability of the mini-protein may be ascertained by monitoring the changes in absorption at various wavelengths as a function of temperature, pH, etc.; buried residues become exposed as the protein unfolds. Similarly, the unfolding of the mini-protein as a result of denaturing conditions results in changes in NMR line positions and widths. Circular dichroism (CD) spectra are extremely sensitive to conformation.

The variegated disulfide-bonded mini-proteins of the present invention fall into several classes.

Class I mini-proteins are those featuring a single pair of cysteines capable of interacting to form a disulfide bond, said bond having a span of no more than nine residues. This disulfide bridge preferably has a span of at least two residues; this is a function of the geometry of the disulfide bond. When the spacing is two or three residues, one residue is preferably glycine in order to reduce the strain on the bridged residues. The upper limit on spacing is less precise, however, in general, the greater the spacing, the less the constraint on conformation imposed on the linearly intermediate amino acid residues by the disulfide bond.

The main chain of such a peptide has very little freedom, but is not stressed. The free energy released when the disulfide forms exceeds the free energy lost by the main-chain when locked into a conformation that brings the cysteines together. Having lost the free energy of disulfide formation, the proximal ends of the side groups are held in more or less fixed relation to each other. When binding to a target, the domain does not need to


TOP SECRET 56095850


expend free energy getting into the correct conformation. The domain can not jump into some other conformation and bind a non-target.

A disulfide bridge with a span of 4 or 5 is especially preferred. If the span is increased to 6, the constraining influence is reduced. In this case, we prefer that at least one of the enclosed residues be an amino acid that imposes restrictions on the main-chain geometry. Proline imposes the most restriction. Valine and isoleucine restrict the main chain to a lesser extent. The preferred position for this constraining non-cysteine residue is adjacent to one of the invariant cysteines, however, it may be one of the other bridged residues. If the span is seven, we prefer to include two amino acids that limit main-chain conformation. These amino acids could be at any of the seven positions, but are preferably the two bridged residues that are immediately adjacent to the cysteines. If the span is eight or nine, additional constraining amino acids may be provided.

The disulfide bond of a class I mini-proteins is exposed to solvent. Thus, one should avoid exposing the variegated population of GPs that display class I mini-proteins to reagents that rupture disulfides; Creighton names several such reagents (CREI88).

Class II mini-proteins are those featuring a single disulfide bond having a span of greater than nine amino acids. The bridged amino acids form secondary structures which help to stabilize their conformation. Preferably, these intermediate amino acids form hairpin supersecondary structures such as those schematized below:





5 Secondary structures are stabilized by hydrogen bonds
 between amide nitrogen and carbonyl groups, by interac-
 tions between charged side groups and helix dipoles, and
 by van der Waals contacts. One abundant secondary
 structure in proteins is the α -helix. The α helix has 3.6
 10 residues per turn, a 1.5 Å rise per residue, and a helical
 radius of 2.3 Å. All observed α -helices are right-handed.
 The torsion angles ϕ (-57°) and ψ (-47°) are favorable
 for most residues, and the hydrogen bond between the
 backbone carbonyl oxygen of each residue and the backbone
 15 NH of the fourth residue along the chain is 2.86 Å long
 (nearly the optimal distance) and virtually straight.
 Since the hydrogen bonds all point in the same direction,
 the α helix has a considerable dipole moment (carboxy
 terminus negative).

20 The β strand may be considered an elongated helix
 with 2.3 residues per turn, a translation of 3.3 Å per
 residue, and a helical radius of 1.0 Å. Alone, a β strand
 forms no main-chain hydrogen bonds. Most commonly, β
 strands are found in twisted (rather than planar) paral-
 25 lel, antiparallel, or mixed parallel/antiparallel sheets.

A peptide chain can form a sharp reverse turn. A
 reverse turn may be accomplished with as few as four amino
 acids. Reverse turns are very abundant, comprising a
 quarter of all residues in globular proteins. In pro-
 30 teins, reverse turns commonly connect β strands to form β
 sheets, but may also form other connections. A peptide
 can also form other turns that are less sharp.

Based on studies of known proteins, one may calculate
 the propensity of a particular residue, or of a particular

dipeptide or tripeptide, to be found in an α helix, β strand or reverse turn. The normalized frequencies of occurrence of the amino acid residues in these secondary structures is given in Table 6-4 of CREI84. For a more
 5 detailed treatment on the prediction of secondary structure from the amino acid sequence, see Chapter 6 of SCHU79.

In designing a suitable hairpin structure, one may copy an actual structure from a protein whose three-
 10 dimensional conformation is known, design the structure using frequency data, or combine the two approaches. Preferably, one or more actual structures are used as a model, and the frequency data is used to determine which mutations can be made without disrupting the structure.

15 Preferably, no more than three amino acids lie between the cysteine and the beginning or end of the α helix or β strand.

More complex structures (such as a double hairpin) are also possible.

20 Class III mini-proteins are those featuring a plurality of disulfide bonds. They optionally may also feature secondary structures such as those discussed above with regard to Class II mini-proteins. Since the number of possible disulfide bond topologies increases rapidly
 25 with the number of bonds (two bonds, three topologies; three bonds, 15 topologies; four bonds, 105 topologies) the number of disulfide bonds preferably does not exceed four. With two or more disulfide bonds, the disulfide bridge spans preferably do not exceed 50, and the largest
 30 intercysteine chain segment preferably does not exceed 20.

Naturally occurring class III mini-proteins, such as heat-stable enterotoxin ST-Ia frequently have pairs of cysteines that are adjacent in the amino-acid sequence.

TOPSECRET-5503660

Adjacent cysteines are very unlikely to form an intramolecular disulfide and cysteines separated by a single amino acid form an intramolecular disulfide with difficulty and only for certain intervening amino acids.

- 5 Thus, clustering cysteines within the amino-acid sequence reduces the number of realizable disulfide bonding schemes. We utilize such clustering in the class III mini-protein disclosed herein.

Metal Finger Mini-Proteins. The mini-proteins of the
 10 present invention are not limited to those crosslinked by disulfide bonds. Another important class of mini-proteins are analogues of finger proteins. Finger proteins are characterized by finger structures in which a metal ion is coordinated by two Cys and two His residues, forming a
 15 tetrahedral arrangement around it. The metal ion is most often zinc(II), but may be iron, copper, cobalt, etc. The "finger" has the consensus sequence (Phe or Tyr)-(1 AA)-Cys-(2-4 AAs)-Cys-(3 AAs)-Phe-(5 AAs)-Leu-(2 AAs)-His-(3 AAs)-His-(5 AAs) (BERG88; GIBS88). While finger proteins
 20 typically contain many repeats of the finger motif, it is known that a single finger will fold in the presence of zinc ions (FRAN87; PARR88). There is some dispute as to whether two fingers are necessary for binding to DNA. The present invention encompasses mini-proteins with either
 25 one or two fingers. It is to be understood that the target need not be a nucleic acid.

G. Modified PBSS

There exist a number of enzymes and chemical reagents that can selectively modify certain side groups of
 30 proteins, including: a) protein-tyrosine kinase, Ellmans reagent, methyl transferases (that methylate GLU side groups), serine kinases, proline hydroxylases, vitamin-K dependent enzymes that convert GLU to GLA, maleic anhydride, and alkylating agents. Treatment of the variegated

FOI 5696860

population of GP(PBD)s with one of these enzymes or reagents will modify the side groups affected by the chosen enzyme or reagent. Enzymes and reagents that do not kill the GP are much preferred. Such modification of side groups can directly affect the binding properties of the displayed PBDs. Using affinity separation methods, we enrich for the modified GPs that bind the predetermined target. Since the active binding domain is not entirely genetically specified, we must repeat the post-morphogenesis modification at each enrichment round. This approach is particularly appropriate with mini-protein IPBDs because we envision chemical synthesis of these SBDs.

III. VARIATION STRATEGY -- MUTAGENESIS TO OBTAIN POTENTIAL BINDING DOMAINS WITH DESIRED DIVERSITY

III.A. Generally

Using standard genetic engineering techniques, a molecule of variegated DNA can be introduced into a vector so that it constitutes part of a gene (OLIP86, OLIP87, AUSU87, REID88a). When vector containing variegated DNA are used to transform bacteria, each cell makes a version of the original protein. Each colony of bacteria may produce a different version from any other colony. If the variegations of the DNA are concentrated at loci known to be on the surface of the protein or in a loop, a population of proteins will be generated, many members of which will fold into roughly the same 3D structure as the parent protein. The specific binding properties of each member, however, may be different from each other member.

We now consider the manner in which we generate a diverse population of potential binding domains in order to facilitate selection of a PBD-bearing GP which binds with the requisite affinity to the target of choice. The

FOIb290"56096560

potential binding domains are first designed at the amino acid level. Once we have identified which residues are to be mutagenized, and which mutations to allow at those positions, we may then design the variegated DNA which is to encode the various PBDs so as to assure that there is a reasonable probability that if a PBD has an affinity for the target, it will be detected. Of course, the number of independent transformants obtained and the sensitivity of the affinity separation technology will impose limits on the extent of variegation possible within any single round of variegation.

There are many ways to generate diversity in a protein. (See RICH86, CARU85, and OLIP86.) At one extreme, we vary a few residues of the protein as much as possible (inter alia see CARU85, CARU87, RICH86, and WHAR86). We will call this approach "Focused Mutagenesis". A typical "Focused Mutagenesis" strategy is to pick a set of five to seven residues and vary each through 13-20 possibilities. An alternative plan of mutagenesis ("Diffuse Mutagenesis") is to vary many more residues through a more limited set of choices (See VERS86a and PAKU86). The variegation pattern adopted may fall between these extremes, e.g., two residues varied through all twenty amino acids, two more through only two possibilities, and a fifth into ten of the twenty amino acids.

There is no fixed limit on the number of codons which can be mutated simultaneously. However, it is desirable to adopt a mutagenesis strategy which results in a reasonable probability that a possible PBD sequence is in fact displayed by at least one genetic package. When the size of the set of amino acids potentially encoded by each variable codon is the same for all variable codons and within the set all amino acids are equiprobable, this probability may be calculated as follows: Let $\Gamma(k,q)$ be

the probability that amino acid number k will occur at variegated codon q ; these codons need not be contiguous. The probability that a particular vgDNA molecule will encode a PBD containing n variegated amino acids k_1, \dots, k_n is:

$$p(k_1, \dots, k_n) = \Gamma(k_1, 1) \cdot \dots \cdot \Gamma(k_n, n)$$

Consider a library of N_{it} independent transformants prepared with said vgDNA; the probability that the sequence k_1, \dots, k_n is absent is:

$$P(\text{missing } k_1, \dots, k_n) = \exp\{-N_{it} \cdot p(k_1, \dots, k_n)\}.$$

$$P(k_1, \dots, k_n \text{ in lib}) = 1 - \exp\{-N_{it} \cdot p(k_1, \dots, k_n)\}.$$

Preferably, the probability that a mutein encoded by the vgDNA and composed of the least favored amino acids at each variegated position will be displayed by at least one independent transformant in the library is at least 0.50, and more preferably at least 0.90. (Muteins composed of more favored amino acids would of course be more likely to occur in the same library.)

Preferably, the variegation is such as will cause a typical transformant population to display 10^6 - 10^7 different amino acid sequences by means of preferably not more than 10-fold more (more preferably not more than 3-fold) different DNA sequences.

For a mini-protein that lacks α helices and β strands, one will, in any given round of mutation, preferably variegate each of 4-6 non-cysteine codons so that they each encode at least eight of the 20 possible amino acids. The variegation at each codon could be customized to that position. Preferably, cysteine is not one of the potential substitutions, though it is not excluded.

When the mini-protein is a metal finger protein, in a

FOI b6 b7C b7D

typical variegation strategy, the two Cys and two His residues, and optionally also the aforementioned Phe/Tyr, Phe and Leu residues, are held invariant and a plurality (usually 5-10) of the other residues are varied.

5 When the mini-protein is of the type featuring one or more α helices and β strands, the set of potential amino acid modifications at any given position is picked to favor those which are less likely to disrupt the secondary structure at that position. Since the number of possibilities at each variable amino acid is more limited, the
10 total number of variable amino acids may be greater without altering the sampling efficiency of the selection process.

For the last-mentioned class of mini-proteins, as
15 well as domains other than mini-proteins, preferably not more than 20 and more preferably 5-10 codons will be variegated. However, if diffuse mutagenesis is employed, the number of codons which are variegated can be higher.

The decision as to which residues to modify is eased
20 by knowledge of which residues lie on the surface of the domain and which are buried in the interior.

We choose residues in the IPBD to vary through consideration of several factors, including: a) the 3D structure of the IPBD, b) sequences homologous to IPBD,
25 and c) modeling of the IPBD and mutants of the IPBD. When the number of residues that could strongly influence binding is greater than the number that should be varied simultaneously, the user should pick a subset of those residues to vary at one time. The user picks trial levels
30 of variegation and calculate the abundances of various sequences. The list of varied residues and the level of variegation at each varied residue are adjusted until the composite variegation is commensurate with the sensitivity
-- of the affinity separation and the number of independent

FD6250-5696360

transformants that can be made.

Preferably, the abundance of PPBD-encoding DNA is 3 to 10 times higher than both $1/M_{ntv}$ and $1/C_{sensi}$ to provide a margin of redundancy. M_{ntv} is the number of transformants that can be made from Y_{D100} DNA. With current technology M_{ntv} is approximately $5 \cdot 10^8$, but the exact value depends on the details of the procedures adapted by the user. Improvements in technology that allow more efficient: a) synthesis of DNA, b) ligation of DNA, or c) transformation of cells will raise the value of M_{ntv} . C_{sensi} is the sensitivity of the affinity separation; improvements in affinity separation will raise C_{sensi} . If the smaller of M_{ntv} and C_{sensi} is increased, higher levels of variegation may be used. For example, if C_{sensi} is 1 in 10^9 and M_{ntv} is 10^8 , then improvements in C_{sensi} are less valuable than improvements in M_{ntv} .

While variegation normally will involve the substitution of one amino acid for another at a designated variable codon, it may involve the insertion or deletion of amino acids as well.

III.B. Identification of Residues to be Varied

We now consider the principles that guide our choice of residues of the IPBD to vary. A key concept is that only structured proteins exhibit specific binding, *i.e.* can bind to a particular chemical entity to the exclusion of most others. Thus the residues to be varied are chosen with an eye to preserving the underlying IPBD structure. Substitutions that prevent the PBD from folding will cause GPs carrying those genes to bind indiscriminately so that they can easily be removed from the population.

Sauer and colleagues (PAKU86, REID88), and Caruthers and colleagues (EISE85) have shown that some residues on the polypeptide chain are more important than others in

T06250-5696360

determining the 3D structure of a protein. The 3D structure is essentially unaffected by the identity of the amino acids at some loci; at other loci only one or a few types of amino acid is allowed. In most cases, loci where wide variety is allowed have the amino acid side group directed toward the solvent. Loci where limited variety is allowed frequently have the side group directed toward other parts of the protein. Thus substitutions of amino acids that are exposed to solvent are less likely to affect the 3D structure than are substitutions at internal loci. (See also SCHU79, p169-171 and CREI84, p239-245, 314-315).

The residues that join helices to helices, helices to sheets, and sheets to sheets are called turns and loops and have been classified by Richardson (RICH81), Thornton (THOR88), Sutcliffe et al. (SUTC87a) and others. Insertions and deletions are more readily tolerated in loops than elsewhere. Thornton et al. (THOR88) have summarized many observations indicating that related proteins usually differ most at the loops which join the more regular elements of secondary structure. (These observations are relevant not only to the variegation of potential binding domains but also to the insertion of binding domains into an outer surface protein of a genetic package, as discussed in a later section.)

Burial of hydrophobic surfaces so that bulk water is excluded is one of the strongest forces driving the binding of proteins to other molecules. Bulk water can be excluded from the region between two molecules only if the surfaces are complementary. We should test as many surface variations as possible to find one that is complementary to the target. The selection-through-binding isolates those proteins that are more nearly complementary to some surface on the target.

Proteins do not have distinct, countable faces. Therefore we define an "interaction set" to be a set of residues such that all members of the set can simultaneously touch one molecule of the target material without any atom of the target coming closer than van der Waals distance to any main-chain atom of the IPBD. The concept of a residue "touching" a molecule of the target is discussed below. From a picture of BPTI (such as Figure 6-10, p. 225 of CREI84) we can see that residues 3, 7, 8, 10, 13, 39, 41, and 42 can all simultaneously contact a molecule the size and shape of myoglobin. We also see that residue 49 can not touch a single myoglobin molecule simultaneously with any of the first set even though all are on the surface of BPTI. (It is not the intent of the present invention, however, to suggest that use of models is required to determine which part of the target molecule will actually be the site of binding by PBD.)

Variations in the position, orientation and nature of the side chains of the residues of the interaction set will alter the shape of the potential binding surface defined by that set. Any individual combination of such variations may result in a surface shape which is a better or a worse fit for the target surface. The effective diversity of a variegated population is measured by the number of distinct shapes the potentially complementary surfaces of the PBD can adopt, rather than the number of protein sequences. Thus, it is preferable to maximize the former number, when our knowledge of the IPBD permits us to do so.

To maximize the number of surface shapes generated for when N residues are varied, all residues varied in a given round of variegation should be in the same interaction set because variation of several residues in one

interaction set generates an exponential number of different shapes of the potential binding surface.

If cassette mutagenesis is to be used to introduce the variegated DNA into the ipbd gene, the protein residues to be varied are, preferably, close enough together in sequence that the variegated DNA (vgDNA) encoding all of them can be made in one piece. The present invention is not limited to a particular length of vgDNA that can be synthesized. With current technology, a stretch of 60 amino acids (180 DNA bases) can be spanned.

Further, when there is reason to mutate residues further than sixty residues apart, one can use other mutational means, such as single-stranded-oligonucleotide-directed mutagenesis (BOTS85) using two or more mutating primers.

Alternatively, to vary residues separated by more than sixty residues, two cassettes may be mutated as follows: 1) vg DNA having a low level of variegation (for example, 20 to 400 fold variegation) is introduced into one cassette in the OCV, 2) cells are transformed and cultured, 3) vg OCV DNA is obtained, 4) a second segment of vgDNA is inserted into a second cassette in the OCV, and 5) cells are transformed and cultured, GPs are harvested and subjected to selection-through-binding.

The composite level of variation preferably does not exceed the prevailing capabilities to a) produce very large numbers of independently transformed cells or b) detect small components in a highly varied population. The limits on the level of variegation are discussed later.

Data about the IPBD and the target that are useful in deciding which residues to vary in the variegation cycle include: 1) 3D structure, or at least a list of residues

TOP SECRET 56096860

on the surface of the IPBD, 2) list of sequences homologous to IPBD, and 3) model of the target molecule or a stand-in for the target.

These data and an understanding of the behavior of
5 different amino acids in proteins will be used to answer two questions:

- 1) which residues of the IPBD are on the outside and close enough together in space to touch the target simultaneously?
- 10 2) which residues of the IPBD can be varied with high probability of retaining the underlying IPBD structure?

Although an atomic model of the target material (obtained through X-ray crystallography, NMR, or other
15 means) is preferred in such examination, it is not necessary. For example, if the target were a protein of unknown 3D structure, it would be sufficient to know the molecular weight of the protein and whether it were a soluble globular protein, a fibrous protein, or a membrane
20 protein. Physical measurements, such as low-angle neutron diffraction, can determine the overall molecular shape, viz. the ratios of the principal moments of inertia. One can then choose a protein of known structure of the same class and similar size and shape to use as a molecular
25 stand-in and yardstick. It is not essential to measure the moments of inertia of the target because, at low resolution, all proteins of a given size and class look much the same. The specific volumes are the same, all are more or less spherical and therefore all proteins of the same size
30 and class have about the same radius of curvature. The radii of curvature of the two molecules determine how much of the two molecules can come into contact.

The most appropriate method of picking the residues of the protein chain at which the amino acids should be

T06250-5606660

varied is by viewing, with interactive computer graphics, a model of the IPBD. A stick-figure representation of molecules is preferred. A suitable set of hardware is an Evans & Sutherland PS390 graphics terminal (Evans & Sutherland Corporation, Salt Lake City, UT) and a MicroVAX II supermicro computer (Digital Equipment Corp., Maynard, MA). The computer should, preferably, have at least 150 megabytes of disk storage, so that the Brookhaven Protein Data Bank can be kept on line. A FORTRAN compiler, or some equally good higher-level language processor is preferred for program development. Suitable programs for viewing and manipulating protein models include: a) PS-FRODO, written by T. A. Jones (JONE85) and distributed by the Biochemistry Department of Rice University, Houston, TX; and b) PROTEUS, developed by Dayringer, Tramantano, and Fletterick (DAYR86). Important features of PS-FRODO and PROTEUS that are needed to view and manipulate protein models for the purposes of the present invention are the abilities to: 1) display molecular stick figures of proteins and other molecules, 2) zoom and clip images in real time, 3) prepare various abstract representations of the molecules, such as a line joining C α s and side group atoms, 4) compute and display solvent-accessible surfaces reasonably quickly, 5) point to and identify atoms, and 6) measure distance between atoms.

In addition, one could use theoretical calculations, such as dynamic simulations of proteins, to estimate whether a substitution at a particular residue of a particular amino-acid type might produce a protein of approximately the same 3D structure as the parent protein. Such calculations might also indicate whether a particular substitution will greatly affect the flexibility of the protein; calculations of this sort may be useful but are not required.

Residues whose mutagenesis is most likely to affect binding to a target molecule, without destabilizing the protein, are called the "principal set". Using the knowledge of which residues are on the surface of the IPBD

5 (as noted above), we pick residues that are close enough together on the surface of the IPBD to touch a molecule of the target simultaneously without having any IPBD main-chain atom come closer than van der Waals distance (viz. 4.0 to 5.0 Å) from any target atom. For the purposes of

10 the present invention, a residue of the IPBD "touches" the target if: a) a main-chain atom is within van der Waals distance; viz. 4.0 to 5.0 Å of any atom of the target molecule, or b) the C_β is within D_{cutoff} of any atom of the target molecule so that a side-group atom could make

15 contact with that atom.

Because side groups differ in size (cf. Table 35), some judgment is required in picking D_{cutoff} . In the preferred embodiment, we will use $D_{\text{cutoff}} = 8.0$ Å, but other values in the range 6.0 Å to 10.0 Å could be used.

20 If IPBD has G at a residue, we construct a pseudo C_β with the correct bond distance and angles and judge the ability of the residue to touch the target from this pseudo C_β .

Alternatively, we choose a set of residues on the

25 surface of the IPBD such that the curvature of the surface defined by the residues in the set is not so great that it would prevent contact between all residues in the set and a molecule of the target. This method is appropriate if the target is a macromolecule, such as a protein, because

30 the PBDs derived from the IPBD will contact only a part of the macromolecular surface. The surfaces of macromolecules are irregular with varying curvatures. If we pick residues that define a surface that is not too convex, then there will be a region on a macromolecular target

T05290-5095860

with a compatible curvature.

In addition to the geometrical criteria, we prefer that there be some indication that the underlying IPBD structure will tolerate substitutions at each residue in the principal set of residues. Indications could come from various sources, including: a) homologous sequences, b) static computer modeling, or c) dynamic computer simulations.

The residues in the principal set need not be contiguous in the protein sequence and usually are not. The exposed surfaces of the residues to be varied do not need to be connected. We desire only that the amino acids in the residues to be varied all be capable of touching a molecule of the target material simultaneously without having atoms overlap. If the target were, for example, horse heart myoglobin, and if the IPBD were BPTI, any set of residues in one interaction set of BPTI defined in Table 34 could be picked.

The secondary set comprises those residues not in the primary set that touch residues in the primary set. These residues might be excluded from the primary set because: a) the residue is internal; b) the residue is highly conserved, or c) the residue is on the surface, but the curvature of the IPBD surface prevents the residue from being in contact with the target at the same time as one or more residues in the primary set.

Internal residues are frequently conserved and the amino acid type can not be changed to a significantly different type without substantial risk that the protein structure will be disrupted. Nevertheless, some conservative changes of internal residues, such as I to L or F to Y, are tolerated. Such conservative changes subtly affect the placement and dynamics of adjacent protein residues and such "fine tuning" may be useful once an SBD is found.

T06290-5603660

Surface residues in the secondary set are most often located on the periphery of the principal set. Such peripheral residues can not make direct contact with the target simultaneously with all the other residues of the principal set. The charge on the amino acid in one of these residues could, however, have a strong effect on binding. Once an SBD is found, it is appropriate to vary the charge of some or all of these residues. For example, the variegated codon containing equimolar A and G at base 1, equimolar C and A at base 2, and A at base 3 yields amino acids T, A, K, and E with equal probability.

The assignment of residues to the primary and secondary sets may be based on: a) geometry of the IPBD and the geometrical relationship between the IPBD and the target (or a stand-in for the target) in a hypothetical complex, and b) sequences of proteins homologous to the IPBD. However, it should be noted that the distinction between the principal set and the secondary set is one more of convenience than of substance; we could just as easily have assigned each amino acid residue in the domain a preference score that weighed together the different considerations affecting whether they are suitable for variegation, and then ranked the residues in order, from most preferred to least.

For any given round of variegation, it may be necessary to limit the variegation to a subset of the residues in the primary and secondary sets, based on geometry and on the maximum allowed level of variegation that assures progressivity. The allowed level of variegation determines how many residues can be varied at once; geometry determines which ones.

The user may pick residues to vary in many ways. For example, pairs of residues are picked that are diametrically opposed across the face of the principal set. Two

T06250"5605650

The number of residues picked is coupled to the range through which each can be varied by the restrictions discussed below. In the first round, we do not assume any binding between IPBD and the target and so progressivity is not an issue. At the first round, the user may elect to produce a level of variegation such that each molecule of vgDNA is potentially different through. for example, unlimited variegation of 10 codons (20^{10} approx. = 10^{13}). One run of the DNA synthesizer produces approximately 10^{13} molecules of length 100 nts. Inefficiencies in ligation and transformation will reduce the number of proteins actually tested to between 10^7 and $5 \cdot 10^8$. Multiple replications of the process with such very high levels of variegation will not yield repeatable results; the user decides whether this is important.

Having picked which residues to vary, we now decide the range of amino acids to allow at each variable residue. The total level of variegation is the product of the number of variants at each varied residue. Each varied residue can have a different scheme of variegation, producing 2 to 20 different possibilities. The set of amino acids which are potentially encoded by a given

variegated codon are called its "substitution set".

The computer that controls a DNA synthesizer, such as the Milligen 7500, can be programmed to synthesize any base of an oligo-nt with any distribution of nts by taking
 5 some nt substrates (e.g. nt phosphoramidites) from each of two or more reservoirs. Alternatively, nt substrates can be mixed in any ratios and placed in one of the extra reservoir for so called "dirty bottle" synthesis. Each codon could be programmed differently. The "mix" of bases
 10 at each nucleotide position of the codon determines the relative frequency of occurrence of the different amino acids encoded by that codon.

Simply variegated codons are those in which those nucleotide positions which are degenerate are obtained
 15 from a mixture of two or more bases mixed in equimolar proportions. These mixtures are described in this specification by means of the standardized "ambiguous nucleotide" code (Table 1 and 37 CFR §1.822). In this code, for example, in the degenerate codon "SNT", "S"
 20 denotes an equimolar mixture of bases G and C, "N", an equimolar mixture of all four bases, and "T", the single invariant base thymidine.

Complexly variegated codons are those in which at least one of the three positions is filled by a base from
 25 an other than equimolar mixture of two of more bases.

Either simply or complexly variegated codons may be used to achieve the desired substitution set.

If we have no information indicating that a particular amino acid or class of amino acid is appropriate, we
 30 strive to substitute all amino acids with equal probability because representation of one mini-protein above the detectable level is wasteful. Equal amounts of all four nts at each position in a codon (NNN) yields the amino

TOP SECRET 560560

acid distribution in which each amino acid is present in proportion to the number of codons that code for it. This distribution has the disadvantage of giving two basic residues for every acidic residue. In addition, six times
 5 as much R, S, and L as W or M occur. If five codons are synthesized with this distribution, each of the 243 sequences encoding some combination of L, R, and S are 7776-times more abundant than each of the 32 sequences encoding some combination of W and M. To have five Ws
 10 present at detectable levels, we must have each of the (L,R,S) sequences present in 7776-fold excess...

Preferably, we also consider the interactions between the sites of variegation and the surrounding DNA. If the method of mutagenesis to be used is replacement of a
 15 cassette, we consider whether the variegation will generate gratuitous restriction sites and whether they seriously interfere with the intended introduction of diversity. We reduce or eliminate gratuitous restriction sites by appropriate choice of variegation pattern and
 20 silent alteration of codons neighboring the sites of variegation.

It is generally accepted that the sequence of amino acids in a protein or polypeptide determine the three-dimensional structure of the molecule, including the
 25 possibility of no definite structure. Among polypeptides of definite length and sequence, some have a defined tertiary structure and most do not.

Particular amino acid residues can influence the tertiary structure of a defined polypeptide in several
 30 ways, including by:

- a) affecting the flexibility of the polypeptide main chain,
- b) adding hydrophobic groups,
- c) adding charged groups,

T06230-5096360

d) allowing hydrogen bonds, and

e) forming cross-links, such as disulfides, chelation to metal ions, or bonding to prosthetic groups.

Most works on proteins classify the twenty amino acids into categories such as hydrophobic/hydrophilic, positive/negative/neutral, or large/small. These classifications are useful rules of thumb, but one must be careful not to oversimplify. Proteins contain a variety of identifiable secondary structural features, including: a) α helices, b) 3-10 helices, c) anti-parallel β sheets, d) parallel β sheets, e) Ω loops, f) reverse turns, and g) various cross links. Many people have analyzed proteins of known structures and assigned each amino-acid to one category or another. Using the frequency at which particular amino acids occur in various types of secondary structures, people have a) tried to predict the secondary structures of proteins for which only the amino-acid sequence is known (CHOU74, CHOU78a, CHOU78b), and b) designed proteins de novo that have a particular set of secondary structural elements (DEGR87, HECH90). Although some amino acids show definite predilection for one secondary form (e.g. VAL for β structure and ALA for α helices), these preferences are not very strong; Creighton has tabulated the preferences (CREI84). In only seven cases does the tendency exceed 2.0:

Amino acid	distinction	ratio
MET	α /turn	3.7
PRO	turn/ α	3.7
VAL	β /turn	3.2
GLY	turn/ α	2.9
ILE	β /turn	2.8
PHE	β /turn	2.3
LEU	α /turn	2.2

Every amino-acid type has been observed in every identified secondary structural motif. ARG is particularly indiscriminate.

5

10

Flexibility:

20

25

30

tendency to exist in a β sheet.

The main chain of proline is particularly constrained by the cyclic side group. The ϕ angle is always close to -60° . Most prolines are found near the surface of the protein.

Charge:

LYS and ARG carry a single positive charge at any pH below 10.4 or 12.0, respectively. Nevertheless, the methylene groups, four and three respectively, of these amino acids are capable of hydrophobic interactions. The guanidinium group of ARG is capable of donating five hydrogens simultaneously, while the amino group of LYS can donate only three. Furthermore, the geometries of these groups is quite different, so that these groups are often not interchangeable.

ASP and GLU carry a single negative charge at any pH above ≈ 4.5 and 4.6 , respectively. Because ASP has but one methylene group, few hydrophobic interactions are possible. The geometry of ASP lends itself to forming hydrogen bonds to main-chain nitrogens which is consistent with ASP being found very often in reverse turns and at the beginning of helices. GLU is more often found in α helices and particularly in the amino-terminal portion of these helices because the negative charge of the side group has a stabilizing interaction with the helix dipole (NICH88, SALI88).

HIS has an ionization pK in the physiological range, viz. 6.2. This pK can be altered by the proximity of charged groups or of hydrogen donors or acceptors. HIS is capable of forming bonds to metal ions such as zinc, copper, and iron.

Hydrogen bonds:

Aside from the charged amino acids, SER, THR, ASN,

T06290"56096860

GLN, TYR, and TRP can participate in hydrogen bonds.

Cross links:

The most important form of cross link is the disulfide bond formed between two thiols, especially the thiols of CYS residues. In a suitably oxidizing environment, these bonds form spontaneously. These bonds can greatly stabilize a particular conformation of a protein or mini-protein. When a mixture of oxidized and reduced thiol reagents are present, exchange reactions take place that allow the most stable conformation to predominate. Concerning disulfides in proteins and peptides, see also KATZ90, MATS89, PERR84, PERR86, SAUE86, WELL86, JANA89, HORV89, KISH85, and SCHN86.

Other cross links that form without need of specific enzymes include:

- | | | |
|----|---|---|
| 1, | (CYS) ₄ :Fe | Rubredoxin (in CREI84, P.376) |
| 2) | (CYS) ₄ :Zn | Aspartate Transcarbamylase (in CREI84, P.376) and Zn-fingers (HARD90) |
| 3) | (HIS) ₂ (MET)(CYS):Cu | Azurin (in CREI84, P.376) and Basic "Blue" Cu Cucumber protein (GUSS88) |
| 4) | (HIS) ₄ :Cu | CuZn superoxide dismutase |
| 5) | (CYS) ₄ :(Fe ₄ S ₄) | Ferredoxin (in CREI84, P.376) |
| 6) | (CYS) ₂ (HIS) ₂ :Zn | Zinc-fingers (GIBS88) |
| 7) | (CYS) ₃ (HIS):Zn | Zinc-fingers (GAUS87, GIBS88) |

Cross links having (HIS)₂(MET)(CYS):Cu has the potential advantage that HIS and MET can not form other cross links without Cu.

Simply Variegated Codons

The following simply variegated codons are useful because they encode a relatively balanced set of amino acids:

- 1) SNT which encodes the set [L,P,H,R,V,A,D,G]: a) one acidic (D) and one basic (R), b) both aliphatic (L,V) and aromatic hydrophobics (H), c) large (L,R,H) and small (G,A) side groups, d) ridged (P) and flexible (G) amino acids, e) each amino acid encoded once.
- 2) RNG which encodes the set [M,T,K,R,V,A,E,G]: a) one acidic and two basic (not optimal, but acceptable), b) hydrophilics and hydrophobics, c) each amino acid encoded once.
- 3) RMG which encodes the set [T,K,A,E]: a) one acidic, one basic, one neutral hydrophilic, b) three favor α helices, c) each amino acid encoded once.
- 4) VNT which encodes the set [L,P,H,R,I,T,N,S,V,A,D,G]: a) one acidic, one basic, b) all classes: charged, neutral hydrophilic, hydrophobic, ridged and flexible, etc., c) each amino acid encoded once.
- 5) RRS which encodes the set [N,S,K,R,D,E,G²]: a) two acidics, two basics, b) two neutral hydrophilics, c) only glycine encoded twice.
- 6) NNT which encodes the set [F,S,Y,C,L,P,H,R,I,T,N,V,A,D,G]: a) sixteen DNA sequences provide fifteen different amino acids; only serine is repeated, all others are present in equal amounts (This allows very efficient sampling of the library.), b) there are equal numbers of acidic and basic amino acids (D and R, once each), c) all major classes of amino acids are present: acidic, basic, aliphatic hydrophobic, aromatic hydrophobic, and neutral hydrophilic.
- 7) NNG, which encodes the set [L²,R²,S,W,P,Q,M,T,K,V,A,-E,G, stop]: a) fair preponderance of residues that favor formation of α -helices [L,M,A,Q,K,E; and, to a lesser extent, S,R,T]; b) encodes 13 different amino acids. (VHG encodes a subset of the set encoded by NNG which encodes 9 amino acids in nine different DNA

sequences, with equal acids and bases, and 5/9 being α helix-favoring.)

For the initial variegation, NNT is preferred, in most cases. However, when the codon is encoding an amino acid to be incorporated into an α helix, NNG is preferred.

Below, we analyze several simple variegations as to the efficiency with which the libraries can be sampled.

Libraries of random hexapeptides encoded by (NNK)⁶ have been reported (SCOT90, CWIR90). Table 130 shows the expected behavior of such libraries. NNK produces single codons for PHE, TYR, CYS, TRP, HIS, GLN, ILE, MET, ASN, LYS, ASP, and GLU (α set); two codons for each of VAL, ALA, PRO, THR, and GLY (Φ set); and three codons for each of LEU, ARG, and SER (Ω set). We have separated the 64,000,000 possible sequences into 28 classes, shown in Table 130A, based on the number of amino acids from each of these sets. The largest class is $\Phi\Omega\alpha\alpha\alpha\alpha$ with $\approx 14.6\%$ of the possible sequences. Aside from any selection, all the sequences in one class have the same probability of being produced. Table 130B shows the probability that a given DNA sequence taken from the (NNK)⁶ library will encode a hexapeptide belonging to one of the defined classes; note that only $\approx 6.3\%$ of DNA sequences belong to the $\Phi\Omega\alpha\alpha\alpha\alpha$ class.

Table 130C shows the expected numbers of sequences in each class for libraries containing various numbers of independent transformants (viz. 10^6 , $3 \cdot 10^6$, 10^7 , $3 \cdot 10^7$, 10^8 , $3 \cdot 10^8$, 10^9 , and $3 \cdot 10^9$). At 10^6 independent transformants (ITs), we expect to see 56% of the $\Omega\Omega\Omega\Omega\Omega\Omega$ class, but only 0.1% of the $\alpha\alpha\alpha\alpha\alpha\alpha$ class. The vast majority of sequences seen come from classes for which less than 10% of the class is sampled. Suppose a peptide from, for example, class $\Phi\Phi\Omega\Omega\alpha\alpha$ is isolated by fractionating the library for binding to a target. Consider how much we

09896095-62901
T06290-5696860

5 LEU at position 2, but ARG or SER could be better. Even if we isolate a peptide of the $\Omega\Omega\Omega\Omega\Omega$ class, there is a noticeable chance that better members of the class were not present in the library.

With a library of 10^7 ITs, we see that several
10 classes have been completely sampled, but that the $\alpha\alpha\alpha\alpha\alpha$
class is only 1.1% sampled. At $7.6 \cdot 10^7$ ITs, we expect
display of 50% of all amino-acid sequences, but the
classes containing three or more amino acids of the α set
are still poorly sampled. To achieve complete sampling of
15 the (NNK)⁶ library requires about $3 \cdot 10^9$ ITs, 10-fold
larger than the largest (NNK)⁶ library so far reported.

Table 131 shows expectations for a library encoded by (NNT)⁴(NNG)². The expectations of abundance are independent of the order of the codons or of interspersed
20 unvaried codons. This library encodes 0.133 times as many amino-acid sequences, but there are only 0.0165 times as many DNA sequences. Thus $5.0 \cdot 10^7$ ITs (i.e. 60-fold fewer
than required for (NNK)⁶) gives almost complete sampling of the library. The results would be slightly better for
25 (NNT)⁶ and slightly, but not much, worse for (NNG)⁶. The controlling factor is the ratio of DNA sequences to amino-acid sequences.

Table 132 shows the ratio of #DNA sequences/#AA sequences for codons NNK, NNT, and NNG. For NNK and NNG, we have assumed that the PBD is displayed as part of an essential gene, such as gene III in Ff phage, as is indicated by the phrase "assuming stops vanish". It is not in any way required that such an essential gene be used. If a non-essential gene is used, the analysis would

be slightly different; sampling of NNK and NNG would be slightly less efficient. Note that (NNT)⁶ gives 3.6-fold more amino-acid sequences than (NNK)⁵ but requires 1.7-fold fewer DNA sequences. Note also that (NNT)⁷ gives
 5 twice as many amino-acid sequences as (NNK)⁶, but 3.3-fold fewer DNA sequences.

Thus, while it is possible to use a simple mixture (NNS, NNK or NNN) to obtain at a particular position all twenty amino acids, these simple mixtures lead to a highly
 10 biased set of encoded amino acids. This problem can be overcome by use of complexly variegated codons.

Complexly Variegated Codons

Let Abun(x) be the abundance of DNA sequences coding for amino acid x, defined by the distribution of nts at
 15 each base of the codon. For any distribution, there will be a most-favored amino acid (mfaa) with abundance Abun(mfaa) and a least-favored amino acid (lfaa) with abundance Abun(lfaa). We seek the nt distribution that allows all twenty amino acids and that yields the largest
 20 ratio Abun(lfaa)/Abun(mfaa) subject, if desirable to further constraints.

We first will present the mixture calculated to be optimal when the nt distribution is subject to two constraints: equal abundances of acidic and basic amino
 25 acids and the least possible number of stop codons. Thus only nt distributions that yield Abun(E)+Abun(D) = Abun(R)+Abun(K) are considered, and the function maximized is:

$$\{(1-\text{Abun}(\text{stop})) (\text{Abun}(\text{lfaa})/\text{Abun}(\text{mfaa}))\}.$$

30 We have simplified the search for an optimal nt distribution by limiting the third base to T or G (C or G is equivalent). All amino acids are possible and the number of accessible stop codons is reduced because TGA and TAA

T062330-5605660

codons are eliminated. The amino acids F, Y, C, H, N, I, and D require T at the third base while W, M, Q, K, and E require G. Thus we use an equimolar mixture of T and G at the third base. However, it should be noted that the present invention embraces use of complexly variegated codons in which the third base is not limited to T or G (or to C or G).

A computer program, written as part of the present invention and named "Find Optimum vgCodon" (See Table 9), varies the composition at bases 1 and 2, in steps of 0.05, and reports the composition that gives the largest value of the quantity $\{(Abun(lfaa)/Abun(mfaa) (1-Abun(stop)))\}$. A vg codon is symbolically defined by the nucleotide distribution at each base:

15		<u>T</u>	<u>C</u>	<u>A</u>	<u>G</u>
	base #1 =	t1	c1	a1	g1
	base #2 =	t2	c2	a2	g2
	base #3 =	t3	c3	a3	g3
		$t1 + c1 + a1 + g1 = 1.0$			
20		$t2 + c2 + a2 + g2 = 1.0$			
		$t3 = g3 = 0.5, \quad c3 = a3 = 0.$			

The variation of the quantities t1, c1, a1, g1, t2, c2, a2, and g2 is subject to the constraint that:

$$\begin{aligned}
 &Abun(E)+Abun(D) = Abun(K)+Abun(R) \\
 25 \quad &Abun(E)+Abun(D) = g1*a2 \\
 &Abun(K)+Abun(R) = a1*a2/2 + c1*g2 + a1*g2/2 \\
 &g1*a2 = a1*a2/2 + c1*g2 + a1*g2/2
 \end{aligned}$$

Solving for g2, we obtain

$$g2 = (g1*a2 - 0.5*a1*a2)/(c1 + 0.5*a1) .$$

30 In addition,

$$\begin{aligned}
 t1 &= 1 - a1 - c1 - g1 \\
 t2 &= 1 - a2 - c2 - g2 .
 \end{aligned}$$

We vary a1, c1, g1, a2, and c2 and then calculate t1, g2, and t2. Initially, variation is in steps of 5%. Once an approximately optimum distribution of nucleotides is determined, the region is further explored with steps of 1%. The logic of this program is shown in Table 9. The optimum distribution (the "fxS" codon) is shown in Table 10A and yields DNA molecules encoding each type amino acid with the abundances shown.

Note that this chemistry encodes all twenty amino acids, with acidic and basic amino acids being equiprobable, and the most favored amino acid (serine) is encoded only 2.454 times as often as the least favored amino acid (tryptophan). The "fxS" vg codon improves sampling most for peptides containing several of the amino acids [F,Y,C,W,H,Q,I,M,N,K,D,E] for which NNK or NNS provide only one codon. Its sampling advantages are most pronounced when the library is relatively small.

A modification of "Fine Optimum vgCodon" varies the composition at bases 1 and 2, in steps of 0.01, and reports the composition that gives the largest value of the quantity $\{(Abun(lfaa)/Abun(mfaa))\}$ without any restraint on the relative abundance of any amino acids. The results of this optimization is shown in Table 10B. The changes are small, indicating that insisting on equality of acids and bases and minimizing stop codons costs us little. Also note that, without restraining the optimization, the prevalence of acidic and basic amino acids comes out fairly close. On the other hand, relaxing the restriction leaves a distribution in which the least favored amino acid is only .412 times as prevalent as SER.

The advantages of an NNT codon are discussed elsewhere in the present application. Unoptimized NNT provides 15 amino acids encoded by only 16 DNA sequences. It is possible to improve on NNT as follows. First note

that the SER codons occur in the T and A rows of the genetic-code table and in the C and G columns.

$$[\text{SER}] = T_1 \times C_2 + A_1 \times G_2$$

If we reduce the prevalence of SER by reducing T_1 , C_2 , A_1 , and G_2 relative to other bases, then we will also reduce the prevalence of PHE, TYR, CYS, PRO, THR, ALA, ARG, GLY, ILE, and ASN. The prevalence of LEU, HIS, VAL, and ASP will rise. If we assume that T_1 , C_2 , A_1 , and G_2 are all lowered to the same extent and that C_1 , G_1 , T_2 , and A_2 are increased by the same amount, we can compute a shift that makes the prevalence of SER equal the prevalences of LEU, HIS, VAL, and ASP. The decrease in PHE, TYR, CYS, PRO, THR, ALA, ARG, GLY, ILE, and ASN is not equal; CYS and THR are reduced more than the others.

15 Let the distribution be

	<u>T</u>	<u>C</u>	<u>A</u>	<u>G</u>
base #1 =	.25-q	.25+q	.25-q	.25+q
base #2 =	.25+q	.25-q	.25+q	.25-q
base #3 =	1.00	0.0	0.0	0.0

20 Setting [SER] = [LEU] = [HIS] = [VAL] = [ASP] gives:

$$(.25-q) \cdot (.25-q) + (.25-q) \cdot (.25-q) = (.25+q) \cdot (.25+q)$$

$$2 \cdot (.25 - q)^2 = (.25 + q)^2$$

$$q^2 - 1.5q + .0625 = 0$$

$$q = (3/4) - \sqrt{2}/2 = .0428$$

25 This distribution (shown in Table 10C) gives five amino acids (SER, LEU, HIS, VAL, ASP) in very nearly equal amounts. A further eight amino acids (PHE, TYR, ILE, ASN, PRO, ALA, ARG, GLY) are present at 78% the abundance of SER. THR and CYS remain at half the abundance of SER.

30 When variegating DNA for disulfide-bonded mini-proteins,
it is often desirable to reduce the prevalence of CYS.
This distribution allows 13 amino acids to be seen at high
level and gives no stops; the optimized fxs distribution

The NNG codon can also be optimized. Table 10D shows an approximately optimized NNG codon. When equimolar

20 When synthesizing vgDNA by the "dirty bottle" method,
it is sometimes desirable to use only a limited number of
mixes.. One very useful mixture is called the "optimized
NNS mixture" in which we average the first two positions
of the fxS mixture: $T_1 = 0.24$, $C_1 = 0.17$, $A_1 = 0.33$, $G_1 =$
25 0.26 , the second position is identical to the first, $C_3 =$
 $G_3 = 0.5$. This distribution provides the amino acids ARG,
SER, LEU, GLY, VAL, THR, ASN, and LYS at greater than 5%
plus ALA, ASP, GLU, ILE, MET, and TYR at greater than 4%.

An additional complexly variegated codon is of
30 interest. This codon is identical to the optimized NNT
codon at the first two positions and has T:G::90:10 at the
third position. This codon provides thirteen amino acids
(ALA, ILE, ARG, SER, ASP, LEU, VAL, PHE, ASN, GLY, PRO,
TYR, and HIS) at more than 5.5%. THR at 4.3% and CYS at

3.9% are more common than the LFAAs of NNK (3.125%). The remaining five amino acids are present at less than 1%. This codon has the feature that all amino acids are present; sequences having more than two of the low-
 5 abundance amino acids are rare. When we isolate an SBD using this codon, we can be reasonably sure that the first 13 amino acids were tested at each position. A similar codon, based on optimized NNG, could be used.

Table 10E shows some properties of an unoptimized NNS
 10 (or NNK) codon. Note that there are three equally most-favored amino acids: ARG, LEU, and SER. There are also twelve equally least favored amino acids: PHE, ILE, MET, TYR, HIS, GLN, ASN, LYS, ASP, GLU, CYS, and TRP. Five amino acids (PRO, THR, ALA, VAL, GLY) fall in between.
 15 Note that a six-fold repetition of NNS gives sequences composed of the amino acids [PHE, ILE, MET, TYR, HIS, GLN, ASN, LYS, ASP, GLU, CYS, and TRP] at only $\approx 1\%$ of the sequences composed of [ARG, LEU, and SER]. Not only is this ≈ 20 -fold lower than the prevalence of (TRP/MET)⁶ in
 20 optimized NNG⁶ vgdDNA, but this low prevalence applies to twelve amino acids.

Diffuse Mutagenesis.

Diffuse Mutagenesis can be applied to any part of the protein at any time, but is most appropriate when some
 25 binding to the target has been established. Diffuse Mutagenesis can be accomplished by spiking each of the pure nts activated for DNA synthesis (e.g. nt-phosphoramidites) with a small amount of one or more of the other activated nts.

30 Contrary to general practice, the present invention sets the level of spiking so that only a small percentage (1% to .00001%, for example) of the final product will contain the initial DNA sequence. This will insure that many single, double, triple, and higher mutations occur,

FO5290-5096860

5 that is the majority component, is

$$M = \exp\{ \log_e(Q)/N_b \} = 10^{(\log_{10}(Q)/N_b)}.$$

If, for example, thirty base pairs on the DNA chain were to be varied and 1% of the product is to have the parental sequence, then each mixed nt substrate should contain 86% of the parental nt and 14% of other nts. Table 8 shows the fraction (fn) of DNA molecules having n non-parental bases when 30 bases are synthesized with reagents that contain fraction M of the majority component. When $M=.63096$, f_{24} and higher are less than 10^{-8} . The entry "most" in Table 8 is the number of changes that has the highest probability. Note that substantial probability for multiple substitutions only occurs if the fraction of parental sequence (f_0) is allowed to drop to around 10^{-6} . The N_b base pairs of the DNA chain that are synthesized with mixed reagents need not be contiguous. They are picked so that between $N_b/3$ and N_b codons are affected to various degrees. The residues picked for mutation are picked with reference to the 3D structure of the IPBD, if known. For example, one might pick all or most of the residues in the principal and secondary set. We may impose restrictions on the extent of variation at each of these residues based on homologous sequences or other data. The mixture of non-parental nts need not be random, rather mixtures can be biased to give particular amino acid types specific probabilities of appearance at each codon. For example, one residue may contain a hydrophobic amino acid in all known homologous sequences; in such a case, the first and third base of that codon would be varied, but the second would be set to T. Other examples

of how this might be done are given in the horse heart myoglobin example. This diffuse structure-directed mutagenesis will reveal the subtle changes possible in protein backbone associated with conservative interior changes, such as V to I, as well as some not so subtle changes that require concomitant changes at two or more residues of the protein.

III.D. Special Considerations Relating to Variegation of Mini-Proteins with Essential Cysteines

Several of the preferred simple or complex variegated codons encode a set of amino acids which includes cysteine. This means that some of the encoded binding domains will feature one or more cysteines in addition to the invariant disulfide-bonded cysteines. For example, at each NNT-encoded position, there is a one in sixteen chance of obtaining cysteine. If six codons are so varied, the fraction of domains containing additional cysteines is 0.33. Odd numbers of cysteines can lead to complications, see Perry and Wetzel (PERR84). On the other hand, many disulfide-containing proteins contain cysteines that do not form disulfides, e.g. trypsin. The possibility of unpaired cysteines can be dealt with in several ways:

First, the variegated phage population can be passed over an immobilized reagent that strongly binds free thiols, such as SulfoLink (catalogue number 44895 H from Pierce Chemical Company, Rockford, Illinois, 61105). Another product from Pierce is TNB-Thiol Agarose (Catalogue Code 20409 H). BioRad sells Affi-Gel 401 (catalogue 153-4599) for this purpose.

Second, one can use a variegation that excludes cysteines, such as:

NHT that gives [F,S,Y,L,P,H,I,T,N,V,A,D],

T06230-5605660

VNS that gives

[L², P², H, Q, R³, I, M, T², N, K, S, V², A², E, D, G²],

NNG that gives [L², S, W, P, Q, R², M, T, K, R, V, A, E, G, stop],

SNT that gives [L, P, H, R, V, A, D, G],

5 RNG that gives [M, T, K, R, V, A, E, G],

RMG that gives [T, K, A, E],

VNT that gives [L, P, H, R, I, T, N, S, V, A, D, G], or

RRS that gives [N, S, K, R, D, E, G²].

However, each of these schemes has one or more of the
10 disadvantages, relative to NNT: a) fewer amino acids are
allowed, b) amino acids are not evenly provided, c) acidic
and basic amino acids are not equally likely), or d) stop
codons occur. Nonetheless, NNG, NHT, and VNT are almost
as useful as NNT. NNG encodes 13 different amino acids
15 and one stop signal. Only two amino acids appear twice in
the 16-fold mix.

Thirdly, one can enrich the population for binding to
the preselected target, and evaluate selected sequences
post hoc for extra cysteines. Those that contain more
20 cysteines than the cysteines provided for conformational
constraint may be perfectly usable. It is possible that
a disulfide linkage other than the designed one will
occur. This does not mean that the binding domain defined
by the isolated DNA sequence is in any way unsuitable.
25 The suitability of the isolated domains is best determined
by chemical and biochemical evaluation of chemically
synthesized peptides.

Lastly, one can block free thiols with reagents, such
as Ellman's reagent, iodoacetate, or methyl iodide, that
30 specifically bind free thiols and that do not react with
disulfides, and then leave the modified phage in the
population. It is to be understood that the blocking
agent may alter the binding properties of the mini-
protein; thus, one might use a variety of blocking reagent

T06290 56096350

III.E Planning the Second and Later Rounds of Variegation

If the first round of variegation results in some binding to the target, but the affinity for the target is still too low, further improvement may be achieved by variegation of the SBDs. Preferably, the process is progressive, i.e. each variegation cycle produces a better starting point for the next variegation cycle than the previous cycle produced. Setting the level of variegation such that the ppbd and many sequences related to the ppbd sequence are present in detectable amounts ensures that

the process is progressive. If the level of variegation is so high that the ppbd sequence is present at such low levels that there is an appreciable chance that no transformant will display the PPBD, then the best SBD of the next round could be worse than the PPBD. At excessively high level of variegation, each round of mutagenesis is independent of previous rounds and there is no assurance of progressivity. This approach can lead to valuable binding proteins, but repetition of experiments with this level of variegation will not yield progressive results. Excessive variation is not preferred.

Progressivity is not an all-or-nothing property. So long as most of the information obtained from previous variegation cycles is retained and many different surfaces that are related to the PPBD surface are produced, the process is progressive. If the level of variegation is so high that the ppbd gene may not be detected, the assurance of progressivity diminishes. If the probability of recovering PPBD is negligible, then the probability of progressive behavior is also negligible.

A level of variegation that allows recovery of the PPBD has two properties:

- 1) we can not regress because the PPBD is available,
- 2) an enormous number of multiple changes related to the PPBD are available for selection and we are able to detect and benefit from these changes.

It is very unlikely that all of the variants will be worse than the PPBD; we desire the presence of PPBD at detectable levels to insure that all the sequences present are indeed related to PPBD.

An opposing force in our design considerations is that PBDs are useful in the population only up to the amount that can be detected; any excess above the detec-

T06250-55095B50

table amount is wasted. Thus we produce as many surfaces related to PPBD as possible within the constraint that the PPBD be detectable.

If the level of variegation in the previous variegation cycle was correctly chosen, then the amino acids selected to be in the residues just varied are the ones best determined. The environment of other residues has changed, so that it is appropriate to vary them again. Because there are often more residues in the principal and secondary sets than can be varied simultaneously, we start by picking residues that either have never been varied (highest priority) or that have not been varied for one or more cycles. If we find that varying all the residues except those varied in the previous cycle does not allow a high enough level of diversity, then residues varied in the previous cycle might be varied again. For example, if M_{ntv} (the number of independent transformants that can be produced from Y_{D100} of DNA) and C_{sensi} (the sensitivity of the affinity separation) were such that seven residues could be varied, and if the principal and secondary sets contained 13 residues, we would always vary seven residues, even though that implies varying some residue twice in a row. In such cases, we would pick the residues just varied that contain the amino acids of highest abundance in the variegated codons used.

It is the accumulation of information that allows the process to select those protein sequences that produce binding between the SBD and the target. Some interfaces between proteins and other molecules involve twenty or
30 more residues. Complete variation of twenty residues would generate 10^{26} different proteins. By dividing the residues that lie close together in space into overlapping groups of five to seven residues, we can vary a large surface but never need to test more than 10^7 to 10^9

candidates at once, a savings of 10^{19} to 10^{17} fold. The power of selection with accumulation of information is well illustrated in Chapter 3 of DAWK86.

Use of NNT or NNG variegated codons leads to very efficient sampling of variegated libraries because the ratio of (different amino-acid sequences)/(different DNA sequences) is much closer to unity than it is for NNK or even the optimized vg codon (fxS). Nevertheless, a few amino acids are omitted in each case. Both NNT and NNG allow members of all important classes of amino acids: hydrophobic, hydrophilic, acidic, basic, neutral hydrophilic, small, and large. After selecting a binding domain, a subsequent variegation and selection may be desirable to achieve a higher affinity or specificity. During this second variegation, amino acid possibilities overlooked by the preceding variegation may be investigated

In the first round, we assume that the parental protein has no known affinity for the target material. For example, consider the parental mini-protein, similar to that discussed in Example 11, having the structure X_1 - C_2 - X_3 - X_4 - X_5 - X_6 - C_7 - X_8 in which C_2 and C_7 form a disulfide bond. Introduction of extra cysteines may cause alternative structures to form which might be disadvantageous. Accidental cysteines at positions 4 or 5 are thought to be potentially more troublesome than at the other positions. We adopt the pattern of variegation: X_1 :NNT, X_3 :NNT, X_4 :NNG, X_5 :NNG, X_6 :NNT, and X_8 :NNT, so that cysteine can not occur at positions 4 and 5. (Table 131 shows the number of different amino acids expected in libraries prepared with DNA variegated in this way and comprising different numbers of independent transformants.)

In the second round of variegation, a preferred strategy is to vary each position through a new set of residues which includes the amino acid(s) which were found

FOUO 56096860

5
10

15

25

THR is another possibility with the NNT codon. If THR is selected, we switch to NNG because that includes the previously excluded possibilities and includes THR. Suppose the successful amino acid encoded by the NNT codon

was ASP. We use RRS at the next variegation because this includes both acidic amino acids plus LYS and ARG. One could also use VRS to allow GLN.

Thus, later rounds of variegation test both amino acid positions not previously mutated, and amino acid substitutions at a previously mutated position which were not within the previous substitution set.

If the first round of variegation is entirely unsuccessful, a different pattern of variegation should be used. For example, if more than one interaction set can be defined within a domain, the residues varied in the next round of variegation should be from a different set than that probed in the initial variegation. If repeated failures are encountered, one may switch to a different IPBD.

IV. DISPLAY STRATEGY: DISPLAYING FOREIGN BINDING DOMAINS ON THE SURFACE OF A "GENETIC PACKAGE"

IV.A. General Requirements for Genetic Packages

It is emphasized that the GP on which selection-through-binding will be practiced must be capable, after the selection, either of growth in some suitable environment or of in vitro amplification and recovery of the encapsulated genetic message. During at least part of the growth, the increase in number is preferably approximately exponential with respect to time. The component of a population that exhibits the desired binding properties may be quite small, for example, one in 10^6 or less. Once this component of the population is separated from the non-binding components, it must be possible to amplify it. Culturing viable cells is the most powerful amplification of genetic material known and is preferred. Genetic messages can also be amplified in vitro, e.g. by PCR, but this is not the most preferred method.

T05290-5603660

Preferred GPs are vegetative bacterial cells, bacterial spores and bacterial DNA viruses. Eukaryotic cells could be used as genetic packages but have longer dividing times and more stringent nutritional requirements than do bacteria and it is much more difficult to produce a large number of independent transformants. They are also more fragile than bacterial cells and therefore more difficult to chromatograph without damage. Eukaryotic viruses could be used instead of bacteriophage but must be propagated in eukaryotic cells and therefore suffer from some of the amplification problems mentioned above.

Nonetheless, a strain of any living cell or virus is potentially useful if the strain can be: 1) genetically altered with reasonable facility to encode a potential binding domain, 2) maintained and amplified in culture, 3) manipulated to display the potential binding protein domain where it can interact with the target material during affinity separation, and 4) affinity separated while retaining the genetic information encoding the displayed binding domain in recoverable form. Preferably, the GP remains viable after affinity separation.

When the genetic package is a bacterial cell, or a phage which is assembled periplasmically, the display means has two components. The first component is a secretion signal which directs the initial expression product to the inner membrane of the cell (a host cell when the package is a phage). This secretion signal is cleaved off by a signal peptidase to yield a processed, mature, potential binding protein. The second component is an outer surface transport signal which directs the package to assemble the processed protein into its outer surface. Preferably, this outer surface transport signal is derived from a surface protein native to the genetic package.

For example, in a preferred embodiment, the hybrid gene comprises a DNA encoding a potential binding domain operably linked to a signal sequence (e.g., the signal sequences of the bacterial phoA or bla genes or the signal sequence of M13 phage geneIII) and to DNA encoding a coat protein (e.g., the M13 gene III or gene VIII proteins) of a filamentous phage (e.g., M13). The expression product is transported to the inner membrane (lipid bilayer) of the host cell, whereupon the signal peptide is cleaved off to leave a processed hybrid protein. The C-terminus of the coat protein-like component of this hybrid protein is trapped in the lipid bilayer, so that the hybrid protein does not escape into the periplasmic space. (This is typical of the wild-type coat protein.) As the single-stranded DNA of the nascent phage particle passes into the periplasmic space, it collects both wild-type coat protein and the hybrid protein from the lipid bilayer. The hybrid protein is thus packaged into the surface sheath of the filamentous phage, leaving the potential binding domain exposed on its outer surface. (Thus, the filamentous phage, not the host bacterial cell, is the "replicable genetic package" in this embodiment.)

If a secretion signal is necessary for the display of the potential binding domain, in an especially preferred embodiment the bacterial cell in which the hybrid gene is expressed is of a "secretion-permissive" strain.

When the genetic package is a bacterial spore, or a phage whose coat is assembled intracellularly, a secretion signal directing the expression product to the inner membrane of the host bacterial cell is unnecessary. In these cases, the display means is merely the outer surface transport signal, typically a derivative of a spore or phage coat protein.

FOI 250 56096850

20 The fusion of ipbd and osp fragments may also include
fragments of random or pseudorandom DNA to produce a
population, members of which may display IPBD on the GP
surface. The members displaying IPBD are isolated by
screening or selection for the display-of-binding pheno-
25 type.

Preferably, the GP is readily stored, for example, by freezing. If the GP is a cell, it should have a short

doubling time, such as 20-40 minutes. If the GP is a virus, it should be prolific, e.g., a burst size of at least 100/infected cell. GPs which are finicky or expensive to culture are disfavored. The GP should be
 5 easy to harvest, preferably by centrifugation. The GP is preferably stable for a temperature range of -70 to 42°C (stable at 4°C for several days or weeks); resistant to shear forces found in HPLC; insensitive to UV; tolerant of desiccation; and resistant to a pH of 2.0 to 10.0, surface
 10 active agents such as SDS or Triton, chaotropes such as 4M urea or 2M guanidinium HCl, common ions such as K⁺, Na⁺, and SO₄²⁻, common organic solvents such as ether and acetone, and degradative enzymes. Finally, there must be a suitable OCV.

15 Although knowledge of specific OSPs may not be required for vegetative bacterial cells and endospores, the user of the present invention, preferably, will know: Is the sequence of any osp known? (preferably yes, at least one required for phage). How does the OSP arrive at
 20 the surface of GP? (knowledge of route necessary, different routes have different uses, no route preferred per se). Is the OSP post-translationally processed? (no processing most preferred, predictable processing preferred over unpredictable processing). What rules are known
 25 governing this processing, if there is any processing? (no processing most preferred, predictable processing acceptable). What function does the OSP serve in the outer surface? (preferably not essential). Is the 3D structure of an OSP known? (highly preferred). Are fusions between
 30 fragments of osp and a fragment of x known? Does expression of these fusions lead to X appearing on the surface of the GP? (fusion data is as preferred as knowledge of a 3D structure). Is a "2D" structure of an OSP available? (in this context, a "2D" structure indicates which
 35 residues are exposed on the cell surface) (2D structure

105250" 55095560

less preferred than 3D structure). Where are the domain boundaries in the OSP? (not as preferred as a 2D structure, but acceptable). Could IPBD go through the same process as OSP and fold correctly? (IPBD might need
 5 prosthetic groups) (preferably IPBD will fold after same process). Is the sequence of an osp promoter known? (preferably yes). Is osp gene controlled by regulatable promoter available? (preferably yes). What activates this promoter? (preferably a diffusible chemical, such as
 10 IPTG). How many different OSPs do we know? (the more the better). How many copies of each OSP are present on each package? (more is better).

The user will want knowledge of the physical attributes of the GP: How large is the GP? (knowledge useful
 15 in deciding how to isolate GPs) (preferably easy to separate from soluble proteins such as IgGs). What is the charge on the GP? (neutral preferred). What is the sedimentation rate of the GP? (knowledge preferred, no particular value preferred).

20 The preferred GP, OCV and OSP are those for which the fewest serious obstacles can be seen, rather than the one that scores highest on any one criterion.

Viruses are preferred over bacterial cells and spores (cp. LUIT85 and references cited therein). The virus is
 25 preferably a DNA virus with a genome size of 2 kb to 10 kb base pairs, such as (but not limited to) the filamentous (Ff) phage M13, fd, and f1 (inter alia see RASC86, BOEK80, BOEK82, DAYL88, GRAY81b, KUHN88, LOPE85, WEBS85, MARV75, MARV80, MOSE82, CRIS84, SMIT88a, SMIT88b) ; the IncN
 30 specific phage Ike and If1 (NAKA81, PEET85, PEET87, THOM83, THOM88a); IncP-specific Pseudomonas aeruginosa phage Pf1 (THOM83, THOM88a) and Pf3 (LUIT83, LUIT85, LUTI87, THOM88a); and the Xanthomonas oryzae phage Xf (THOM83, THOM88a). Filamentous phage are especially

FOI 2005-03095

Preferred OSPs for several GPs are given in Table 2. References to osp-ipbd fusions in this section should be taken to apply, mutatis mutandis, to osp-pbd and osp-sbd fusions as well.

IV.B. Phages for Use as GPs:

The morphogenetic pathway of the phage determines the

5

10

20

The filamentous phages (e.g., M13) are of particular interest.

25

The user must choose a site in the candidate OSP gene for inserting a ipbd gene fragment. The coats of most bacteriophage are highly ordered. Filamentous phage can be described by a helical lattice; isometric phage, by an icosahedral lattice. Each monomer of each major coat protein sits on a lattice point and makes defined interactions with each of its neighbors. Proteins that fit into the lattice by making some, but not all, of the normal lattice contacts are likely to destabilize the virion by: a) aborting formation of the virion, b) making the virion unstable, or c) leaving gaps in the virion so that the nucleic acid is not protected. Thus in bacteriophage, unlike the cases of bacteria and spores, it is important to retain in engineered OSP-IPBD fusion proteins those residues of the parental OSP that interact with other proteins in the virion. For M13 gVIII, we retain the entire mature protein, while for M13 gIII, it might suffice to retain the last 100 residues (or even fewer).

Such a truncated gIII protein would be expressed in parallel with the complete gIII protein, as gIII protein is required for phage infectivity.

Il'ichev *et al.* (ILIC89) have reported viable phage
 5 having alterations in gene VIII. In one case, a point mutation changed one amino acid near the amino terminus of the mature gVIII protein from GLU to ASP. In the other case, five amino acids were inserted at the site of the first mutation. They suggested that similar constructions
 10 could be used for vaccines. They did not report on any binding properties of the modified phage, nor did they suggest mutagenizing the inserted material. Furthermore, they did not insert a binding domain, nor did they suggest inserting such a domain.

15 Further considerations on the design of the ipbd::osp gene is discussed in section IV.F.

Filamentous phage:

Compared to other bacteriophage, filamentous phage in general are attractive and M13 in particular is especially
 20 attractive because: 1) the 3D structure of the virion is known; 2) the processing of the coat protein is well understood; 3) the genome is expandable; 4) the genome is small; 5) the sequence of the genome is known; 6) the virion is physically resistant to shear, heat, cold, urea,
 25 guanidinium Cl, low pH, and high salt; 7) the phage is a sequencing vector so that sequencing is especially easy; 8) antibiotic-resistance genes have been cloned into the genome with predictable results (HINE80); 9) It is easily cultured and stored (FRIT85), with no unusual or expensive
 30 media requirements for the infected cells, 10) it has a high burst size, each infected cell yielding 100 to 1000 M13 progeny after infection; and 11) it is easily harvested and concentrated (SALI64, FRIT85).

The filamentous phage include M13, f1, fd, If1, Ike, Xf, Pf1, and Pf3.

The entire life cycle of the filamentous phage M13, a common cloning and sequencing vector, is well understood.

5 M13 and fl are so closely related that we consider the properties of each relevant to both (RASC86); any differentiation is for historical accuracy. The genetic structure (the complete sequence (SCHA78), the identity and function of the ten genes, and the order of transcription and location of the promoters) of M13 is well known
10 as is the physical structure of the virion (BANN81, BOEK80, CHAN79, ITOK79, KAPL78, KUHN85b, KUHN87, MAK080, MARV78, MESS78, OHKA81, RASC86, RUSS81, SCHA78, SMIT85, WEBS78, and ZIMM82); see RASC86 for a recent review of the structure and function of the coat proteins. Because the
15 genome is small (6423 bp), cassette mutagenesis is practical on RF M13 (AUSU87), as is single-stranded oligonucleotide directed mutagenesis (FRIT85). M13 is a plasmid and transformation system in itself, and an ideal sequencing
20 vector. M13 can be grown on Rec⁻ strains of E. coli. The M13 genome is expandable (MESS78, FRIT85) and M13 does not lyse cells. Because the M13 genome is extruded through the membrane and coated by a large number of identical protein molecules, it can be used as a cloning vector
25 (WATS87 p278, and MESS77). Thus we can insert extra genes into M13 and they will be carried along in a stable manner.

Marvin and collaborators (MARV78, MAK080, BANN81) have determined an approximate 3D virion structure of f1 by a combination of genetics, biochemistry, and X-ray diffraction from fibers of the virus. Figure 4 is drawn after the model of Banner et al. (BANN81) and shows only the C α s of the protein. The apparent holes in the cylindrical sheath are actually filled by protein side

5

10

20

30

structure of M13 coat protein is implicit in the 3D structure. Mature M13 gene VIII protein has only one domain.

When the GP is M13 the gene III and the gene VIII proteins are highly preferred as OSP (see Examples I through IV). The proteins from genes VI, VII, and IX may also be used.

As discussed in the Examples, we have constructed a tripartite gene comprising:

- 1) DNA encoding a signal sequence directing secretion of parts (2) and (3) through the inner membrane,
- 2) DNA encoding the mature BPTI sequence, and
- 3) DNA encoding the mature M13 gVIII protein.

This gene causes BPTI to appear in active form on the surface of M13 phage.

The gene VIII protein is a preferred OSP because it is present in many copies and because its location and orientation in the virion are known (BANN81). Preferably, the PBD is attached to the amino terminus of the mature M13 coat protein. Had direct fusion of PBD to M13 CP failed to cause PBD to be displayed on the surface of M13, we would have varied part of the mini-protein sequence and/or insert short random or nonrandom spacer sequences between mini-protein and M13 CP. The 3D model of f1 indicates strongly that fusing IPBD to the amino terminus of M13 CP is more likely to yield a functional chimeric protein than any other fusion site.

Similar constructions could be made with other filamentous phage. Pf3 is a well known filamentous phage that infects Pseudomonas aeruginosa cells that harbor an IncP-1 plasmid. The entire genome has been sequenced (LUIT85) and the genetic signals involved in replication and assembly are known (LUIT87). The major coat protein of PF3 is unusual in having no signal peptide to direct

T05250-56096850

its secretion. The sequence has charged residues ASP₇, ARG₃₇, LYS₄₀, and PHE₄₄-COO⁻ which is consistent with the amino terminus being exposed. Thus, to cause an IPBD to appear on the surface of Pf3, we construct a tripartite

5 gene comprising:

- 1) a signal sequence known to cause secretion in P. aeruginosa (preferably known to cause secretion of IPBD) fused in-frame to,
- 2) a gene fragment encoding the IPBD sequence, fused
10 in-frame to,
- 3) DNA encoding the mature Pf3 coat protein.

Optionally, DNA encoding a flexible linker of one to 10 amino acids is introduced between the ipbd gene fragment and the Pf3 coat-protein gene. Optionally, DNA encoding
15 the recognition site for a specific protease, such as tissue plasminogen activator or blood clotting Factor Xa, is introduced between the ipbd gene fragment and the Pf3 coat-protein gene. Amino acids that form the recognition site for a specific protease may also serve the function
20 of a flexible linker. This tripartite gene is introduced into Pf3 so that it does not interfere with expression of any Pf3 genes. To reduce the possibility of genetic recombination, part (3) is designed to have numerous silent mutations relative to the wild-type gene. Once the
25 signal sequence is cleaved off, the IPBD is in the periplasm and the mature coat protein acts as an anchor and phage-assembly signal. It matters not that this fusion protein comes to rest in the lipid bilayer by a route different from the route followed by the wild-type
30 coat protein.

The amino-acid sequence of M13 pre-coat (SCHA78), called AA_seq1, is

T06230-56096850

AA_seq1

	1	1	2	2	3	3	4	4	5
5	0	5	0	↓	5	0	5	0	5

5 MKKSLVLKASVAVATLVPMLSFAAEGDDPAKAAFNSLQASATEYIGYAWA

5	6	6	7	7
5	0	5	0	3

10 MVVVIVGATIGIKLFKKFTSKAS

The single-letter codes for amino acids and the codes for ambiguous DNA are given in Table 1. The best site for inserting a novel protein domain into M13 CP is after A23 because SP-I cleaves the precoat protein after A23, as indicated by the arrow. Proteins that can be secreted will appear connected to mature M13 CP at its amino terminus. Because the amino terminus of mature M13 CP is located on the outer surface of the virion, the introduced domain will be displayed on the outside of the virion.

20 The uncertainty of the mechanism by which M13CP appears in the lipid bilayer raises the possibility that direct insertion of bpti into gene VIII may not yield a functional fusion protein. It may be necessary to change the signal sequence of the fusion to, for example, the phoA

25 signal sequence (MKQSTIALALLPLLFTPVTKA.....). Marks et al. (MARK86) showed that the phoA signal peptide could direct mature BPTI to the E. coli periplasm.

Another vehicle for displaying the IPBD is by expressing it as a domain of a chimeric gene containing part or all of gene III. This gene encodes one of the minor coat proteins of M13. Genes VI, VII, and IX also encode minor coat proteins. Each of these minor proteins is present in about 5 copies per virion and is related to morphogenesis or infection. In contrast, the major coat protein is present in more than 2500 copies per virion.

35 The gene VI, VII, and IX proteins are present at the ends of the virion; these three proteins are not post-translationally processed (RASC86).

The single-stranded circular phage DNA associates with about five copies of the gene III protein and is then extruded through the patch of membrane-associated coat protein in such a way that the DNA is encased in a helical sheath of protein (WEBS78). The DNA does not base pair (that would impose severe restrictions on the virus genome); rather the bases intercalate with each other independent of sequence.

Smith (SMIT85) and de la Cruz et al. (DELA88) have shown that insertions into gene III cause novel protein domains to appear on the virion outer surface. The mini-protein's gene may be fused to gene III at the site used by Smith and by de la Cruz et al., at a codon corresponding to another domain boundary or to a surface loop of the protein, or to the amino terminus of the mature protein.

All published works use a vector containing a single modified gene III of fd. Thus, all five copies of gIII are identically modified. Gene III is quite large (1272 b.p. or about 20% of the phage genome) and it is uncertain whether a duplicate of the whole gene can be stably inserted into the phage. Furthermore, all five copies of gIII protein are at one end of the virion. When bivalent target molecules (such as antibodies) bind a pentavalent phage, the resulting complex may be irreversible. Irreversible binding of the GP to the target greatly interferes with affinity enrichment of the GPs that carry the genetic sequences encoding the novel polypeptide having the highest affinity for the target.

To reduce the likelihood of formation of irreversible complexes, we may use a second, synthetic gene that encodes carboxy-terminal parts of III. We might, for example, engineer a gene that consists of (from 5' to 3'):

- 1) a promoter (preferably regulated),
- 2) a ribosome-binding site,

09896095.062901

- 5

10

15

- 25

30

cated fusions are placed downstream of a known promoter, preferably a regulated promoter such as lacUV5, tac, or trp. Fusions (1) and (2) are appropriate when the minor coat protein attaches to the phage from the cytoplasm or by autonomous insertion into the lipid bilayer. Fusion (1) is appropriate if the amino terminus of the minor coat protein is free and (2) is appropriate if the carboxy terminus is free. Fusions (3) and (4) are appropriate if the minor coat protein attaches to the phage from the periplasm or from within the lipid bilayer. Fusion (3) is appropriate if the amino terminus of the minor coat protein is free and (4) is appropriate if the carboxy terminus is free.

Bacteriophage ϕ X174:

The bacteriophage ϕ X174 is a very small icosahedral virus which has been thoroughly studied by genetics, biochemistry, and electron microscopy (See The Single-Stranded DNA Phages (DENH78)). To date, no proteins from ϕ X174 have been studied by X-ray diffraction. ϕ X174 is not used as a cloning vector because ϕ X174 can accept very little additional DNA; the virus is so tightly constrained that several of its genes overlap. Chambers et al. (CHAM82) showed that mutants in gene G are rescued by the wild-type G gene carried on a plasmid so that the host supplies this protein.

Three gene products of ϕ X174 are present on the outside of the mature virion: F (capsid), G (major spike protein, 60 copies per virion), and H (minor spike protein, 12 copies per virion). The G protein comprises 175 amino acids, while H comprises 328 amino acids. The F protein interacts with the single-stranded DNA of the virus. The proteins F, G, and H are translated from a single mRNA in the viral infected cells. If the G protein is supplied from a plasmid in the host, then the

viral g gene is no longer essential. We introduce one or more stop codons into g so that no G is produced from the viral gene. We fuse a pbd gene fragment to h, either at the 3' or 5' terminus. We eliminate an amount of the
 5 viral g gene equal to the size of pbd so that the size of the genome is unchanged.

Large DNA Phages

Phage such as λ or T4 have much larger genomes than do M13 or ϕ X174. Large genomes are less conveniently
 10 manipulated than small genomes. Phage λ has such a large genome that cassette mutagenesis is not practicable. One can not use annealing of a mutagenic oligonucleotide either, because there is no ready supply of single-stranded λ DNA. (λ DNA is packaged as double-stranded
 15 DNA.) Phage such as λ and T4 have more complicated 3D capsid structures than M13 or ϕ X174, with more OSs to choose from. Intracellular morphogenesis of phage λ could cause protein domains that contain disulfide bonds in their folded forms not to fold.

20 Phage λ virions and phage T4 virions form intracellularly, so that IPBDs requiring large or insoluble prosthetic groups might fold on the surfaces of these phage.

RNA Phages

25 RNA phage are not preferred because manipulation of RNA is much less convenient than is the manipulation of DNA. If the RNA phage MS2 were modified to make room for an osp-ipbd gene and if a message containing the A protein binding site and the gene for a chimera of coat protein
 30 and a PBD were produced in a cell that also contained A protein and wild-type coat protein (both produced from regulated genes on a plasmid), then the RNA coding for the chimeric protein would get packaged. A package comprising

5

15

IV.C. Bacterial Cells as Genetic Packages:

20

25

5

10

15

30

and the signal peptide moiety is extruded into the periplasmic space. There, it is cleaved off by a "signal peptidase", and the remaining "mature" protein can now enter the periplasm. Once there, other cellular mechanisms recognize structures in the mature protein which indicate that its proper place is on the outer membrane, and transport it to that location.

It is well known that the DNA coding for the leader or signal peptide from one protein may be attached to the DNA sequence coding for another protein, protein X, to form a chimeric gene whose expression causes protein X to appear free in the periplasm (BECK83, INOU86 Ch10, LEEC86, MARK86, and BOQU87). That is, the leader causes the chimeric protein to be secreted through the lipid bilayer; once in the periplasm, it is cleaved off by the signal peptidase SP-I.

The use of export-permissive bacterial strains (LISS85, STAD89) increases the probability that a signal-sequence-fusion will direct the desired protein to the cell surface. Liss et al. (LISS85) showed that the mutation prlA4 makes E. coli more permissive with respect to signal sequences. Similarly, Stader et al. (STAD89) found a strain that bears a prlG mutation and that permits export of a protein that is blocked from export in wild-type cells. Such export-permissive strains are preferred.

OSP-IPBD fusion proteins need not fill a structural role in the outer membranes of Gram-negative bacteria because parts of the outer membranes are not highly ordered. For large OSPs there is likely to be one or more sites at which osp can be truncated and fused to ipbd such that cells expressing the fusion will display IPBDs on the cell surface. Fusions of fragments of omp genes with fragments of an x gene have led to X appearing on the outer membrane (CHAR88b, BENS84, CLEM81). When such

fusions have been made, we can design an osp-ipbd gene by substituting ipbd for x in the DNA sequence. Otherwise, a successful OMP-IPBD fusion is preferably sought by fusing fragments of the best omp to an ipbd, expressing the fused gene, and testing the resultant GPs for display-of-IPBD phenotype. We use the available data about the OMP to pick the point or points of fusion between omp and ipbd to maximize the likelihood that IPBD will be displayed. (Spacer DNA encoding flexible linkers, made, e.g., of GLY, SER, and ASN, may be placed between the osp- and ipbd-derived fragments to facilitate display.) Alternatively, we truncate osp at several sites or in a manner that produces osp fragments of variable length and fuse the osp fragments to ipbd; cells expressing the fusion are screened or selected which display IPBDs on the cell surface. Freudl *et al.* (FREU89) have shown that fragments of OSPs (such as OmpA) above a certain size are incorporated into the outer membrane. An additional alternative is to include short segments of random DNA in the fusion of omp fragments to ipbd and then screen or select the resulting variegated population for members exhibiting the display-of-IPBD phenotype.

In E. coli, the LamB protein is a well understood OSP and can be used (BENS84, CHAR90, RONC90, VAND90, CHAP90, MOLL90, CHAR88b, CHAR88c, CLEM81, DARG88, FERE82a, FERE82b, FERE83, FERE84, FERE86a, FERE86b, FERE89a, FERE89b, GEHR87, HALL82, NAKA86a, STAD86, HEIN88, BENS87b, BENS87c, BOUG84, BOUL86a, CHAR84) . The E. coli LamB has been expressed in functional form in S. typhimurium (DEV84, BARB85, HARK87), V. cholerae (HARK86), and K. pneumonia (DEV84, WEHM89), so that one could display a population of PBDs in any of these species as a fusion to E. coli LamB. K. pneumonia expresses a maltoporin similar to LamB (WEHM89) which could also be used. In P. aeruginosa, the D1 protein (a homologue of LamB) can be

used (TRIA88).

LamB of E. coli is a porin for maltose and malto-dextrin transport, and serves as the receptor for adsorption of bacteriophages λ and K10. LamB is transported to the outer membrane if a functional N-terminal sequence is present; further, the first 49 amino acids of the mature sequence are required for successful transport (BENS84). As with other OSPs, LamB of E. coli is synthesized with a typical signal-sequence which is subsequently removed.

10 Homology between parts of LamB protein and other outer membrane proteins OmpC, OmpF, and PhoE has been detected (NIKA84), including homology between LamB amino acids 39-49 and sequences of the other proteins. These subsequences may label the proteins for transport to the outer

15 membrane.

The amino acid sequence of LamB is known (CLEM81), and a model has been developed of how it anchors itself to the outer membrane (Reviewed by, among others, BENZ88b). The location of its maltose and phage binding domains are

20 also known (HEIN88). Using this information, one may identify several strategies by which a PBD insert may be incorporated into LamB to provide a chimeric OSP which displays the PBD on the bacterial outer membrane.

When the PBDs are to be displayed by a chimeric transmembrane protein like LamB, the PBD could be inserted into a loop normally found on the surface of the cell (cp. BECK83, MANO86). Alternatively, we may fuse a 5' segment of the osp gene to the ipbd gene fragment; the point of fusion is picked to correspond to a surface-exposed loop

30 of the OSP and the carboxy terminal portions of the OSP are omitted. In LamB, it has been found that up to 60 amino acids may be inserted (CHAR88b) with display of the foreign epitope resulting; the structural features of OmpC, OmpA, OmpF, and PhoE are so similar that one expects

It should be noted that while LamB may be characterized as a binding protein, it is used in the present invention to provide an OSTs; its binding domains are not variegated.

Other bacterial outer surface proteins, such as OmpA, OmpC, OmpF, PhoE, and pilin, may be used in place of LamB and its homologues. OmpA is of particular interest because it is very abundant and because homologues are known in a wide variety of gram-negative bacterial species. Baker et al. (BAKE87) review assembly of proteins into the outer membrane of E. coli and cite a topological model of OmpA (VOGE86) that predicts that residues 19-32, 62-73, 105-118, and 147-158 are exposed on the cell surface. Insertion of a ipbd encoding fragment at about codon 111 or at about codon 152 is likely to cause the IPBD to be displayed on the cell surface. Concerning OmpA, see also MACI88 and MANO88. Porin Protein F of Pseudomonas aeruginosa has been cloned and has sequence homology to OmpA of E. coli (DUCH88). Although this homology is not sufficient to allow prediction of surface-exposed residues on Porin Protein F, the methods used to determine the topological model of OmpA may be applied to Porin Protein F. Works related to use of OmpA as an OSP include BECK80 and MACI88.

Misra and Benson (MISR88a, MISR88b) disclose a topological model of E. coli OmpC that predicts that, among others, residues GLY₁₆₄ and LEU₂₅₀ are exposed on the cell surface. Thus insertion of an ipbd gene fragment at about codon 164 or at about codon 250 of the E. coli ompC gene or at corresponding codons of the S. typhimurium ompC gene is likely to cause IPBD to appear on the cell surface. The ompC genes of other bacterial species may be used. Other works related to OmpC include CATR87 and

CLIC88.

OmpF of E. coli is a very abundant OSP, $\geq 10^4$ copies/cell. Pages et al. (PAGE90) have published a model of OmpF indicating seven surface-exposed segments. Fusion of an ipbd gene fragment, either as an insert or to replace the 3' part of ompF, in one of the indicated regions is likely to produce a functional ompF::ipbd gene the expression of which leads to display of IPBD on the cell surface. In particular, fusion at about codon 111, 177, 217, or 245 should lead to a functional ompF::ipbd gene. Concerning OmpF, see also REID88b, PAGE88, BENS88, TOMM82, and SODE85.

Pilus proteins are of particular interest because piliated cells express many copies of these proteins and because several species (N. gonorrhoeae, P. aeruginosa, Moraxella bovis, Bacteroides nodosus, and E. coli) express related pilins. Getzoff and coworkers (GETZ88, PARG87, SOME85) have constructed a model of the gonococcal pilus that predicts that the protein forms a four-helix bundle having structural similarities to tobacco mosaic virus protein and myohemerythrin. On this model, both the amino and carboxy termini of the protein are exposed. The amino terminus is methylated. Elleman (ELLE88) has reviewed pilins of Bacteroides nodosus and other species and serotype differences can be related to differences in the pilin protein and that most variation occurs in the C-terminal region. The amino-terminal portions of the pilin protein are highly conserved. Jennings et al. (JENN89) have grafted a fragment of foot-and-mouth disease virus (residues 144-159) into the B. nodosus type 4 fimbrial protein which is highly homologous to gonococcal pilin. They found that expression of the 3'-terminal fusion in P. aeruginosa led to a viable strain that makes detectable amounts of the fusion protein. Jennings et al. did not vary the foreign epitope nor did they suggest any varia-

TOE90"5696660

5

10

15

25

30

phospholipases. E. coli phospholipase A, product of the pldA gene, has been cloned and sequenced by de Geus et al. (DEGE84). They found that the protein appears at the cell surface without any posttranslational processing. A ipbd gene fragment can be attached at either terminus or inserted at positions predicted to encode loops in the protein. That phospholipase A arrives on the outer surface without removal of a signal sequence does not prove that a PldA::IPBD fusion protein will also follow this route. Thus we might cause a PldA::IPBD or IPBD:-PldA fusion to be secreted into the periplasm by addition of an appropriate signal sequence. Thus, in addition to simple binary fusion of an ipbd fragment to one terminus of pldA, the constructions:

- 1) ss::ipbd::pldA
- 2) ss::pldA::ipbd

should be tested. Once the PldA::IPBD protein is free in the periplasm it does not remember how it got there and the structural features of PldA that cause it to localize on the outer surface will direct the fusion to the same destination.

IV.D. Bacterial Spores as Genetic Packages:

Bacterial spores have desirable properties as GP candidates. Spores are much more resistant than vegetative bacterial cells or phage to chemical and physical agents, and hence permit the use of a great variety of affinity selection conditions. Also, Bacillus spores neither actively metabolize nor alter the proteins on their surface. Spores have the disadvantage that the molecular mechanisms that trigger sporulation are less well worked out than is the formation of M13 or the export of protein to the outer membrane of E. coli.

Bacteria of the genus Bacillus form endospores that are extremely resistant to damage by heat, radiation,

5
10

15

20

25

30

dipeptides and one isolated K. There are also 20 tyrosines (Y) of which 10 appear as 5 Y-Y dipeptides. Peptides rich in Y and K are known to become crosslinked in oxidizing environments (DEVO78, WAIT83, WAIT85, WAIT86). CotC contains 16 D and E amino acids that nearly equals the 19 Ks. There are no A, F, R, I, L, N, P, Q, S, or W amino acids in CotC. Neither CotC nor CotD is post-translationally cleaved, but the proteins CotA and CotB are.

10 Since, in B. subtilis, some of the spore coat proteins are post-translationally processed by specific proteases, it is valuable to know the sequences of precursors and mature coat proteins so that we can avoid incorporating the recognition sequence of the specific
15 protease into our construction of an OSP-IPBD fusion. The sequence of a mature spore coat protein contains information that causes the protein to be deposited in the spore coat; thus gene fusions that include some or all of a mature coat protein sequence are preferred for screening
20 or selection for the display-of-IPBD phenotype.

Fusions of ipbd fragments to cotC or cotD fragments are likely to cause IPBD to appear on the spore surface. The genes cotC and cotD are preferred osp genes because CotC and CotD are not post-translationally cleaved.
25 Subsequences from cotA or cotB could also be used to cause an IPBD to appear on the surface of B. subtilis spores, but we must take the post-translational cleavage of these proteins into account. DNA encoding IPBD could be fused to a fragment of cotA or cotB at either end of the coding
30 region or at sites interior to the coding region. Spores could then be screened or selected for the display-of-IPBD phenotype.

The promoter of a spore coat protein is most active:
a) when spore coat protein is being synthesized and

5 during sporulation. Ray et al. (RAYC87) have shown that the G4 promoter of B. subtilis is directly controlled by RNA polymerase bound to σ^E . To date, no Bacillus sporulation promoter has been shown to be inducible by an exogenous chemical inducer as the lac promoter of E. coli.
10 Nevertheless, the quantity of protein produced from a sporulation promoter can be controlled by other factors, such as the DNA sequence around the Shine-Dalgarno sequence or codon usage. Chemically inducible sporulation promoters can be developed if necessary.

It is generally preferable to use as the genetic package a cell, spore or virus for which an outer surface protein which can be engineered to display a IPBD has already been identified. However, the present invention is not limited to such genetic packages.

It is believed that the conditions for an outer surface transport signal in a bacterial cell or spore are not particularly stringent, i.e., a random polypeptide of appropriate length (preferably 30-100 amino acids) has a reasonable chance of providing such a signal. Thus, by constructing a chimeric gene comprising a segment encoding the IPBD linked to a segment of random or pseudorandom DNA (the potential OSTs), and placing this gene under control of a suitable promoter, there is a possibility that the chimeric protein so encoded will function as an OSP-IPBD.

This possibility is greatly enhanced by constructing numerous such genes, each having a different potential OSTS, cloning them into a suitable host, and selecting for transformants bearing the IPBD (or other marker) on their

outer surface. Use of secretion-permissive mutants, such as prlA4 (LISS85) or prlG (STAD89), can increase the probability of obtaining a working OSP-IPBD.

When seeking to display a IPBD on the surface of a
 5 bacterial cell, as an alternative to choosing a natural
 OSP and an insertion site in the OSP, we can construct a
 gene (the "display probe") comprising: a) a regulatable
 promoter (e.g. lacUV5), b) a Shine-Dalgarno sequence, c) a
 periplasmic transport signal sequence, d) a fusion of the
 10 ipbd gene with a segment of random DNA (as in Kaiser et
al. (KAIS87)), e) a stop codon, and f) a transcriptional
 terminator.

When the genetic package is a spore, we can use the
 approach described above for attaching a IPBD to an E.
 15 coli cell, except that: a) a sporulation promoter is used,
 and b) no periplasmic signal sequence should be present

For phage, because the OSP-IPBD fulfills a structural
 role in the phage coat, it is unlikely that any particular
 random DNA sequence coupled to the ipbd gene will produce
 20 a fusion protein that fits into the coat in a functional
 way. Nevertheless, random DNA inserted between large
 fragments of a coat protein gene and the pbd gene will
 produce a population that is likely to contain one or more
 members that display the IPBD on the outside of a viable
 25 phage.

As previously stated, the purpose of the random DNA
 is to encode an OSTS, like that embodied in known OSPs.
 The fusion of ipbd and the random DNA could be in either
 order, but ipbd upstream is slightly preferred. Isolates
 30 from the population generated in this way can be screened
 for display of the IPBD. Preferably, a version of
 selection-through-binding is used to select GPs that
 display IPBD on the GP surface. Alternatively, clonal
 isolates of GPs may be screened for the display-of-IPBD

T06230" 56096860

phenotype.

The preference for ipbd upstream of the random DNA arises from consideration of the manner in which the successful GP(IPBD) will be used. The present invention
 5 contemplates introducing numerous mutations into the pbd region of the osp-pbd gene, which, depending on the variegation scheme, might include gratuitous stop codons. If pbd precedes the random DNA, then gratuitous stop codons in pbd lead to no OSP-PBD protein appearing on the
 10 cell surface. If pbd follows the random DNA, then gratuitous stop codons in pbd might lead to incomplete OSP-PBD proteins appearing on the cell surface. Incomplete proteins often are non-specifically sticky so that GPs displaying incomplete PBDs are easily removed from the
 15 population.

The random DNA may be obtained in a variety of ways. Degenerate synthetic DNA is one possibility. Alternatively, pseudorandom DNA can be generated from any DNA having high sequence diversity, e.g., the genome of the organism,
 20 by partially digesting with an enzyme that cuts very often, e.g., Sau3AI. Alternatively, one could shear DNA having high sequence diversity, blunt the sheared DNA with the large fragment of E. coli DNA polymerase I (hereinafter referred to as Klenow fragment), and clone the
 25 sheared and blunted DNA into blunt sites of the vector (MANI82, p295, AUSU87).

If random DNA and phenotypic selection or screening are used to obtain a GP(IPBD), then we clone random DNA into one of the restriction sites that was designed into
 30 the display probe. A plasmid carrying the display probe is digested with the appropriate restriction enzyme and the fragmented, random DNA is annealed and ligated by standard methods. The ligated plasmids are used to transform cells that are grown and selected for expression

0095095-063901

of the antibiotic-resistance gene. Plasmid-bearing GPs are then selected for the display-of-IPBD phenotype by the affinity selection methods described hereafter, using AfM(IPBD) as if it were the target.

- 5 As an alternative to selecting GP(IPBD)s through binding to an affinity column, we can isolate colonies or plaques and screen for successful artificial OSPs through use of one of the methods listed below for verification of the display strategy.

10 IV.F Designing the osp-ipbd gene insert:

Genetic Construction and Expression Considerations

- The (i)pbd-osp gene may be: a) completely synthetic, b) a composite of natural and synthetic DNA, or c) a composite of natural DNA fragments. The important point
15 is that the pbd segment be easily variegated so as to encode a multitudinous and diverse family of PBDs as previously described. A synthetic ipbd segment is preferred because it allows greatest control over placement of restriction sites. Primers complementary to
20 regions abutting the osp-ipbd gene on its 3' flank and to parts of the osp-ipbd gene that are not to be varied are needed for sequencing.

- The sequences of regulatory parts of the gene are taken from the sequences of natural regulatory elements:
25 a) promoters, b) Shine-Dalgarno sequences, and c) transcriptional terminators. Regulatory elements could also be designed from knowledge of consensus sequences of natural regulatory regions. The sequences of these regulatory elements are connected to the coding regions; restriction
30 sites are also inserted in or adjacent to the regulatory regions to allow convenient manipulation.

The essential function of the affinity separation is to separate GPs that bear PBDs (derived from IPBD) having

039605-062904
TOP SECRET

high affinity for the target from GPs bearing PBDs having low affinity for the target. If the elution volume of a GP depends on the number of PBDs on the GP surface, then a GP bearing many PBDs with low affinity, GP(PBD_w), might co-elute with a GP bearing fewer PBDs with high affinity, GP(PBD_s). Regulation of the osp-pbd gene preferably is such that most packages display sufficient PBD to effect a good separation according to affinity. Use of a regulatable promoter to control the level of expression of the osp-pbd allows fine adjustment of the chromatographic behavior of the variegated population.

Induction of synthesis of engineered genes in vegetative bacterial cells has been exercised through the use of regulated promoters such as lacUV5, trpP, or tac (MANI82). The factors that regulate the quantity of protein synthesized include: a) promoter strength (cf. HOOP87), b) rate of initiation of translation (cf. GOLD87), c) codon usage, d) secondary structure of mRNA, including attenuators (cf. LAND87) and terminators (cf. YAGE87), e) interaction of proteins with mRNA (cf. MCPH86, MILL87b, WINT87), f) degradation rates of mRNA (cf. BRAW87, KING86), g) proteolysis (cf. GOTT87). These factors are sufficiently well understood that a wide variety of heterologous proteins can now be produced in E. coli, B. subtilis and other host cells in at least moderate quantities (SKER88, BETT88). Preferably, the promoter for the osp-ipbd gene is subject to regulation by a small chemical inducer. For example, the lac promoter and the hybrid trp-lac (tac) promoter are regulatable with isopropyl thiogalactoside (IPTG). Hereinafter, we use "XINDUCE" as a generic term for a chemical that induces expression of a gene. The promoter for the constructed gene need not come from a natural osp gene; any regulatable bacterial promoter can be used.

Transcriptional regulation of gene expression is best understood and most effective, so we focus our attention on the promoter. If transcription of the osp-ipbd gene is controlled by the chemical XINDUCE, then the number of
 5 OSP-IPBDs per GP increases for increasing concentrations of XINDUCE until a fall-off in the number of viable packages is observed or until sufficient IPBD is observed on the surface of harvested GP(IPBD)s. The attributes that affect the maximum number of OSP-IPBDs per GP are
 10 primarily structural in nature. There may be steric hindrance or other unwanted interactions between IPBDs if OSP-IPBD is substituted for every wild-type OSP. Excessive levels of OSP-IPBD may also adversely affect the solubility or morphogenesis of the GP. For cellular and
 15 viral GPs, as few as five copies of a protein having affinity for another immobilized molecule have resulted in successful affinity separations (FERE82a, FERE82b, and SMIT85).

A non-leaky promoter is preferred. Non-leakiness is
 20 useful: a) to show that affinity of GP(osp-ipbd)s for AfM(IPBD) is due to the osp-ipbd gene, and b) to allow growth of GP(osp-ipbd) in the absence of XINDUCE if the expression of osp-ipbd is disadvantageous. The lacUV5 promoter in conjunction with the LacI^Q repressor is a
 25 preferred example.

An exemplary osp-ipbd gene has the DNA sequence shown in Table 25 and there annotated to explain the useful restriction sites and biologically important features, viz. the lacUV5 promoter, the lacO operator, the Shine-
 30 Dalgarno sequence, the amino acid sequence, the stop codons, and the trp attenuator transcriptional terminator.

The present invention is not limited to a single method of gene design. The osp-ipbd gene need not be synthesized in toto; parts of the gene may be obtained

from nature. One may use any genetic engineering method to produce the correct gene fusion, so long as one can easily and accurately direct mutations to specific sites in the pbd DNA subsequence. In all of the methods of

5 mutagenesis considered in the present invention, however, it is necessary that the coding sequence for the osp-ipbd gene be different from any other DNA in the OCV. The degree and nature of difference needed is determined by the method of mutagenesis to be used. If the method of

10 mutagenesis is to be replacement of subsequences coding for the PBD with vgDNA, then the subsequences to be mutagenized are preferably bounded by restriction sites that are unique with respect to the rest of the OCV. Use of non-unique sites involves partial digestion which is

15 less efficient than complete digestion of a unique site and is not preferred. If single-stranded-oligonucleotide-directed mutagenesis is to be used, then the DNA sequence of the subsequence coding for the IPBD must be unique with respect to the rest of the OCV.

20 The coding portions of genes to be synthesized are designed at the protein level and then encoded in DNA. The amino acid sequences are chosen to achieve various goals, including: a) display of a IPBD on the surface of a GP, b) change of charge on a IPBD, and c) generation of a

25 population of PBDs from which to select an SBD. These issues are discuss in more detail below. The ambiguity in the genetic code is exploited to allow optimal placement of restriction sites and to create various distributions of amino acids at variegated codons.

30 While the invention does not require any particular number or placement of restriction sites, it is generally preferable to engineer restriction sites into the gene to facilitate subsequent manipulations. Preferably, the gene provides a series of fairly uniformly spaced unique

TOP SECRET 5096860

restriction sites with no more than a preset maximum number of bases, for example 100, between sites. Preferably, the gene is designed so that its insertion into the OCV does not destroy the uniqueness of unique restriction sites of the OCV. Preferred recognition sites are those for restriction enzymes which a) generate cohesive ends, b) have unambiguous recognition, or c) have higher specific activity.

The ambiguity of the DNA between the restriction sites is resolved from the following considerations. If the given amino acid sequence occurs in the recipient organism, and if the DNA sequence of the gene in the organism is known, then, preferably, we maximize the differences between the engineered and natural genes to minimize the potential for recombination. In addition, the following codons are poorly translated in E. coli and, therefore, are avoided if possible: cta(L), cga (R), cgg (R), and agg (R). For other host species, different codon restrictions would be appropriate. Finally, long repeats of any one base are prone to mutation and thus are avoided. Balancing these considerations, we can design a DNA sequence.

Structural Considerations

The design of the amino-acid sequence for the ipbd-osp gene to encode involves a number of structural considerations. The design is somewhat different for each type of GP. In bacteria, OSPs are not essential, so there is no requirement that the OSP domain of a fusion have any of its parental functions beyond lodging in the outer membrane.

Relationship between PBD and OSP

It is not required that the PBD and OSP domains have any particular spatial relationship; hence the process of this invention does not require use of the method of US

Patent '692.

It is, in fact, desirable that the OSP not constrain the orientation of the PBD domain; this is not to be confused with lack of constraint within the PBD. Cwirla et al. (CWIR90), Scott and Smith (SCOT90), and Devlin et al. (DEVL90), have taught that variable residues in phage-displayed random peptides should be free of influence from the phage OSP. We teach that binding domains having a moderate to high degree of conformational constraint will exhibit higher specificity and that higher affinity is also possible. Thus, we prescribe picking codons for variegation that specify amino acids that will appear in a well-defined framework. The nature of the side groups is varied through a very wide range due to the combinatorial replacement of multiple amino acids. The main chain conformations of most PBDs of a given class is very similar. The movement of the PBD relative to the OSP should not, however, be restricted. Thus it is often appropriate to include a flexible linker between the PBD and the OSP. Such flexible linkers can be taken from naturally occurring proteins known to have flexible regions. For example, the gIII protein of M13 contains glycine-rich regions thought to allow the amino-terminal domains a high degree of freedom. Such flexible linkers may also be designed. Segments of polypeptides that are rich in the amino acids GLY, ASN, SER, and ASP are likely to give rise to flexibility. Multiple glycines are particularly preferred.

Constraints imposed by OSP

When we choose to insert the PBD into a surface loop of an OSP such as LamB, OmpA, or M13 gIII protein, there are a few considerations that do not arise when PBD is joined to the end of an OSP. In these cases, the OSP exerts some constraining influence on the PBD; the ends of

the PBD are held in more or less fixed positions. We could insert a highly varied DNA sequence into the osp gene at codons that encode a surface-exposed loop and select for cells that have a specific-binding phenotype.

- 5 When the identified amino-acid sequence is synthesized (by any means), the constraint of the OSP is lost and the peptide is likely to have a much lower affinity for the target and a much lower specificity. Tan and Kaiser (TANN77) found that a synthetic model of BPTI containing
 10 all the amino acids of BPTI that contact trypsin has a K_d for trypsin $\approx 10^7$ higher than BPTI. Thus, it is strongly preferred that the varied amino acids be part of a PBD in which the structural constraints are supplied by the PBD.

It is known that the amino acids adjoining foreign
 15 epitopes inserted into LamB influence the immunological properties of these epitopes (VAND90). We expect that PBDs inserted into loops of LamB, OmpA, or similar OSPs will be influenced by the amino acids of the loop and by the OSP in general. To obtain appropriate display of the
 20 PBD, it may be necessary to add one or more linker amino acids between the OSP and the PBD. Such linkers may be taken from natural proteins or designed on the basis of our knowledge of the structural behavior of amino acids. Sequences rich in GLY, SER, ASN, ASP, ARG, and THR are
 25 appropriate. One to five amino acids at either junction are likely to impart the desired degree of flexibility between the OSP and the PBD.

Phage OSP

- A preferred site for insertion of the ipbd gene into
 30 the phage osp gene is one in which: a) the IPBD folds into its original shape, b) the OSP domains fold into their original shapes, and c) there is no interference between the two domains.

If there is a model of the phage that indicates that

FOG290 55096860

either the amino or carboxy terminus of an OSP is exposed to solvent, then the exposed terminus of that mature OSP becomes the prime candidate for insertion of the ipbd gene. A low resolution 3D model suffices.

- 5 In the absence of a 3D structure, the amino and carboxy termini of the mature OSP are the best candidates for insertion of the ipbd gene. A functional fusion may require additional residues between the IPBD and OSP domains to avoid unwanted interactions between the
- 10 domains. Random-sequence DNA or DNA coding for a specific sequence of a protein homologous to the IPBD or OSP, can be inserted between the osp fragment and the ipbd fragment if needed.

- Fusion at a domain boundary within the OSP is also a
- 15 good approach for obtaining a functional fusion. Smith exploited such a boundary when subcloning heterologous DNA into gene III of f1 (SMIT85).

- The criteria for identifying OSP domains suitable for causing display of an IPBD are somewhat different from
- 20 those used to identify and IPBD. When identifying an OSP, minimal size is not so important because the OSP domain will not appear in the final binding molecule nor will we need to synthesize the gene repeatedly in each variegation round. The major design concerns are that: a) the
- 25 OSP::IPBD fusion causes display of IPBD, b) the initial genetic construction be reasonably convenient, and c) the osp::ipbd gene be genetically stable and easily manipulated. There are several methods of identifying domains. Methods that rely on atomic coordinates have been reviewed
- 30 by Janin and Chothia (JANI85). These methods use matrices of distances between α carbons (C_α), dividing planes (cf. ROSE85), or buried surface (RASH84). Chothia and collaborators have correlated the behavior of many natural proteins with domain structure (according to their

definition). Rashin correctly predicted the stability of a domain comprising residues 206-316 of thermolysin (VITA84, RASH84).

Many researchers have used partial proteolysis and protein sequence analysis to isolate and identify stable domains. (See, for example, VITA84, POTE83, SCOT87a, and PABO79.) Pabo et al. used calorimetry as an indicator that the cI repressor from the coliphage λ contains two domains; they then used partial proteolysis to determine the location of the domain boundary.

If the only structural information available is the amino acid sequence of the candidate OSP, we can use the sequence to predict turns and loops. There is a high probability that some of the loops and turns will be correctly predicted (cf. Chou and Fasman, (CHOU74)); these locations are also candidates for insertion of the ipbd gene fragment.

Bacterial OSPs

In bacterial OSPs, the major considerations are: a) that the PBD is displayed, and b) that the chimeric protein not be toxic.

From topological models of OSPs, we can determine whether the amino or carboxy termini of the OSP is exposed. If so, then these are excellent choices for fusion of the osp fragment to the ipbd fragment.

The lamB gene has been sequenced and is available on a variety of plasmids (CLEM81, CHAR88). Numerous fusions of fragments of lamB with a variety of other genes have been used to study export of proteins in E. coli. From various studies, Charbit et al. (CHAR88) have proposed a model that specifies which residues of LamB are: a) embedded in the membrane, b) facing the periplasm, and c) facing the cell surface; we adopt the numbering of this

model for amino acids in the mature protein. According to this model, several loops on the outer surface are defined, including: 1) residues 88 through 111, 2) residues 145 through 165, and 3) 236 through 251.

- 5 Consider a mini-protein embedded in LamB. For example, insertion of DNA encoding $G_1NXCX_5XXXCX_{10}SG_{12}$ between codons 153 and 154 of lamB is likely to lead to a wide variety of LamB derivatives being expressed on the surface of E. coli cells. G_1 , N_2 , S_{11} , and G_{12} are
- 10 supplied to allow the mini-protein sufficient orientation-al freedom that is can interact optimally with the target. Using affinity enrichment (involving, for example, FACS via a fluorescently labeled target, perhaps through several rounds of enrichment), we might obtain a strain
- 15 (named, for example, BEST) that expresses a particular LamB derivative that shows high affinity for the predetermined target. An octapeptide having the sequence of the inserted residues 3 through 10 from BEST is likely to have an affinity and specificity similar to that observed in
- 20 BEST because the octapeptide has an internal structure that keeps the amino acids in a conformation that is quite similar in the LamB derivative and in the isolated mini-protein.

Consideration of the Signal Peptide

- 25 Fusing one or more new domains to a protein may make the ability of the new protein to be exported from the cell different from the ability of the parental protein. The signal peptide of the wild-type coat protein may function for authentic polypeptide but be unable to direct
- 30 export of a fusion. To utilize the Sec-dependent pathway, one may need a different signal peptide. Thus, to express and display a chimeric BPTI/M13 gene VIII protein, we found it necessary to utilize a heterologous signal peptide (that of phoA).

09393095-062903

Provision of a means to remove PBD from the GP

5 GPs that display peptides having high affinity for the target may be quite difficult to elute from the target, particularly a multivalent target. (Bacteria that are bound very tightly can simply multiply in situ.) For phage, one can introduce a cleavage site for a specific protease, such as blood-clotting Factor Xa, into the fusion OSP protein so that the binding domain can be cleaved from the genetic package. Such cleavage has the advantage that all resulting phage have identical OSPs and therefore are equally infective, even if polypeptide-
10 displaying phage can be eluted from the affinity matrix without cleavage. This step allows recovery of valuable genes which might otherwise be lost. To our knowledge, no one has disclosed or suggested using a specific protease as a means to recover an information-containing genetic package or of converting a population of phage that vary in infectivity into phage having identical infectivity.

IV.G. Synthesis of Gene Inserts

20 The present invention is not limited as to how a designed DNA sequence is divided for easy synthesis. An established method is to synthesize both strands of the entire gene in overlapping segments of 20 to 50 nucleotides (nts) (THER88). An alternative method that is more
25 suitable for synthesis of vgdNA is an adaptation of methods published by Oliphant et al. (OLIP86 and OLIP87) and Ausubel et al. (AUSU87). It differs from previous methods in that it: a) uses two synthetic strands, and b) does not cut the extended DNA in the middle. Our goals
30 are: a) to produce longer pieces of dsDNA than can be synthesized as ssDNA on commercial DNA synthesizers, and b) to produce strands complementary to single-stranded vgdNA. By using two synthetic strands, we remove the requirement for a palindromic sequence at the 3' end.

FO6290 5609630

DNA synthesizers can currently produce oligo-nts of lengths up to 200 nts in reasonable yield, $M_{DNA} = 200$. The parameters N_W (the length of overlap needed to obtain efficient annealing) and N_S (the number of spacer bases needed so that a restriction enzyme can cut near the end of blunt-ended dsDNA) are determined by DNA and enzyme chemistry. $N_W = 10$ and $N_S = 5$ are reasonable values. Larger values of N_W and N_S are allowed but add to the length of ssDNA that is to be synthesized and reduce the net length of dsDNA that can be produced.

Let A_L be the actual length of dsDNA to be synthesized, including any spacers. A_L must be no greater than $(2 M_{DNA} - N_W)$. Let Q_W be the number of nts that the overlap window can deviate from center,

$$Q_W = (2 M_{DNA} - N_W - A_L)/2 .$$

Q_W is never negative. It is preferred that the two fragments be approximately the same length so that the amounts synthesized will be approximately equal. This preference may be overridden by other considerations. The overall yield of dsDNA is usually dominated by the synthetic yield of the longer oligo-nt.

We use the following procedure to generate dsDNA of lengths up to $(2 M_{DNA} - N_W)$ nts through the use of Klenow fragment to extend synthetic ss DNA fragments that are not more than M_{DNA} nts long. When a pair of long oligo-nts, complementary for N_W nts at their 3' ends, are annealed there will be a free 3' hydroxyl and a long ssDNA chain continuing in the 5' direction on either side. We will refer to this situation as a 5' superoverhang. The procedure comprises:

- 1) picking a non-palindromic subsequence of N_W to N_W+4 nts near the center of the dsDNA to be synthesized;

this region is called the overlap (typically, N_w is 10),

- 2) synthesizing a ss DNA molecule that comprises that part of the anti-sense strand from its 5' end up to and including the overlap,
- 3) synthesizing a ss DNA molecule that comprises that part of the sense strand from its 5' end up to and including the overlap,
- 4) annealing the two synthetic strands that are complementary throughout the overlap region, and
- 5) extending both superoverhangs with Klenow fragment and all four deoxynucleotide triphosphates.

Because M_{DNA} is not rigidly fixed at 200, the current limits of 390 ($= 2 M_{DNA} - N_w$) nts overall and 200 in each fragment are not rigid, but can be exceeded by 5 or 10 nts. Going beyond the limits of 390 and 200 will lead to lower yields, but these may be acceptable in certain cases.

Restriction enzymes do not cut well at sites closer than about five base pairs from the end of blunt ds DNA fragments (OLIP87 and p.132 New England BioLabs 1990-1991 Catalogue). Therefore N_s nts (with N_s typically set to 5) of spacer are added to ends that we intend to cut with a restriction enzyme. If the plasmid is to be cut with a blunt-cutting enzyme, then we do not add any spacer to the corresponding end of the ds DNA fragment.

To choose the optimum site of overlap for the oligo-nt fragments, first consider the anti-sense strand of the DNA to be synthesized, including any spacers at the ends, written (in upper case) from 5' to 3' and left-to-right.

N.B.: The N_w nt long overlap window can never include bases that are to be variegated. N.B.: The N_w nt long overlap should not be palindromic lest single DNA molecules prime themselves. Place a N_w nt long window as

20 Underscore the anti-sense strand from the 5' end up to the right edge of the window. Write the complementary sense sequence 3'-to-5' and left-to-right and in lower case letters, under the anti-sense strand starting at the left edge of the window and continuing all the way to the right end of the anti-sense strand.

The present invention is not limited to any parti-

cular method of DNA synthesis or construction. Conventional DNA synthesizers may be used, with appropriate reagent modifications for production of variegated DNA (similar to that now used for production of mixed probes).

- 5 For example, the Milligen 7500 DNA synthesizer has seven vials from which phosphoramidites may be taken. Normally, the first four contain A, C, T, and G. The other three vials may contain unusual bases such as inosine or mixtures of bases, the so-called "dirty bottle". The
10 standard software allows programmed mixing of two, three, or four bases in equimolar quantities.

The synthesized DNA may be purified by any art recognized technique, e.g., by high-pressure liquid chromatography (HPLC) or PAGE.

- 15 The osp-pbd genes may be created by inserting vgDNA into an existing parental gene, such as the osp-ipbd shown to be displayable by a suitably transformed GP. The present invention is not limited to any particular method of introducing the vgDNA, however, two techniques are
20 discussed below.

- In the case of cassette mutagenesis, the restriction sites that were introduced when the gene for the inserted domain was synthesized are used to introduce the synthetic vgDNA into a plasmid or other OCV. Restriction digestions
25 and ligations are performed by standard methods (AUSU87).

In the case of single-stranded-oligonucleotide-directed mutagenesis, synthetic vgDNA is used to create diversity in the vector (BOTS85).

- The modes of creating diversity in the population of
30 GPs discussed herein are not the only modes possible. Any method of mutagenesis that preserves at least a large fraction of the information obtained from one selection and then introduces other mutations in the same domain will work. The limiting factors are the number of

independent transformants that can be produced and the amount of enrichment one can achieve through affinity separation. Therefore the preferred embodiment uses a method of mutagenesis that focuses mutations into those
 5 residues that are most likely to affect the binding properties of the PBD and are least likely to destroy the underlying structure of the IPBD.

Other modes of mutagenesis might allow other GPs to be considered. For example, the bacteriophage λ is not a
 10 useful cloning vehicle for cassette mutagenesis because of the plethora of restriction sites. One can, however, use single-stranded-oligo-nt-directed mutagenesis on λ without the need for unique restriction sites. No one has used single-stranded-oligo-nt-directed mutagenesis to
 15 introduce the high level of diversity called for in the present invention, but if it is possible, such a method would allow use of phage with large genomes.

IV.H. Operative Cloning Vector

The operative cloning vector (OCV) is a replicable
 20 nucleic acid used to introduce the chimeric ipbd-osp or ipbd-osp gene into the genetic package. When the genetic package is a virus, it may serve as its own OCV. For cells and spores, the OCV may be a plasmid, a virus, a phagemid, or a chromosome.

25 The OCV is preferably small (less than 10 KB), stable (even after insertion of at least 1 kb DNA), present in multiple copies within the host cell, and selectable with appropriate media. It is desirable that cassette mutagenesis be practical in the OCV; preferably, at least 25
 30 restriction enzymes are available that do not cut the OCV. It is likewise desirable that single-stranded mutagenesis be practical. If a suitable OCV does not already exist, it may be engineered by manipulation of available vectors.

FOIA b 7 - 56056860

5

10

20

25

present invention. Phagemids may be entirely suitable for developing a gene that causes an IPBD to appear on the surface of phage-like genetic packages.

It is also well known that plasmids containing the
 5 ColE1 origin of replication can be greatly amplified if protein synthesis is halted in a log-phase culture. Protein synthesis can be halted by addition of chloramphenicol or other agents (MANI82).

The bacteriophage M13 bla 61 (ATCC 37039) is derived
 10 from wild-type M13 through the insertion of the β lactamase gene (HINE80). This phage contains 8.13 kb of DNA. M13 bla cat 1 (ATCC 37040) is derived from M13 bla 61 through the additional insertion of the chloramphenicol resistance gene (HINE80); M13 bla cat 1 contains 9.88 kb
 15 of DNA. Although neither of these variants of M13 contains the ColE1 origin of replication, either could be used as a starting point to construct a cloning vector with this feature.

IV.I. Transformation of cells:

20 When the GP is a cell, the population of GPs is created by transforming the cells with suitable OCVs. When the GP is a phage, the phage are genetically engineered and then transfected into host cells suitable for amplification. When the GP is a spore, cells capable of
 25 sporulation are transformed with the OCV while in a normal metabolic state, and then sporulation is induced so as to cause the OSP-PBDs to be displayed. The present invention is not limited to any one method of transforming cells with DNA. The procedure given in the examples is a
 30 modification of that of Maniatis (p250, MANI82). One preferably obtains at least 10^7 and more preferably at least 10^8 transformants/ μ g of CCC DNA.

The transformed cells are grown first under non-

T06230-5606860

selective conditions that allow expression of plasmid genes and then selected to kill untransformed cells. Transformed cells are then induced to express the osp-pbd gene at the appropriate level of induction. The GPs
 5 carrying the IPBD or PBDs are then harvested by methods appropriate to the GP at hand, generally, centrifugation to pelletize GPs and resuspension of the pellets in sterile medium (cells) or buffer (spores or phage). They are then ready for verification that the display strategy
 10 was successful (where the GPs all display a "test" IPBD) or for affinity selection (where the GPs display a variety of different PBDs).

IV.J. Verification of Display Strategy:

The harvested packages are tested to determine
 15 whether the IPBD is present on the surface. In any tests of GPs for the presence of IPBD on the GP surface, any ions or cofactors known to be essential for the stability of IPBD or AfM(IPBD) are included at appropriate levels. The tests can be done: a) by affinity labeling, b)
 20 enzymatically, c) spectrophotometrically, d) by affinity separation, or e) by affinity precipitation. The AfM(IPBD) in this step is one picked to have strong affinity (preferably, $K_d < 10^{-11}$ M) for the IPBD molecule and little or no affinity for the wtGP. For example, if BPTI
 25 were the IPBD, trypsin, anhydrotrypsin, or antibodies to BPTI could be used as the AfM(BPTI) to test for the presence of BPTI. Anhydrotrypsin, a trypsin derivative with serine 195 converted to dehydroalanine, has no proteolytic activity but retains its affinity for BPTI
 30 (AKOH72 and HUBE77).

Preferably, the presence of the IPBD on the surface of the GP is demonstrated through the use of a soluble, labeled derivative of a AfM(IPBD) with high affinity for IPBD. The label could be: a) a radioactive atom such as

TOP SECRET S6096360

5 1) mix AfM(IPBD)* with GPs that are to be tested for the
presence of IPBD; conditions of mixing should favor
binding of IPBD to AfM(IPBD)*,
2) separate GPs from unbound AfM(IPBD)* by use of:
10 a) a molecular sizing filter that will pass
AfM(IPBD)* but not GPs,
b) centrifugation, or
c) a molecular sizing column (such as Sepharose or
Sephadex) that retains free AfM(IPBD)* but not
15 GPs,
3) quantitate the AfM(IPBD)* bound by GPs.

If the IPBD has strong, characteristic absorption bands in the visible or UV that are distinct from absorption by the wtGP, then another alternative for measuring the IPBD displayed on the GP is a spectrophotometric measurement. For example, if IPBD were azurin, the visible absorption could be used to identify GPs that display azurin.

Another alternative is to label the GPs and measure
30 the amount of label retained by immobilized AfM(IPBD).
For example, the GPs could be grown with a radioactive
precursor, such as ^{32}P or ^3H -thymidine, and the radioac-
tivity retained by immobilized AfM(IPBD) measured.

Another alternative is to use affinity chromatography; the ability of a GP bearing the IPBD to bind a matrix that supports a AfM(IPBD) is measured by reference to the wtGP.

- 5 Another alternative for detecting the presence of IPBD on the GP surface is affinity precipitation.

If random DNA has been used, then affinity selection procedures are used to obtain a clonal isolate that has the display-of-IPBD phenotype. Alternatively, clonal
10 isolates may be screened for the display-of-IPBD phenotype. The tests of this step are applied to one or more of these clonal isolates.

If no isolates that bind to the affinity molecule are obtained we take corrective action as disclosed below.

- 15 If one or more of the tests above indicates that the IPBD is displayed on the GP surface, we verify that the binding of molecules having known affinity for IPBD is due to the chimeric osp-ipbd gene through the use of standard genetic and biochemical techniques, such as:

- 20 1) transferring the osp-ipbd gene into the parent GP to verify that osp-ipbd confers binding,
2) deleting the osp-ipbd gene from the isolated GP to verify that loss of osp-ipbd causes loss of binding,
3) showing that binding of GPs to AfM(IPBD) correlates
25 with [XINDUCE] (in those cases that expression of osp-ipbd is controlled by [XINDUCE]), and
4) showing that binding of GPs to AfM(IPBD) is specific to the immobilized AfM(IPBD) and not to the support matrix.

- 30 Variation of: a) binding of GPs by soluble AfM(IPBD-)*, b) absorption caused by IPBD, and c) biochemical reactions of IPBD are linear in the amount of IPBD displayed. Presence of IPBD on the GP surface is indi-

cated by a strong correlation between [XINDUCE] and the reactions that are linear in the amount of IPBD. Leakiness of the promoter is not likely to present problems of high background with assays that are linear in the amount of IPBD. These experiments may be quicker and easier than the genetic tests. Interpreting the effect of [XINDUCE] on binding to a {AfM(IPBD)} column, however, may be problematic unless the regulated promoter is completely repressed in the absence of [XINDUCE]. The affinity retention of GP(IPBD)s is not linear in the number of IPBDs/GP and there may be, for example, little phenotypic difference between GPs bearing 5 IPBDs and GPs bearing 50 IPBDs. The demonstration that binding is to AfM(IPBD) and the genetic tests are essential; the tests with XINDUCE are optional.

We sequence the relevant ipbd gene fragment from each of several clonal isolates to determine the construction. We also establish the maximum salt concentration and pH range for which the GP(IPBD) binds the chosen AfM(IPBD). This is preferably done by measuring, as a function of salt concentration and pH, the retention of AfM(IPBD)* on molecular sizing filters that pass AfM(IPBD)* but not GP. This information will be used in refining the affinity selection scheme.

25 IV.K. Analysis and Correction of Display Problems

If the IPBD is displayed on the outside of the GP, and if that display is clearly caused by the introduced osp-ipbd gene, we proceed with variegation, otherwise we analyze the result and adopt appropriate corrective measures. If we have unsuccessfully attempted to fuse an ipbd fragment to a natural osp fragment, our options are :1) pick a different fusion to the same osp by a) using opposite end of osp, b) keeping more or fewer residues from osp in the fusion; for example, in increments of 3 or

4 residues, c) trying a known or predicted domain boundary, d) trying a predicted loop or turn position, 2) pick a different osp, or 3) switch to random DNA method. If we have just tried the random DNA method unsuccessfully, our options are: 1) choose a different relationship between ipbd fragment and random DNA (ipbd first, random DNA second or vice versa), 2) try a different degree of partial digestion, a different enzyme for partial digestion, a different degree of shearing or a different source of natural DNA, or 3) switch to the natural OSP method. If all reasonable OSPs of the current GP have been tried and the random DNA method has been tried, both without success, we pick a new GP.

We may illustrate the ways in which problems may be attacked by using the example of BPTI as the IPBD, the M13 phage as the GP, and the major coat (gene VIII) protein as the OSP. The following amino-acid sequence, called AA_seq2, illustrates how the sequence for mature BPTI (shown underscored) may be inserted immediately after the signal sequence of M13 precoat protein (indicated by the arrow) and before the sequence for the M13 CP.

AA_seq2

25		1	1	2	2	3	3	4	4	5
	5	0	5	0	5	0	5	0	5	0
	<u>MKKSILVLKASVAVATLVPMLSFARPDFCLEPPYTGPCKARIIRYFYNAKA</u>									
	5	6	6	7	7	8	8	9	9	10
30	5	0	5	0	5	0	5	0	5	0
	<u>GLCOTFVYGGCRAKRNNFKSAEDCMRTCGGAAEGDDPAKA AFNSLQASAT</u>									
	10	11	11	12	12	13				
	5	0	5	0	5	0				
35	EYIGYAWAMVVVIVGATIGIKLFKKFTSKAS									

We adopt the convention that sequence numbers of fusion proteins refer to the fusion, as coded, unless otherwise noted. Thus the alanine that begins M13 CP is referred to as "number 82", "number 1 of M13 CP", or

"number 59 of the mature BPTI-M13 CP fusion".

It is desirable to determine where, exactly, the BPTI binding domain is being transported: is it remaining in the cytoplasm? Is it free within the periplasm? Is it
 5 attached to the inner membrane? Proteins in the periplasm can be freed through spheroplast formation using lysozyme and EDTA in a concentrated sucrose solution (BIRD67, MALA64). If BPTI were free in the periplasm, it would be found in the supernatant. Trypsin labeled with
 10 ^{125}I would be mixed with supernatant and passed over a non-denaturing molecular sizing column and the radioactive fractions collected. The radioactive fractions would then be analyzed by SDS-PAGE and examined for BPTI-sized bands by silver staining.

15 Spheroplast formation exposes proteins anchored in the inner membrane. Spheroplasts would be mixed with AHTrp* and then either filtered or centrifuged to separate them from unbound AHTrp*. After washing with hypertonic buffer, the spheroplasts would be analyzed for extent of
 20 AHTrp* binding.

If BPTI were found free in the periplasm, then we would expect that the chimeric protein was being cleaved both between BPTI and the M13 mature coat sequence and between BPTI and the signal sequence. In that case, we
 25 should alter the BPTI/M13 CP junction by inserting vgDNA at codons for residues 78-82 of AA_seq2.

If BPTI were found attached to the inner membrane, then two hypotheses can be formed. The first is that the chimeric protein is being cut after the signal sequence,
 30 but is not being incorporated into LG7 virion; the treatment would also be to insert vgDNA between residues 78 and 82 of AA_seq2. The alternative hypothesis is that BPTI could fold and react with trypsin even if signal sequence is not cleaved. N-terminal amino acid sequencing

T06230-5605660

of trypsin-binding material isolated from cell homogenate determines what processing is occurring. If signal sequence were being cleaved, we would use the procedure above to vary residues between C78 and A82; subsequent
 5 passes would add residues after residue 81. If signal sequence were not being cleaved, we would vary residues between 23 and 27 of AA_seq2. Subsequent passes through that process would add residues after 23.

If BPTI were found neither in the periplasm nor on
 10 the inner membrane, then we would expect that the fault was in the signal sequence or the signal-sequence-to-BPTI junction. The treatment in this case would be to vary residues between 23 and 27.

Analytical experiments to determine what has gone
 15 wrong take time and effort and, for the foreseen outcomes, indicate variations in only two regions. Therefore, we believe it prudent to try the synthetic experiments described below without doing the analysis. For example, these six experiments that introduce variegation into the
 20 bpti-gene VIII fusion could be tried:

- 1) 3 variegated codons between residues 78 and 82 using olig#12 and olig#13,
- 2) 3 variegated codons between residues 23 and 27 using olig#14 and olig#15,
- 25 3) 5 variegated codons between residues 78 and 82 using olig#13 and olig#12a,
- 4) 5 variegated codons between residues 23 and 27 using olig#15 and olig#14a,
- 5) 7 variegated codons between residues 78 and 82 using
 30 olig#13 and olig#12b, and
- 6) 7 variegated codons between residues 23 and 27 using olig#15 and olig#14b.

To alter the BPTI-M13 CP junction, we introduce DNA variegated at codons for residues between 78 and 82 into

T05250-56096860

the SphI and SfiI sites of pLG7. The residues after the last cysteine are highly variable in amino acid sequences homologous to BPTI, both in composition and length; in Table 25 these residues are denoted as G79, G80, and A81.

- 5 The first part of the M13 CP is denoted as A82, E83, and G84. One of the oligo-nts olig#12, olig#12a, or olig#12b and the primer olig#13 are synthesized by standard methods. The oligo-nts are:

```

10      residue 75 76 77 78 79 80 81 82 83
      5' gc|gag|cgc|ATG|CGT|ACC|TGC|qfk|qfk|qfk|GCT|GAA|-
      84 85 86 87 88 89 90 91
      GGT|GAT|GAT|CCG|GCC|AAA|GCG|GCC|gcg|cc 3' olig#12
15
      residue 75 76 77 78 79 80 81 81a 81b
      5' gc|gag|cgc|ATG|CGT|ACC|TGC|qfk|qfk|qfk|qfk|qfk|-
20
      82 83 84 85 86 87
      GCT|GAA|GGT|GAT|GAT|CCG|-
      88 89 90 91
      GCC|AAA|GCG|GCC|gcg|cc 3' olig#12a
25
      residue 75 76 77 78 79 80 81 81a 81b
      5' gc|gag|cgc|ATG|CGT|ACC|TGC|qfk|qfk|qfk|qfk|qfk|-
      81c 81d 82 83 84 85 86 87
      qfk|qfk|GCT|GAA|GGT|GAT|GAT|CCG|-
30
      88 89 90 91
      GCC|AAA|GCG|GCC|gcg|cc 3' olig#12b
35
      residue 91 90 89 88 87 86
      5' gg|cgc|GGC|CGC|TTT|GGC|CGG|ATC 3' olig#13

```

- where q is a mixture of (0.26 T, 0.18C, 0.26 A, and 0.30 G), f is a mixture of (0.22 T, 0.16 C, 0.40 A, and 0.22 G), and k is a mixture of equal parts of T and G. The bases shown in lower case at either end are spacers and are not incorporated into the cloned gene. The primer is complementary to the 3' end of each of the longer oligo-nts. One of the variegated oligo-nts and the primer
- 45 olig#13 are combined in equimolar amounts and annealed.

The dsDNA is completed with all four (nt)TPs and Klenow fragment. The resulting dsDNA and RF pLG7 are cut with both SfiI and SphI, purified, mixed, and ligated. We then select a transformed clone that, when induced with IPTG, 5 binds AHTrp.

To vary the junction between M13 signal sequence and BPTI, we introduce DNA variegated at codons for residues between 23 and 27 into the KpnI and XhoI sites of pLG7. The first three residues are highly variable in amino acid sequences homologous to BPTI. Homologous sequences also vary in length at the amino terminus. One of the oligo-nts olig#14, olig#14a, or olig#14b and the primer olig#15 are synthesized by standard methods. The oligo-nts are:

15 residue : 17 18 19 20 21 22 23 24 25
5' g|gcc|gcG|GTA|CCG|ATG|CTG|TCT|TTT|GCT|fxk|fxk|-

20 26 27 28 29 30
|fxk|TTC|TGT|CTC|GAG|cgc|ccg|cga| 3' olig#14

25 residue 17 18 19 20 21 22 23 24 25 26
5'g|gcc|gcG|GTA|CCG|ATG|CTG|TCT|TTT|GCT|fxk|fxk|fxk|-

26a 26b 27 28 29 30
|fxk|fxk|TTC|TGT|CTC|GAG|cgc|ccg|cga| 3' olig#14a,

30 residue 17 18 19 20 21 22 23 24 25 26
5'g|gcc|gcG|GTA|CCG|ATG|CTG|TCT|TTT|GCT|fxk|fxk|fxk|-

26a 26b 26c 26d 27 28 29 30
|fxk|fxk|fxk|fxk|TTC|TGT|CTC|GAG|cgc|ccg|cga| 3' olig#14b

35

5' |tcg|cgg|gcg|CTC|GAG|ACA|GAA| 3' olig#15

40 where f is a mixture of (0.26 T, 0.18 C, 0.26 A, and 0.30 G), x is a mixture of (0.22 T, 0.16 C, 0.40 A, and 0.22 G), and k is a mixture of equal parts of T and G. The bases shown in lower case at either end are spacers and are not incorporated into the cloned gene. One of the

099600-509600

variegated oligo-nts and the primer are combined in equimolar amounts and annealed. The ds DNA is completed with all four (nt)TPs and Klenow fragment. The resulting dsDNA and RF pLG7 are cut with both KpnI and XhoI,
 5 purified, mixed, and ligated. We select a transformed clone that, when induced with IPTG, binds AHTrp or trp.

Other numbers of variegated codons could be used.

If none of these approaches produces a working chimeric protein, we may try a different signal sequence.
 10 If that doesn't work, we may try a different OSP.

V. AFFINITY SELECTION OF TARGET-BINDING MUTANTS

V.A. Affinity Separation Technology, Generally

Affinity separation is used initially in the present invention to verify that the display system is working,
 15 i.e., that a chimeric outer surface protein has been expressed and transported to the surface of the genetic package and is oriented so that the inserted binding domain is accessible to target material. When used for this purpose, the binding domain is a known binding domain
 20 for a particular target and that target is the affinity molecule used in the affinity separation process. For example, a display system may be validated by using inserting DNA encoding BPTI into a gene encoding an outer surface protein of the genetic package of interest, and
 25 testing for binding to anhydrotrypsin, which is normally bound by BPTI.

If the genetic packages bind to the target, then we have confirmation that the corresponding binding domain is indeed displayed by the genetic package. Packages which
 30 display the binding domain (and thereby bind the target) are separated from those which do not.

Once the display system is validated, it is possible to use a variegated population of genetic packages which

T06290"55096860

display a variety of different potential binding domains, and use affinity separation technology to determine how well they bind to one or more targets. This target need not be one bound by a known binding domain which is parental to the displayed binding domains, i.e., one may select for binding to a new target.

For example, one may variegate a BPTI binding domain and test for binding, not to trypsin, but to another serine protease, such as human neutrophil elastase or cathepsin G, or even to a wholly unrelated target, such as horse heart myoglobin.

The term "affinity separation means" includes, but is not limited to: a) affinity column chromatography, b) batch elution from an affinity matrix material, c) batch elution from an affinity material attached to a plate, d) fluorescence activated cell sorting, and e) electrophoresis in the presence of target material. "Affinity material" is used to mean a material with affinity for the material to be purified, called the "analyte". In most cases, the association of the affinity material and the analyte is reversible so that the analyte can be freed from the affinity material once the impurities are washed away.

The procedures described in sections V.H, V.I and V.J are not required for practicing the present invention, but may facilitate the development of novel binding proteins thereby.

V.B. Affinity Chromatography, Generally

Affinity column chromatography, batch elution from an affinity matrix material held in some container, and batch elution from a plate are very similar and hereinafter will be treated under "affinity chromatography."

If affinity chromatography is to be used, then:

- 1) the molecules of the target material must be of sufficient size and chemical reactivity to be applied to a solid support suitable for affinity separation,
- 2) after application to a matrix, the target material preferably does not react with water,
- 3) after application to a matrix, the target material preferably does not bind or degrade proteins in a non-specific way, and
- 4) the molecules of the target material must be sufficiently large that attaching the material to a matrix allows enough unaltered surface area (generally at least 500 Å², excluding the atom that is connected to the linker) for protein binding.

Affinity chromatography is the preferred separation means, but FACS, electrophoresis, or other means may also be used.

V.C. Fluorescent-Activated Cell Sorting. Generally

Fluorescent-activated cell sorting involves use of an affinity material that is fluorescent per se or is labeled with a fluorescent molecule. Current commercially available cell sorters require 800 to 1000 molecules of fluorescent dye, such as Texas red, bound to each cell. FACS can sort 10³ cells or viruses/sec.

FACS (e.g. FACStar from Beckton-Dickinson, Mountain View, CA) is most appropriate for bacterial cells and spores because the sensitivity of the machines requires approximately 1000 molecules of fluorescent label bound to each GP to accomplish a separation. OSPs such as OmpA, OmpF, OmpC are present at $\geq 10^4$ /cell, often as much as 10⁵/cell. Thus use of FACS with PBDs displayed on one of the OSPs of a bacterial cell is attractive. This is particularly true if the target is quite small so that attachment to a matrix has a much greater effect than would attachment to a dye. To optimize FACS separation of

GPs, we use a derivative of Afm(IPBD) that is labeled with
 a fluorescent molecule, denoted Afm(IPBD)*. The vari-
 ables to be optimized include: a) amount of IPBD/GP, b)
 concentration of Afm(IPBD)*, c) ionic strength, d)
 5 concentration of GPs, and e) parameters pertaining to
 operation of the FACS machine. Because Afm(IPBD)* and GPs
 interact in solution, the binding will be linear in both
 [Afm(IPBD)*] and [displayed IPBD]. Preferably, these two
 parameters are varied together. The other parameters can
 10 be optimized independently.

If FACS is to be used as the affinity separation means, then:

- 1) the molecules of the target material must be of
 sufficient size and chemical reactivity to be
 15 conjugated to a suitable fluorescent dye or the
 target must itself be fluorescent,
- 2) after any necessary fluorescent labeling, the target
 preferably does not react with water,
- 3) after any necessary fluorescent labeling, the target
 20 material preferably does not bind or degrade proteins
 in a non-specific way, and
- 4) the molecules of the target material must be suffi-
 ciently large that attaching the material to a
 suitable dye allows enough unaltered surface area
 25 (generally at least 500 Å², excluding the atom that
 is connected to the linker) for protein binding.

V.D. Affinity Electrophoresis, Generally

Electrophoretic affinity separation involves electro-
 phoresis of viruses or cells in the presence of target
 30 material, wherein the binding of said target material
 changes the net charge of the virus particles or cells.
 It has been used to separate bacteriophages on the basis
 of charge. (SERW87).

Electrophoresis is most appropriate to bacteriophage -

T06290-56096860

5 For example, chloroacetate ions contain only seven atoms and would be essentially altered by any linkage. GPs that bind chloroacetate would become more negatively charged than GPs that do not bind the ion and so these classes of GPs could be separated.

- 1) the target must either be charged or of such a nature that its binding to a protein will change the charge of the protein,
- 2) the target material preferably does not react with water,
- 3) the target material preferably does not bind or degrade proteins in a non-specific way, and
- 4) the target must be compatible with a suitable gel material.

20 The present invention makes use of affinity separation of bacterial cells, or bacterial viruses (or other genetic packages) to enrich a population for those cells or viruses carrying genes that code for proteins with
25 desirable binding properties.

V.E. Target Materials

The present invention may be used to select for binding domains which bind to one or more target materials, and/or fail to bind to one or more target materials. Specificity, of course, is the ability of a binding molecule to bind strongly to a limited set of target materials, while binding more weakly or not at all to another set of target materials from which the first set

must be distinguished.

The target materials may be organic macromolecules, such as polypeptides, lipids, polynucleic acids, and polysaccharides, but are not so limited. Almost any molecule that is stable in aqueous solvent may be used as a target. The following list of possible targets is given as illustration and not as limitation. The categories are not strictly mutually exclusive. The omission of any category is not to be construed to imply that said category is unsuitable as a target.

A. Peptides

- 1) human β endorphin (Merck Index 3528)
- 2) dynorphin (MI 3458)
- 3) Substance P (MI 8834)
- 4) Porcine somatostatin (MI 8671)
- 5) human atrial natriuretic factor (MI 887)
- 6) human calcitonin
- 7) glucagon

B. Proteins

I. Soluble Proteins

a. Hormones

- 1) human TNF (MI 9411)
- 2) Interleukin-1 (MI 4895)
- 3) Interferon- γ (MI 4894)
- 4) Thyrotropin (MI 9709)
- 5) Interferon- α (MI 4892)
- 6) Insulin (MI 4887, p.789)

b. Enzymes

- 1) human neutrophil elastase
- 2) Human thrombin
- 3) human Cathepsin G
- 4) human tryptase
- 5) human chymase
- 6) human blood clotting Factor Xa

TOP SECRET

- 7) any retro-viral Pol protease
 8) any retro-viral Gag protease
 9) dihydrofolate reductase
 10) Pseudomonas putida cytochrome P450_{CAM}
 5 11) human pyruvate kinase
 12) E. coli pyruvate kinase
 13) jack bean urease
 14) aspartate transcarbamylase (E. coli)
 15) ras protein
 10 16) any protein-tyrosine kinase
- c. Inhibitors
- 1) aprotinin (MI 784)
 2) human α 1-anti-trypsin
 3) phage λ cI (inhibits DNA transcription)
- 15 d. Receptors
- 1) TNF receptor
 2) IgE receptor
 3) LamB
 4) CD4
 20 5) IL-1 receptor
- e. Toxins
- 1) ricin (also an enzyme)
 2) α Conotoxin GI
 3) mellitin
 25 4) Bordetella pertussis adenylate cyclase
 (also an enzyme)
 5) Pseudomonas aeruginosa hemolysin
- f. Other proteins
- 1) horse heart myoglobin
 30 2) human sickle-cell haemoglobin
 3) human deoxy haemoglobin
 4) human CO haemoglobin
 5) human low-density lipoprotein (a lipoprotein)
 35 6) human IgG (combining site removed or

T06290"56096860

5

- 10

15

- ### C. Nucleic acids

20

- 25

35

- 1) yeast Phe tRNA
- 2) ribosomal RNA
- 3) segment of mRNA

D. Organic molecules (not peptide, protein, or nucleic acid)

I. Small and monomeric

- 5 1) cholesterol
- 2) aspartame
- 3) bilirubin
- 4) morphine
- 5) codeine
- 10 6) heroine
- 7) dichlorodiphenyltrichlorethane (DDT)
- 8) prostaglandin PGE2
- 9) actinomycin
- 10) 2,2,3 trimethyldecane
- 15 11) Buckminsterfullerene
- 12) cortavazol (MI 2536, p.397)

II. Polymers

- 1) cellulose
- 2) chitin

III. Others

- 1) O-antigen of Salmonella enteritidis (a lipopolysaccharide)

E. Inorganic compounds

- 1) asbestos
- 25 2) zeolites
- 3) hydroxylapatite
- 4) 111 face of crystalline silicon
- 5) paulingite
- 6) U(IV) (uranium ions)
- 30 7) Au(III) (gold ions)

F. Organometallic compounds

- 1) iron(III) haem
- 2) cobalt haem
- 3) cobalamine
- 35 4) (isopropylamino)₆Cr(III)

T06290-55096860

15



25

30

35

HNE is not the only serine protease for which an inhibitor would be valuable. Works concerning uses of protease inhibitors and diseases thought to result from inappropriate protease activity include: NADE87, REST88, 5 SOMM90, and SOMM89. Trypsin and chymase may be involved in asthma, see FRAN89 and VAND89. There are reports that suggest that Proteinase 3 (also known as p29) is as important or even more important than HNE; see NILE89, ARNA90, KAOR88, CAMP90, and GUPT90. Cathepsin G is 10 another protease that may cause disease when present in excess; see FERR90, PETE89, SALV87, and SOMM90. These works indicate that a problem exists and that blocking one or another protease might well alleviate a disease state. Some of the cited works report inhibitors having measur- 15 able affinity for a target protease, but none report truly excellent inhibitors that have K_d in the range of 10^{-12} M as may be obtained by the method of the present invention. The same IPBDs used for HNE can be used for any serine protease.

20 The present invention is not, however, limited to any of the above-identified target materials. The only limitation is that the target material be suitable for affinity separation.

A supply of several milligrams of pure target 25 material is desired. With HNE (as discussed in Examples II and III), 400 μ g of enzyme is used to prepare 200 μ l of ReactiGel beads. This amount of beads is sufficient for as many as 40 fractionations. Impure target material could be used, but one might obtain a protein that binds

09895095 062904

to a contaminant instead of to the target.

The following information about the target material is highly desirable: 1) stability as a function of temperature, pH, and ionic strength, 2) stability with respect to chaotropes such as urea or guanidinium Cl, 3) pI, 4) molecular weight, 5) requirements for prosthetic groups or ions, such as haem or Ca^{+2} , and 6) proteolytic activity, if any. It is also potentially useful to know: 1) the target's sequence, if the target is a macro-molecule, 2) the 3D structure of the target, 3) enzymatic activity, if any, and 4) toxicity, if any.

The user of the present invention specifies certain parameters of the intended use of the binding protein: 1) the acceptable temperature range, 2) the acceptable pH range, 3) the acceptable concentrations of ions and neutral solutes, and 4) the maximum acceptable dissociation constant for the target and the SBD:

$$K_T = [\text{Target}][\text{SBD}]/[\text{Target:SBD}].$$

In some cases, the user may require discrimination between T, the target, and N, some non-target. Let

$$K_T = [\text{T}][\text{SBD}]/[\text{T:SBD}], \text{ and}$$

$$K_N = [\text{N}][\text{SBD}]/[\text{N:SBD}],$$

$$\text{then } K_T/K_N = ([\text{T}][\text{N:SBD}])/([\text{N}][\text{T:SBD}]).$$

The user then specifies a maximum acceptable value for the ratio K_T/K_N .

The target material preferably is stable under the specified conditions of pH, temperature, and solution conditions.

If the target material is a protease, one considers the following points:

- 1) a highly specific protease can be treated like any other target,
- 2) a general protease, such as subtilisin, may degrade

- the OSPs of the GP including OSP-PBDs; there are several alternative ways of dealing with general proteases, including: a) use a protease inhibitor as PPBD so that the SBD is an inhibitor of the protease,
- 5 b) a chemical inhibitor may be used to prevent proteolysis (e.g. phenylmethylfluorosulfate (PMFS) that inhibits serine proteases), c) one or more active-site residues may be mutated to create an inactive protein (e.g. a serine protease in which the
- 10 active serine is mutated to alanine), or d) one or more active-site amino-acids of the protein may be chemically modified to destroy the catalytic activity (e.g. a serine protease in which the active serine is converted to anhydroserine),
- 15 3) SBDs selected for binding to a protease need not be inhibitors; SBDs that happen to inhibit the protease target are a fairly small subset of SBDs that bind to the protease target,
- 4) the more we modify the target protease, the less like
- 20 we are to obtain an SBD that inhibits the target protease, and
- 5) if the user requires that the SBD inhibit the target protease, then the active site of the target protease must not be modified any more than necessary;
- 25 inactivation by mutation or chemical modification are preferred methods of inactivation and a protein protease inhibitor becomes a prime candidate for IPBD. For example, BPTI has been mutated, by the methods of the present invention, to bind to pro-
- 30 teases other than trypsin.

Example III - VI disclose that uninhibited serine proteases may be used as targets quite successfully and that protein protease inhibitors derived from BPTI and selected for binding to these immobilized proteases are

35 excellent inhibitors.

V.F. Immobilization or Labeling of Target Material

For chromatography, FACS, or electrophoresis there may be a need to covalently link the target material to a second chemical entity. For chromatography the second entity is a matrix, for FACS the second entity is a fluorescent dye, and for electrophoresis the second entity is a strongly charged molecule. In many cases, no coupling is required because the target material already has the desired property of: a) immobility, b) fluorescence, or c) charge. In other cases, chemical or physical coupling is required.

Various means may be used to immobilize or label the target materials. The means of immobilization or labeling is, in part, determined by the nature of the target. In particular, the physical and chemical nature of the target and its functional groups of the target material determine which types of immobilization reagents may be most easily used.

For the purpose of selecting an immobilization method, it may be more helpful to classify target materials as follows: (a) solid, whether crystalline or amorphous, and insoluble in an aqueous solvent (e.g., many minerals, and fibrous organics such as cellulose and silk); (b) solid, whether crystalline or amorphous, and soluble in an aqueous solvent; (c) liquid, but insoluble in aqueous phase (e.g., 2,3,3-trimethyldecane); or (d) liquid, and soluble in aqueous media.

It is not necessary that the actual target material be used in preparing the immobilized or labeled analogue that is to be used in affinity separation; rather, suitable reactive analogues of the target material may be more convenient. If 2,3,3-trimethyldecane were the target material, for example, then 2,3,3-trimethyl-10-aminodecane would be far easier to immobilize than the parental

compound. Because the latter compound is modified at one end of the chain, it retains almost all of the shape and charge attributes that differentiate the former compound from other alkanes.

5 Target materials that do not have reactive functional groups may be immobilized by first creating a reactive functional group through the use of some powerful reagent, such as a halogen. For example, an alkane can be immobilized for affinity by first halogenating it and then
10 reacting the halogenated derivative with an immobilized or immobilizable amine.

In some cases, the reactive groups of the actual target material may occupy a part on the target molecule that is to be left undisturbed. In that case, additional
15 functional groups may be introduced by synthetic chemistry. For example, the most reactive groups in cholesterol are on the steroid ring system, viz, -OH and >C=C. We may wish to leave this ring system as it is so that it binds to the novel binding protein. In this case, we prepare an
20 analogue having a reactive group attached to the aliphatic chain (such as 26-aminocholesterol) and immobilize this derivative in a manner appropriate to the reactive group so attached.

Two very general methods of immobilization are widely
25 used. The first is to biotinylate the compound of interest and then bind the biotinylated derivative to immobilized avidin. The second method is to generate antibodies to the target material, immobilize the antibodies by any of numerous methods, and then bind the
30 target material to the immobilized antibodies. Use of antibodies is more appropriate for larger target materials; small targets (those comprising, for example, ten or fewer non-hydrogen atoms) may be so completely engulfed by an antibody that very little of the target is exposed in

the target-antibody complex.

Non-covalent immobilization of hydrophobic molecules without resort to antibodies may also be used. A compound, such as 2,3,3-trimethyldecane is blended with a matrix precursor, such as sodium alginate, and the mixture is extruded into a hardening solution. The resulting beads will have 2,3,3-trimethyldecane dispersed throughout and exposed on the surface.

Other immobilization methods depend on the presence of particular chemical functionalities. A polypeptide will present -NH_2 (N-terminal; Lysines), -COOH (C-terminal; Aspartic Acids; Glutamic Acids), -OH (Serines; Threonines; Tyrosines), and -SH (Cysteines). A polysaccharide has free -OH groups, as does DNA, which has a sugar backbone.

The following table is a nonexhaustive review of reactive functional groups and potential immobilization reagents:

Group	Reagent
R-NH ₂	Derivatives of 2,4,6-trinitrobenzene sulfonates (TNBS), (CREI84, p.11)
R-NH ₂	Carboxylic acid anhydrides, e.g. derivatives of succinic anhydride, maleic anhydride, citraconic anhydride (CREI84, p.11)
R-NH ₂	Aldehydes that form reducible Schiff bases (CREI84, p.12)
guanido	cyclohexanedione derivatives (CREI84, p.14)
R-CO ₂ H	

		Diazo cmpds (CREI84, p.10)
	R-CO ₂ ⁻	
5	R-OH	Epoxides (CREI84, p.10)
		Carboxylic acid anhydrides
10	Aryl-OH	Carboxylic acid anhydrides
	Indole ring	
		Benzyl halide and sulphenyl halides (CREI84, p.19)
15	R-SH	
		N-alkylmaleimides (CREI84, p.21)
20	R-SH	
		ethyleneimine derivatives (CREI84, p.21)
25	R-SH	
		Aryl mercury compounds, (CREI84, P.21)
	R-SH	
		Disulfide reagents, (CREI84, p.23)
30	Thiol ethers	
	Ketones	
35	Aldehydes	Make Schiff's base and reduce with NaBH ₄ . (CREI84, p.12-13)
		Oxidize to COOH, <u>vide supra</u> .
40	R-SO ₃ H	
		Convert to R-SO ₂ Cl and react with immobilized alcohol or amine.
45	R-PO ₃ H	
		Convert to R-PO ₂ Cl and react with immobilized alcohol or amine.
50	CC double bonds	
		Add HBr and then make amine or thiol.

00996630 5096660

The next table identifies the reactive groups of a number of potential targets.

5			Reactive groups or [derivatives]
	<u>Compound (Item#, page)*</u>		
	prostaglandin E2 (2893,1251)		-OH, keto, -COOH, C=C
10	aspartame (861,132)		-NH ₂ , -COOH, -COOCH ₃
	haem (4558, 732)		vinyl, -COOH, Fe
15	bilirubin (1235,189)		vinyl, -COOH, keto, -NH-
20	morphine (6186,988)		-OH, -C=C-, reactive phenyl ring
25	codeine (2459,384)		-OH, -C=C-, reactive phenyl ring
	dichlorodiphenyltrichlorethane (2832,446)		aromatic chlorine, aliphatic chlorine
30	benzo(a)pyrene (1113,172)		[Chlorinate->amine, or make sulfonate-> Aryl-SO ₂ Cl]
35	actinomycin D (2804,441)		aryl-NH ₂ , -OH
40	cellulose		self immobilized
	hydroxylapatite		self immobilized
45	cholesterol (2204,341)		-OH, >C=C-

*Note: Item# and page refer to The Merck Index, 11th

F06290"55096860

Edition.

The extensive literature on affinity chromatography and related techniques will provide further examples.

Matrices suitable for use as support materials include polystyrene, glass, agarose and other chromatographic supports, and may be fabricated into beads, sheets, columns, wells, and other forms as desired. Suppliers of support material for affinity chromatography include: Applied Protein Technologies Cambridge, MA; Bio-
 10 Rad Laboratories, Rockville Center, NY; Pierce Chemical Company, Rockford, IL. Target materials are attached to the matrix in accord with the directions of the manufacturer of each matrix preparation with consideration of good presentation of the target.

15 Early in the selection process, relatively high concentrations of target materials may be applied to the matrix to facilitate binding; target concentrations may subsequently be reduced to select for higher affinity SBDs.

20 V.G. Elution of Lower Affinity PBD-Bearing Genetic Packages

The population of GPs is applied to an affinity matrix under conditions compatible with the intended use of the binding protein and the population is fractionated
 25 by passage of a gradient of some solute over the column. The process enriches for PBDs having affinity for the target and for which the affinity for the target is least affected by the eluants used. The enriched fractions are those containing viable GPs that elute from the column at
 30 greater concentration of the eluant.

The eluants preferably are capable of weakening noncovalent interactions between the displayed PBDs and the immobilized target material. Preferably, the eluants

106290" 5509550

do not kill the genetic package; the genetic message corresponding to successful mini-proteins is most conveniently amplified by reproducing the genetic package rather than by in vitro procedures such as PCR. The list of potential eluants includes salts (including Na⁺, NH₄⁺, Rb⁺, SO₄⁻⁻, H₂PO₄⁻, citrate, K⁺, Li⁺, Cs⁺, HSO₄⁻, CO₃⁻⁻, Ca⁺⁺, Sr⁺⁺, Cl⁻, PO₄⁻⁻⁻, HCO₃⁻, Mg⁺⁺, Ba⁺⁺, Br⁻, HPO₄⁻⁻ and acetate), acid, heat, compounds known to bind the target, and soluble target material (or analogues thereof).

Because bacteria continue to metabolize during affinity separation, the choice of buffer components is more restricted for bacteria than for bacteriophage or spores. Neutral solutes, such as ethanol, acetone, ether, or urea, are frequently used in protein purification and are known to weaken non-covalent interactions between proteins and other molecules. Many of these species are, however, very harmful to bacteria and bacteriophage. Urea is known not to harm M13 up to 8 M. Bacterial spores, on the other hand, are impervious to most neutral solutes. Several affinity separation passes may be made within a single round of variegation. Different solutes may be used in different analyses, salt in one, pH in the next, etc.

Any ions or cofactors needed for stability of PBDs (derived from IPBD) or target are included in initial and elution buffers at appropriate levels. We first remove GP(PBD)s that do not bind the target by washing the matrix with the initial buffer. We determine that this phase of washing is complete by plating aliquots of the washes or by measuring the optical density (at 260 nm or 280 nm). The matrix is then eluted with a gradient of increasing: a) salt, b) [H⁺] (decreasing pH), c) neutral solutes, d) temperature (increasing or decreasing), or e)

some combination of these factors. The solutes in each of the first three gradients have been found generally to weaken non-covalent interactions between proteins and bound molecules. Salt is a preferred solute for gradient formation in most cases. Decreasing pH is also a highly preferred eluant. In some cases, the preferred matrix is not stable to low pH so that salt and urea are the most preferred reagents. Other solutes that generally weaken non-covalent interaction between proteins and the target material of interest may also be used.

The uneluted genetic packages contain DNA encoding binding domains which have a sufficiently high affinity for the target material to resist the elution conditions. The DNA encoding such successful binding domains may be recovered in a variety of ways. Preferably, the bound genetic packages are simply eluted by means of a change in the elution conditions. Alternatively, one may culture the genetic package in situ, or extract the target-containing matrix with phenol (or other suitable solvent) and amplify the DNA by PCR or by recombinant DNA techniques. Additionally, if a site for a specific protease has been engineered into the display vector, the specific protease is used to cleave the binding domain from the GP.

V.H. Optimization of Affinity Chromatography Separation:

For linear gradients, elution volume and eluant concentration are directly related. Changes in eluant concentration cause GPs to elute from the column. Elution volume, however, is more easily measured and specified. It is to be understood that the eluant concentration is the agent causing GP release and that an eluant concentration can be calculated from an elution volume and the specified gradient.

Using a specified elution regime, we compare the elution volumes of GP(IPBD)s with the elution volumes of

TOP SECRET 56056850

wtGP on affinity columns supporting AfM(IPBD). Comparisons are made at various: a) amounts of IPBD/GP, b) densities of AfM(IPBD)/(volume of matrix) (DoAMoM), c) initial ionic strengths, d) elution rates, e) amounts of GP/(volume of support), f) pHs, and g) temperatures, because these are the parameters most likely to affect the sensitivity and efficiency of the separation. We then pick those conditions giving the best separation.

We do not optimize pH or temperature; rather we record optimal values for the other parameters for one or more values of pH and temperature. The pH used must be within the range of pH for which GP(IPBD) binds the AfM(IPBD) that is being used in this step. The conditions of intended use specified by the user may include a specification of pH or temperature. If pH is specified, then pH will not be varied in eluting the column. Decreasing pH may, however, be used to liberate bound GPs from the matrix. Similarly, if the intended use specifies a temperature, we will hold the affinity column at the specified temperature during elution, but we might vary the temperature during recovery. If the intended use specifies the pH or temperature, then we prefer that the affinity separation be optimized for all other parameters at the specified pH and temperature.

In the optimization devised in this step, we preferably use a molecule known to have moderate affinity for the IPBD (K_d in the range 10^{-6} M to 10^{-8} M), for the following reason. When populations of GP(vgPBD)s are fractionated, there will be roughly three subpopulations: a) those with no binding, b) those that have some binding but can be washed off with high salt or low pH, and c) those that bind very tightly and are most easily rescued in situ. We optimize the parameters to separate (a) from (b) rather than (b) from (c). Let PBD_w be a PBD having

T06290" 5505550

weak binding to the target and PBD_S be a PBD having strong binding. Higher DoAMoM might, for example, favor retention of GP(PBD_W) but also make it very difficult to elute viable GP(PBD_S). We will optimize the affinity separation to retain GP(PBD_W) rather than to allow release of GP(PBD_S) because a tightly bound GP(PBD_S) can be rescued by in situ growth. If we find that DoAMoM strongly affects the elution volume, then in part III we may reduce the amount of target on the affinity column when an SBD has been found with moderately strong affinity (K_d on the order of 10^{-7} M) for the target.

In case the promoter of the osp-ipbd gene is not regulated by a chemical inducer, we optimize DoAMoM, the elution rate, and the amount of GP/volume of matrix. If the optimized affinity separation is acceptable, we proceed. If not, we develop a means to alter the amount of IPBD per GP. Among GPs considered in the present invention, this case could arise only for spores because regulatable promoters are available for all other systems.

If the amount of IPBD/spore is too high, we could engineer an operator site into the osp-ipbd gene. We choose the operator sequence such that a repressor sensitive to a small diffusible inducer recognizes the operator. Alternatively, we could alter the Shine-Dalgarno sequence to produce a lower homology with consensus Shine-Dalgarno sequences. If the amount of IPBD/spore is too low, we can introduce variability into the promoter or Shine-Dalgarno sequences and screen colonies for higher amounts of IPBD/spore.

In this step, we measure elution volumes of genetically pure GPs that elute from the affinity matrix as sharp bands that can be detected by UV absorption. Alternatively, samples from effluent fractions can be plated on suitable medium (cells or spores) or on sensi-

T06290-56096860

tive cells (phage) and colonies or plaques counted.

Several values of IPBD/GP, DoAMoM, elution rates, initial ionic strengths, and loadings should be examined. The following is only one of many ways in which the affinity separation could be optimized. We anticipate that optimal values of IPBD/GP and DoAMoM will be correlated and therefore should be optimized together. The effects of initial ionic strength, elution rate, and amount of GP/(matrix volume) are unlikely to be strongly correlated, and so they can be optimized independently.

For each set of parameters to be tested, the column is eluted in a specified manner. For example, we may use a regime called Elution Regime 1: a KCl gradient runs from 10mM to maximum allowed for the GP(IPBD) viability in 100 fractions of 0.05 V_V, followed by 20 fractions of 0.05 V_V at maximum allowed KCl; pH of the buffer is maintained at the specified value with a convenient buffer such as phosphate, Tris, or MOPS. Other elution regimes can be used; what is important is that the conditions of this optimization be similar to the conditions that are used in Part III for selection for binding to target and recovery of GPs from the chromatographic system.

When the osp-ipbd gene is regulated by [XINDUCE], IPBD/GP can be controlled by varying [XINDUCE]. Appropriate values of [XINDUCE] depend on the identity of [XINDUCE] and the promoter; if, for example, XINDUCE is isopropylthiogalactoside (IPTG) and the promoter is lacUV5, then [IPTG] = 0, 0.1 uM, 1.0 uM, 10.0 uM, 100.0 uM, and 1.0 mM would be appropriate levels to test. The range of variation of [XINDUCE] is extended until an optimum is found or an acceptable level of expression is obtained.

DoAMoM is varied from the maximum that the matrix material can bind to 1% or 0.1% of this level in appro-

09895095-06290-5695850

priate steps. We anticipate that the efficiency of separation will be a smooth function of DoAMoM so that it is appropriate to cover a wide range of values for DoAMoM with a coarse grid and then explore the neighborhood of the approximate optimum with a finer grid.

Several values of initial ionic strength are tested, such as 1.0 mM, 5.0 mM, 10.0 mM and 20.0 mM. Low ionic strength favors binding between oppositely charged groups, but could also cause GP to precipitate.

The elution rate is varied, by successive factors of 1/2, from the maximum attainable rate to 1/16 of this value. If the lowest elution rate tested gives the best separation, we test lower elution rates until we find an optimum or adequate separation.

The goal of the optimization is to obtain a sharp transition between bound and unbound GPs, triggered by increasing salt or decreasing pH or a combination of both. This optimization need be performed only: a) for each temperature to be used, b) for each pH to be used, and c) when a new GP(IPBD) is created.

V.I. Measuring the sensitivity of affinity separation:

Once the values of IPBD/GP, DoAMoM, initial ionic strength, elution rate, and amount of GP/(volume of affinity support) have been optimized, we determine the sensitivity of the affinity separation (C_{sensi}) by the following procedure that measures the minimum quantity of GP(IPBD) that can be detected in the presence of a large excess of wtGP. The user chooses a number of separation cycles, denoted N_{chrom} , that will be performed before an enrichment is abandoned; preferably, N_{chrom} is in the range 6 to 10 and N_{chrom} must be greater than 4. Enrichment can be terminated by isolation of a desired GP(SBD) before N_{chrom} passes.

The measurement of sensitivity is significantly expedited if GP(IPBD) and wtGP carry different selectable markers because such markers allow easy identification of colonies obtained by plating fractions obtained from the chromatography column. For example, if wtGP carries kanamycin resistance and GP(IPBD) carries ampicillin resistance, we can plate fractions from a column on non-selective media suitable for the GP. Transfer of colonies onto ampicillin- or kanamycin-containing media will determine the identity of each colony.

Mixtures of GP(IPBD) and wtGP are prepared in the ratios of $1:V_{lim}$, where V_{lim} ranges by an appropriate factor (e.g. 1/10) over an appropriate range, typically 10^{11} through 10^4 . Large values of V_{lim} are tested first; once a positive result is obtained for one value of V_{lim} , no smaller values of V_{lim} need be tested. Each mixture is applied to a column supporting, at the optimal DoAMoM, an AfM(IPBD) having high affinity for IPBD and the column is eluted by the specified elution regime, such as Elution Regime 1. The last fraction that contains viable GPs and an inoculum of the column matrix material are cultured. If GP(IPBD) and wtGP have different selectable markers, then transfer onto selection plates identifies each colony. If GP(IPBD) and wtGP have no selectable markers or the same selectable markers, then a number (e.g. 32) of GP clonal isolates are tested for presence of IPBD. If IPBD is not detected on the surface of any of the isolated GPs, then GPs are pooled from: a) the last few (e.g. 3 to 5) fractions that contain viable GPs, and b) an inoculum taken from the column matrix. The pooled GPs are cultured and passed over the same column and enriched for GP(IPBD) in the manner described. This process is repeated until N_{chrom} passes have been performed, or until the IPBD has been detected on the GPs. If GP(IPBD) is not detected after N_{chrom} passes, V_{lim} is decreased and the process is

repeated.

Once a value for V_{lim} is found that allows recovery of GP(IPBD)s, the factor by which V_{lim} is varied is reduced and additional values are tested until V_{lim} is known to within a factor of two.

C_{sensi} equals the highest value of V_{lim} for which the user can recover GP(IPBD) within N_{chrom} passes. The number of chromatographic cycles (K_{cyc}) that were needed to isolate GP(IPBD) gives a rough estimate of C_{eff} ; C_{eff} is approximately the K_{cyc} th root of V_{lim} :

$$C_{eff} \approx \exp\{ \log_e(V_{lim})/K_{cyc} \}$$

For example, if V_{lim} were 4.0×10^8 and three separation cycles were needed to isolate GP(IPBD), then $C_{eff} \approx 736$.

15 V.J. Measuring the efficiency of separation :

To determine C_{eff} more accurately, we determine the ratio of GP(IPBD)/wtGP loaded onto an AfM(IPBD) column that yields approximately equal amounts of GP(IPBD) and wtGP after elution. We prepare mixtures of GP(IPBD) and wtGP in ratios GP(IPBD):wtGP :: 1:Q; we start Q at twenty times the approximate C_{eff} found above. A 1:Q mixture of GP(IPBD) and wtGP is applied to a AfM(IPBD) column and eluted by the specified elution regime, such as Elution Regime 1. A sample of the last fraction that contains viable GPs is plated at a dilution that gives well separated colonies or plaques. The presence of IPBD or the osp-ipbd gene in each colony or plaque can be determined by a number of standard methods, including: a) use of different selectable markers, b) nitrocellulose filter lift of GPs and detection with AfM(IPBD)* (AUSU87), or c) nitrocellulose filter lift of GPs and detection with radiolabeled DNA that is complementary to the osp-ipbd gene (AUSU87). Let F be the fraction of GP(IPBD) colonies found in the last fraction containing viable GPs. When a

T06290"55096B60

Q is found such that $.20 < F < .80$, then

$$C_{eff} = Q * F.$$

If $F < 0.2$, then we reduce Q by an appropriate factor (e.g. 1/10) and repeat the procedure. If $F > 0.8$,
 5 then we increase Q by an appropriate factor (e.g. 2) and repeat the procedure.

V.K. Reducing selection due to non-specific binding:

When affinity chromatography is used for separating bound and unbound GPs, we may reduce non-specific binding
 10 of GP(PBD)s to the matrix that bears the target in the following ways:

- 1) we treat the column with blocking agents such as genetically defective GPs or a solution of protein before the population of GP(vgPBD)s is chromatographed, and
 15
 - 2) we pass the population of GP(vgPBD)s over a matrix containing no target or a different target from the same class as the actual target prior to affinity chromatography.
- 20 Step (1) above saturates any non-specific binding that the affinity matrix might show toward wild-type GPs or proteins in general; step (2) removes components of our population that exhibit non-specific binding to the matrix or to molecules of the same class as the target. If the
 25 target were horse heart myoglobin, for example, a column supporting bovine serum albumin could be used to trap GPs exhibiting PBDs with strong non-specific binding to proteins. If cholesterol were the target, then a hydrophobic compound, such as p-tertiarybutylbenzyl alcohol,
 30 could be used to remove GPs displaying PBDs having strong non-specific binding to hydrophobic compounds. It is anticipated that PBDs that fail to fold or that are prematurely terminated will be non-specifically sticky.

T.D. 20 36056860

These sequences could outnumber the PBDs having desirable binding properties. Thus, the capacity of the initial column that removes indiscriminately adhesive PBDs should be greater (e.g. 5 fold greater) than the column that
5 supports the target molecule.

Variation in the support material (polystyrene, glass, agarose, cellulose, etc.) in analysis of clones carrying SBDs is used to eliminate enrichment for packages that bind to the support material rather than the target.

10 FACs may be used to separate GPs that bind fluorescent labeled target. We discriminate against artifactual binding to the fluorescent label by using two or more different dyes, chosen to be structurally different. GPs isolated using target labeled with a first dye are
15 cultured. These GPs are then tested with target labeled with a second dye.

Electrophoretic affinity separation uses unaltered target so that only other ions in the buffer can give rise to artifactual binding. Artifactual binding to the gel
20 material gives rise to retardation independent of field direction and so is easily eliminated.

A variegated population of GPs will have a variety of charges. The following 2D electrophoretic procedure accommodates this variation in the population. First the
25 variegated population of GPs is electrophoresed in a gel that contains no target material. The electrophoresis continues until the GP s are distributed along the length of the lane. The gels described by Sewer for phage are very low in agarose and lack mechanical stability. The
30 target-free lane in which the initial electrophoresis is conducted is separate from a square of gel that contains target material by a removable baffle. After the first pass, the baffle is removed and a second electrophoresis is conducted at right angles to the first. GPs that do

T05340-56096360

not bind target migrate with unaltered mobility while GP s that do bind target will separate from the majority that do not bind target. A diagonal line of non-binding GPs will form. This line is excised and discarded. Other parts of the gel are dissolved and the GPs cultured.

V.L. Isolation of GP(PBD)s with binding-to-target phenotypes :

The harvested packages are now enriched for the binding-to-target phenotype by use of affinity separation involving the target material immobilized on an affinity matrix. Packages that fail to bind to the target material are washed away. If the packages are bacteriophage or endospores, it may be desirable to include a bacteriocidal agent, such as azide, in the buffer to prevent bacterial growth. The buffers used in chromatography include: a) any ions or other solutes needed to stabilize the target, and b) any ions or other solutes needed to stabilize the PBDs derived from the IPBD.

V.M. Recovery of packages:

Recovery of packages that display binding to an affinity column may be achieved in several ways, including:

- 1) collect fractions eluted from the column with a gradient as described above; fractions eluting later in the gradient contain GPs more enriched for genes encoding PBDs with high affinity for the column,
- 2) elute the column with the target material in soluble form,
- 3) flood the matrix with a nutritive medium and grow the desired packages in situ,
- 4) remove parts of the matrix and use them to inoculate growth medium,
- 5) chemically or enzymatically degrade the linkage

holding the target to the matrix so that GPs still bound to target are eluted, or

- 6) degrade the packages and recover DNA with phenol or other suitable solvent; the recovered DNA is used to transform cells that regenerate GPs.

It is possible to utilize combinations of these methods. It should be remembered that what we want to recover from the affinity matrix is not the GPs per se, but the information in them. Recovery of viable GPs is very strongly preferred, but recovery of genetic material is essential. If cells, spores, or virions bind irreversibly to the matrix but are not killed, we can recover the information through in situ cell division, germination, or infection respectively. Proteolytic degradation of the packages and recovery of DNA is not preferred.

Although degradation of the bound GPs and recovery of genetic material is a possible mode of operation, inadvertent inactivation of the GPs is very deleterious. It is preferred that maximum limits for solutes that do not inactivate the GPs or denature the target or the column are determined. If the affinity matrices are expendable, one may use conditions that denature the column to elute GPs; before the target is denatured, a portion of the affinity matrix should be removed for possible use as an inoculum. As the GPs are held together by protein-protein interactions and other non-covalent molecular interactions, there will be cases in which the molecular package will bind so tightly to the target molecules on the affinity matrix that the GPs can not be washed off in viable form. This will only occur when very tight binding has been obtained. In these cases, methods (3) through (5) above can be used to obtain the bound packages or the genetic messages from the affinity matrix.

It is possible, by manipulation of the elution

conditions, to isolate SBDs that bind to the target at one pH (pH_p) but not at another pH (pH_o). The population is applied at pH_p and the column is washed thoroughly at pH_p . The column is then eluted with buffer at pH_o and GPs that
 5 come off at the new pH are collected and cultured. Similar procedures may be used for other solution parameters, such as temperature. For example, GP(vgPBD)s could be applied to a column supporting insulin. After eluting with salt to remove GPs with little or no binding
 10 to insulin, we elute with salt and glucose to liberate GPs that display PBDs that bind insulin or glucose in a competitive manner.

V.N. Amplifying the Enriched Packages

Viable GPs having the selected binding trait are
 15 amplified by culture in a suitable medium, or, in the case of phage, infection into a host so cultivated. If the GPs have been inactivated by the chromatography, the OCV carrying the osp-pbd gene are recovered from the GP, and introduced into a new, viable host.

20 V.O. Determining whether further enrichment is needed:

The probability of isolating a GP with improved binding increases by C_{eff} with each separation cycle. Let N be the number of distinct amino-acid sequences produced by the variegation. We want to perform K separation
 25 cycles before attempting to isolate an SBD, where K is such that the probability of isolating a single SBD is 0.10 or higher.

$$K = \text{the smallest integer} \geq \log_{10}(0.10 N) / \log_{10}(C_{eff})$$

For example, if N were $1.0 \cdot 10^7$ and $C_{eff} = 6.31 \cdot 10^2$,
 30 then $\log_{10}(1.0 \cdot 10^6) / \log_{10}(6.31 \cdot 10^2) = 6.0000 / 2.8000 = 2.14$. Therefore we would attempt to isolate SBDs after the third separation cycle. After only two separation cycles, the probability of finding an SBD is

$$(6.31 \times 10^2)^2 / (1.0 \times 10^7) = .04$$

and attempting to isolate SBDs might be profitable.

- Clonal isolates from the last fraction eluted which contained any viable GPs, as well as clonal isolates
 5 obtained by culturing an inoculum taken from the affinity matrix, are cultured in a growth step that is similar to that described previously. Other fractions may be cultured too. If K separation cycles have been completed, samples from a number, e.g. 32, of these clonal isolates
 10 are tested for elution properties on the {target} column. If none of the isolated, genetically pure GPs show improved binding to target, or if K cycles have not yet been completed, then we pool and culture, in a manner similar to the manner set forth previously, the GPs from
 15 the last few fractions eluted that contained viable GPs and from the GPs obtained by culturing an inoculum taken from the column matrix. We then repeat the enrichment procedure described above. This cyclic enrichment may continue N_{chrom} passes or until an SBD is isolated.
- 20 If one or more of the isolated GPs has improved retention on the {target} column, we determine whether the retention of the candidate SBDs is due to affinity for the target material as follows. A second column is prepared using a different support matrix with the target material
 25 bound at the optimal density. The elution volumes, under the same elution conditions as used previously, of candidate GP(SBD)s are compared to each other and to GP(PPBD of this round). If one or more candidate GP(SBD)s has a larger elution volume than GP(PPBD of this round),
 30 then we pick the GP(SBD) having the highest elution volume and proceed to characterize the population. If none of the candidate GP(SBD)s has higher elution volume than GP(PPBD of this round), then we pool and culture, in a manner similar to the manner used previously, the GPs from

059509 062901

the last few fractions that contained viable GPs and the GPs obtained by culturing an inoculum taken from the column matrix. We then repeat the enrichment procedure.

If all of the SBDs show binding that is superior to
 5 PPBD of this round, we pool and culture the GPs from the last fraction that contains viable GPs and from the inoculum taken from the column. This population is re-chromatographed at least one pass to fractionate further the GPs based on K_d .

10 If an RNA phage were used as GP, the RNA would either be cultured with the assistance of a helper phage or be reverse transcribed and the DNA amplified. The amplified DNA could then be sequenced or subcloned into suitable plasmids.

15 V.P. Characterizing the Putative SBDs:

We characterize members of the population showing desired binding properties by genetic and biochemical methods. We obtain clonal isolates and test these strains by genetic and affinity methods to determine genotype and
 20 phenotype with respect to binding to target. For several genetically pure isolates that show binding, we demonstrate that the binding is caused by the artificial chimeric gene by excising the osp-sbd gene and crossing it into the parental GP. We also ligate the deleted backbone of each
 25 GP from which the osp-sbd is removed and demonstrate that each backbone alone cannot confer binding to the target on the GP. We sequence the osp-sbd gene from several clonal isolates. Primers for sequencing are chosen from the DNA flanking the osp-ppbd gene or from parts of the osp-ppbd
 30 gene that are not variegated.

The present invention is not limited to a single method of determining protein sequences, and reference in the appended claims to determining the amino acid sequence

09895095 DEPT

of a domain is intended to include any practical method or combination of methods, whether direct or indirect. The preferred method, in most cases, is to determine the sequence of the DNA that encodes the protein and then to
 5 infer the amino acid sequence. In some cases, standard methods of protein-sequence determination may be needed to detect post-translational processing.

The present invention is not limited to a single method of determining the sequence of nucleotides (nts) in
 10 DNA subsequences. In the preferred embodiment, plasmids are isolated and denatured in the presence of a sequencing primer, about 20 nts long, that anneals to a region adjacent, on the 5' side, to the region of interest. This plasmid is then used as the template in the four sequenc-
 15 ing reactions with one dideoxy substrate in each. Sequencing reactions, agarose gel electrophoresis, and polyacrylamide gel electrophoresis (PAGE) are performed by standard procedures (AUSU87).

For one or more clonal isolates, we may subclone the
 20 sbd gene fragment, without the osp fragment, into an expression vector such that each SBD can be produced as a free protein. Because numerous unique restriction sites were built into the inserted domain, it is easy to subclone the gene at any time. Each SBD protein is
 25 purified by normal means, including affinity chromatography. Physical measurements of the strength of binding are then made on each free SBD protein by one of the following methods: 1) alteration of the Stokes radius as a function of binding of the target material, measured by
 30 characteristics of elution from a molecular sizing column such as agarose, 2) retention of radiolabeled binding protein on a spun affinity column to which has been affixed the target material, or 3) retention of radio-labeled target material on a spun affinity column to which

0095095-063901
 0095095-063901

has been affixed the binding protein. The measurements of binding for each free SBD are compared to the corresponding measurements of binding for the PPBD.

In each assay, we measure the extent of binding as a function of concentration of each protein, and other relevant physical and chemical parameters such as salt concentration, temperature, pH, and prosthetic group concentrations (if any).

In addition, the SBD with highest affinity for the target from each round is compared to the best SBD of the previous round (IPBD for the first round) and to the IPBD (second and later rounds) with respect to affinity for the target material. Successive rounds of mutagenesis and selection-through-binding yield increasing affinity until desired levels are achieved.

If we find that the binding is not yet sufficient, we decide which residues to vary next. If the binding is sufficient, then we now have a expression vector bearing a gene encoding the desired novel binding protein.

20 V.Q. Joint selections:

One may modify the affinity separation of the method described to select a molecule that binds to material A but not to material B. One needs to prepare two selection columns, one with material A and the other with material B. The population of genetic packages is prepared in the manner described, but before applying the population to A, one passes the population over the B column so as to remove those members of the population that have high affinity for B ("reverse affinity chromatography"). In the preceding specification, the initial column supported some other molecule simply to remove GP(PBD)s that displayed PBDs having indiscriminate affinity for surfaces.

It may be necessary to amplify the population that does not bind to B before passing it over A. Amplification would most likely be needed if A and B were in some ways similar and the PPBD has been selected for having affinity for A. The optimum order of interactions might be determined empirically. For example, to obtain an SBD that binds A but not B, three columns could be connected in series: a) a column supporting some compound, neither A nor B, or only the matrix material, b) a column supporting B, and c) a column supporting A. A population of GP(vg-PBD)s is applied to the series of columns and the columns are washed with the buffer of constant ionic strength that is used in the application. The columns are uncoupled, and the third column is eluted with a gradient to isolate GP(PBD)s that bind A but not B.

One can also generate molecules that bind to both A and B. In this case we can use a 3D model and mutate one face of the molecule in question to get binding to A. One can then mutate a different face to produce binding to B. When an SBD binds at least somewhat to both A and B, one can mutate the chain by Diffuse Mutagenesis to refine the binding and use a sequential joint selection for binding to both A and B.

The materials A and B could be proteins that differ at only one or a few residues. For example, A could be a natural protein for which the gene has been cloned and B could be a mutant of A that retains the overall 3D structure of A. SBDs selected to bind A but not B probably bind to A near the residues that are mutated in B. If the mutations were picked to be in the active site of A (assuming A has an active site), then an SBD that binds A but not B will bind to the active site of A and is likely to be an inhibitor of A.

To obtain a protein that will bind to both A and B,

we can, alternatively, first obtain an SBD that binds A and a different SBD that binds B. We can then combine the genes encoding these domains so that a two-domain single-polypeptide protein is produced. The fusion protein will
 5 have affinity for both A and B because one of its domains binds A and the other binds B.

One can also generate binding proteins with affinity for both A and B, such that these materials will compete for the same site on the binding protein. We guarantee
 10 competition by overlapping the sites for A and B. Using the procedures of the present invention, we first create a molecule that binds to target material A. We then vary a set of residues defined as: a) those residues that were varied to obtain binding to A, plus b) those residues
 15 close in 3D space to the residues of set (a) but that are internal and so are unlikely to bind directly to either A or B. Residues in set (b) are likely to make small changes in the positioning of the residues in set (a) such that the affinities for A and B will be changed by small
 20 amounts. Members of these populations are selected for affinity to both A and B.

V.R. Selection for non-binding:

The method of the present invention can be used to select proteins that do not bind to selected targets.
 25 Consider a protein of pharmacological importance, such as streptokinase, that is antigenic to an undesirable extent. We can take the pharmacologically important protein as IPBD and antibodies against it as target. Residues on the surface of the pharmacologically important protein would
 30 be variegated and GP(PBD)s that do not bind to an antibody column would be collected and cultured. Surface residues may be identified in several ways, including: a) from a 3D structure, b) from hydrophobicity considerations, or c) chemical labeling. The 3D structure of the pharmacologi-

FOI 5095850

cally important protein remains the preferred guide to picking residues to vary, except now we pick residues that are widely spaced so that we leave as little as possible of the original surface unaltered.

5 Destroying binding frequently requires only that a single amino acid in the binding interface be changed. If polyclonal antibodies are used, we face the problem that all or most of the strong epitopes must be altered in a single molecule. Preferably, one would have a set of
10 monoclonal antibodies, or a narrow range of antibody species. If we had a series of monoclonal antibody columns, we could obtain one or more mutations that abolish binding to each monoclonal antibody. We could then combine some or all of these mutations in one
15 molecule to produce a pharmacologically important protein recognized by none of the monoclonal antibodies. Such mutants are tested to verify that the pharmacologically interesting properties have not be altered to an unacceptable degree by the mutations.

20 Typically, polyclonal antibodies display a range of binding constants for antigen. Even if we have only polyclonal antibodies that bind to the pharmacologically important protein, we may proceed as follows. We engineer the pharmacologically important protein to appear on the
25 surface of a replicable GP. We introduce mutations into residues that are on the surface of the pharmacologically important protein or into residues thought to be on the surface of the pharmacologically important protein so that a population of GPs is obtained. Polyclonal antibodies
30 are attached to a column and the population of GPs is applied to the column at low salt. The column is eluted with a salt gradient. The GPs that elute at the lowest concentration of salt are those which bear pharmacologically important proteins that have been mutated in a

way that eliminates binding to the antibodies having maximum affinity for the pharmacologically important protein. The GPs eluting at the lowest salt are isolated and cultured. The isolated SBD becomes the PPBD to
 5 further rounds of variegation so that the antigenic determinants are successively eliminated.

V.S. Selection of PBDs for retention of structure:

Let us take an SBD with known affinity for a target as PPBD to a variegation of a region of the PBD that is
 10 far from the residues that were varied to create the SBD. We can use the target as an affinity molecule to select the PBDs that retain binding for the target, and that presumably retain the underlying structure of the IPBD. The variegations in this case could include insertions and
 15 deletions that are likely to disrupt the IPBD structure. We could also use the IPBD and AfM(IPBD) in the same way.

For example, if IPBD were BPTI and AfM(BPTI) were trypsin, we could introduce four or five additional residue after residue 26 and select GPs that display PBDs
 20 having specific affinity for AfM(BPTI). Residue 26 is chosen because it is in a turn and because it is about 25 A from K15, a key amino acid in binding to trypsin.

The underlying structure is most likely to be retained if insertions or deletions are made at loops or
 25 turns.

V.T. Engineering of Antagonists

It may be desirable to provide an antagonist to an enzyme or receptor. This may be achieved by making a molecule that prevents the natural substrate or agonist
 30 from reaching the active site. Molecules that bind directly to the active site may be either agonists or antagonists. Thus we adopt the following strategy. We consider enzymes and receptors together under the designa-

T06290"56096860

tion TER (Target Enzyme or Receptor).

For most TERs, there exist chemical inhibitors that block the active site. Usually, these chemicals are useful only as research tools due to highly toxicity. We make two affinity matrices: one with active TER and one with blocked TER. We make a variegated population of GP(PBD)s and select for SBPs that bind to both forms of the enzyme, thereby obtaining SDPs that do not bind to the active site. We expect that SBDs will be found that bind different places on the enzyme surface. Pairs of the sbd genes are fused with an intervening peptide segment. For example, if SBD-1 and SBD-2 are binding domains that show high affinity for the target enzyme and for which the binding is non-competitive, then the gene sbd-1::linker::sbd-2 encodes a two-domain protein that will show high affinity for the target. We make several fusions having a variety of SBDs and various linkers. Such compounds have a reasonable probability of being an antagonist to the target enzyme.

VI. EXPLOITATION OF SUCCESSFUL BINDING DOMAINS AND CORRESPONDING DNAS

VI.A. Generally

Using the method of the present invention, we can obtain a replicable genetic package that displays a novel protein domain having high affinity and specificity for a target material of interest. Such a package carries both amino-acid embodiments of the binding protein domain and a DNA embodiment of the gene encoding the novel binding domain. The presence of the DNA facilitates expression of a protein comprising the novel binding protein domain within a high-level expression system, which need not be the same system used during the developmental process.

VI.B. Production of Novel Binding Proteins

TOP SECRET S6056360

	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100	2101	2102	2103	2104	2105	2106	2107	2108	2109	2110	2111	2112	2113	2114	2115	2116	2117	2118	2119	2120	2121	2122	2123	2124	2125	2126	2127	2128	2129	2130	2131	2132	2133	2134	2135	2136	2137	2138	2139	2140	2141	2142	2143	2144	2145	2146	2147	2148	2149	2150	2151	2152	2153	2154	2155	2156	2157	2158	2159	2160	2161	2162	2163	2164	2165	2166	2167	2168	2169	2170	2171	2172	2173	2174	2175	2176	2177	2178	2179	2180	2181	2182	2183	2184	2185	2186	2187	2188	2189	2190	2191	2192	2193	2194	2195	2196	2197	2198	2199	2200	2201	2202	2203	2204	2205	2206	2207	2208	2209	2210	2211	2212	2213	2214	2215	2216	2217	2218	2219	2220	2221	2222	2223	2224	2225	2226	2227	2228	2229	2230	2231	2232	2233	2234	2235	2236	2237	2238	2239	2240	2241	2242	2243	2244	2245	2246	2247	2248	2249	2250	2251	2252	2253	2254	2255	2256	2257	2258	2259	2260	2261	2262	2263	2264	2265	2266	2267	2268	2269	2270	2271	2272	2273	2274	2275	2276	2277	2278	2279	2280	2281	2282	2283	2284	2285	2286	2287	2288	2289	2290	2291	2292	2293	2294	2295	2296	2297	2298	2299	2300	2301	2302	2303	2304	2305	2306	2307	2308	2309	2310	2311	2312	2313	2314	2315	2316	2317	2318	2319	2320	2321	2322	2323	2324	2325	2326	2327	2328	2329	2330	2331	2332	2333	2334	2335	2336	2337	2338	2339	2340	2341	2342	2343	2344	2345	2346	2347	2348	2349	2350	2351	2352	2353	2354	2355	2356	2357	2358	2359	2360	2361	2362	2363	2364	2365	2366	2367	2368	2369	2370	2371	2372	2373	2374	2375	2376	2377	2378	2379	2380	2381	2382	2383	2384	2385	2386	2387	2388	2389	2390	2391	2392	2393	2394	2395	2396	2397	2398	2399	2400	2401	2402	2403	2404	2405	2406	2407	2408	2409	2410	2411	2412	2413	2414	2415	2416	2417	2418	2419	2420	2421	2422	2
--	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	---

5

Option (c) may be illustrated as follows. Assume that a novel BPTI derivative has been obtained by selection of M13 derivatives in which a population of BPTI-derived domains are displayed as fusions to mature coat protein. Assume that a specific protease cleavage site (e.g. that of activated clotting factor X) is engineered into the amino-acid sequence between the carboxy terminus of the BPTI-derived domain and the mature coat domain. Furthermore, we alter the display system to maximize the number of fusion proteins displayed on each phage. The desired phage can be produced and purified, for example by centrifugation, so that no bacterial products remain. Treatment of the purified phage with a catalytic amount of factor X cleaves the binding domains from the phage particles. A second centrifugation step separates the cleaved protein from the phage, leaving a very pure protein preparation.

As previously mentioned, an advantage inhering from the use of a mini-protein as an IPBD is that it is likely that the derived SBD will also behave like a mini-protein and will be obtainable by means of chemical synthesis. (The term "chemical synthesis", as used herein, includes the use of enzymatic agents in a cell-free environment.)

It is also to be understood that mini-proteins obtained by the method of the present invention may be

taken as lead compounds for a series of homologues that contain non-naturally occurring amino acids and groups other than amino acids. For example, one could synthesize a series of homologues in which each member of the series
 5 has one amino acid replaced by its D enantiomer. One could also make homologues containing constituents such as β alanine, aminobutyric acid, 3-hydroxyproline, 2-Aminoadipic acid, N-ethylasparagine, norvaline, etc.; these would be tested for binding and other properties of interest,
 10 such as stability and toxicity.

Peptides may be chemically synthesized either in solution or on supports. Various combinations of stepwise synthesis and fragment condensation may be employed.

During synthesis, the amino acid side chains are
 15 protected to prevent branching. Several different protective groups are useful for the protection of the thiol groups of cysteines:

- 1) 4-methoxybenzyl (MBzl; Mob)(NISH82; ZAF88), removable with HF;
- 20 2) acetamidomethyl (Acm)(NISH82; NISH86; BECK89c), removable with iodine; mercury ions (e.g., mercuric acetate); silver nitrate; and
- 3) S-para-methoxybenzyl (HOUG84).

Other thiol protective groups may be found in
 25 standard reference works such as Greene, PROTECTIVE GROUPS IN ORGANIC SYNTHESIS (1981).

Once the polypeptide chain has been synthesized, disulfide bonds must be formed. Possible oxidizing agents include air (HOUG84; NISH86), ferricyanide (NISH82;
 30 HOUG84), iodine (NISH82), and performic acid (HOUG84). Temperature, pH, solvent, and chaotropic chemicals may affect the course of the oxidation.

T06230 55096850

biologically active form: conotoxin G1 (13AA, 4 Cys) (NISH-82); heat-stable enterotoxin ST (18AA, 6 Cys) (HOUG84); analogues of ST (BHAT86); Ω -conotoxin GVIA (27AA, 6Cys) (NISH86; RIVI87b); Ω -conotoxin MVIIA (27 AA, 6 Cys) (OLIV87b); α -conotoxin SI (13 AA, 4 Cys) (ZAF88); μ -conotoxin IIIa (22AA, 6 Cys) (BECK89c, CRUZ89, HATA90). Sometimes, the polypeptide naturally folds so that the correct disulfide bonds are formed. Other times, it must be helped along by use of a differently removable protective group for each pair of cysteines.

VI.D. Uses of Novel Binding Proteins

The successful binding domains of the present invention may, alone or as part of a larger protein, be used for any purpose for which binding proteins are suited, including isolation or detection of target materials. In furtherance of this purpose, the novel binding proteins may be coupled directly or indirectly, covalently or noncovalently, to a label, carrier or support.

When used as a pharmaceutical, the novel binding proteins may be contained with suitable carriers or adjuvanants.

* * * * *

All references cited anywhere in this specification are incorporated by reference to the extent which they may be pertinent.

EXAMPLE I

DISPLAY OF BPTI AS A FUSION TO M13 GENE VIII PROTEIN:

Example I involves display of BPTI on M13 as a fusion to the mature gene VIII coat protein. Each of the DNA constructions was confirmed by restriction digestion analysis and DNA sequencing.

1. Construction of the viii-signal-sequence::bpti::mature-viii-coat-protein Display Vector.

A. Operative cloning vectors (OCV).

10 The operative cloning vectors are M13 and phagemids derived from M13 or f1. The initial construction was in the f1-based phagemid pGEM-3Zf(-)^(TM) (Promega Corp., Madison, WI.).

15 A gene comprising, in order,: i) a modified lacUV5 promoter, ii) a Shine-Dalgarno sequence, iii) DNA encoding the M13 gene VIII signal sequence, iv) a sequence encoding mature BPTI, v) a sequence encoding the mature-M13-gene-VIII coat protein, vi) multiple stop codons, and vii) a transcription terminator, was constructed. This gene is
20 illustrated in Tables 101-105; each table shows the same DNA sequence with different features annotated. There are a number of differences between this gene and the one proposed in the hypothetical example in the generic specification of the parent application. Because the
25 actual construction was made in pGEM-3Zf(-), the ends of the synthetic DNA were made compatible with SalI and BamHI. The lacO operator of lacUV5 was changed to the symmetrical lacO with the intention of achieving tighter repression in the absence of IPTG. Several silent codon
30 changes were made so that the longest segment that is identical to wild-type gene VIII is minimized so that genetic recombination with the co-existing gene VIII is unlikely.

T06299 55095850

i) OCV based upon pGEM-3Zf.

pGEM-3Zf^(TM) (Promega Corp., Madison, WI.) is a plasmid-based vector containing the amp gene, bacterial origin of replication, bacteriophage f1 origin of replication, a lacZ operon containing a multiple cloning site sequence, and the T7 and SP6 polymerase binding sequences.

Two restriction enzyme recognition sites were introduced, by site-directed oligonucleotide mutagenesis, at the boundaries of the lacZ operon. This allowed for the removal of the lacZ operon and its replacement with the synthetic gene. A BamHI recognition site (GGATCC) was introduced at the 5' end of the lacZ operon by the mutation of bases C₃₃₁ and T₃₃₂ to G and A respectively (numbering of Promega). A SalI recognition site (GTCGAC) was introduced at the 3' end of the operon by the mutation of bases C₃₀₂₁ and T₃₀₂₃ to G and C respectively. A construct combining these variants of pGEM-3Zf was designated pGEM-MB3/4.

ii) OCV based upon M13mp18.

M13mp18 (YANI85) is an M13 bacteriophage-based vector (available from, inter alia, New England Biolabs, Beverly, MA.) consisting of the whole of the phage genome into which has been inserted a lacZ operon containing a multiple cloning site sequence (MESS77). Two restriction enzyme sites were introduced into M13mp18 using standard methods. A BamHI recognition site (GGATCC) was introduced at the 5' end of the lacZ operon by the mutation of bases C₆₀₀₃ and G₆₀₀₄ to A and T respectively (numbering of Messing). This mutation also destroyed a unique NarI site. A SalI recognition site (GTCGAC) was introduced at the 3' end of the operon by the mutation of bases A₆₄₃₀ and C₆₄₃₂ to C and A respectively. A construct combining these variants of M13mp18 was designated M13-MB1/2.

T06290" 55096860

B) Synthetic Gene.

A synthetic gene (VIII-signal-sequence::mature-bpti::mature-VIII-coat-protein) was constructed from 16 synthetic oligonucleotides (Table 105), custom synthesized by Genetic Designs Inc. of Houston, Texas, using methods detailed in KIMH89 and ASHM89. Table 101 shows the DNA sequence; Table 102 contains an annotated version of this sequence. Table 103 shows the overlaps of the synthetic oligonucleotides in relationship to the restriction sites and coding sequence. Table 104 shows the synthetic DNA in double-stranded form. Table 105 shows each of the 16 synthetic oligonucleotides from 5'-to-3'. The oligonucleotides were phosphorylated, with the exception of the 5' most molecules, using standard methods, annealed and ligated in stages such that a final synthetic duplex was generated. The overhanging ends of this duplex was filled in with T4 DNA polymerase and it was cloned into the HincII site of pGEM-3Zf(-); the initial construct is called pGEM-MB1 (Table 101a). Double-stranded DNA of pGEM-MB1 was cut with PstI, filled in with T4 DNA polymerase and ligated to a SalI linker (New England BioLabs) so that the synthetic gene is bounded by BamHI and SalI sites (Table 101b and Table 102b). The synthetic gene was obtained on a BamHI-SalI cassette and cloned into pGEM-MB3/4 and M13-MB1/2 utilizing the BamHI and SalI sites previously introduced, to generate the constructs designated pGEM-MB16 and M13-MB15, respectively. The full length of the synthetic insert was sequenced and found to be unambiguously correct except for: 1) a missing G in the Shine-Dalgarno sequence; and 2) a few silent errors in the third bases of some codons (shown as upper case in Table 101). Table 102 shows the Ribosome-binding site A₁₀₄GGAGG but the actual sequence is A₁₀₄GAGG. Efforts to express protein from this construction, in vivo and in vitro, were unavailing.

T06290-56096360

C) Alterations to the synthetic gene.

i) Ribosome binding site (RBS).

Starting with the construct pGEM-MB16, a fragment of DNA bounded by the restriction enzyme sites SacI and NheI (containing the original RBS) was replaced with a synthetic oligonucleotide duplex (with compatible SacI and NheI overhangs) containing the sequence for a new RBS that is very similar to the RBS of E. coli phoA and that has been shown to be functional.

10 Original putative RBS (5'-to-3')

GAGCTCagaggCTTACTATGAAGAAATCTCTGGTTCTTAAGGCTAGC
| SacI | | Nhe I |

New RBS (5'-to-3')

15 GAGCTCTggaggaAATAAAATGAAGAAATCTCTGGTTCTTAAGGCTAGC
| SacI | | Nhe I |

The putative RBSs above are lower case and the initiating methionine codon is underscored and bold. The resulting construct was designated pGEM-MB20. In vitro expression of the gene carried by pGEM-MB20 produced a novel protein species of the expected size, about 14.5 kd.

ii) tac promoter.

25 In order to obtain higher expression levels of the fusion protein, the lacUV5 promoter was changed to a tac promoter. Starting with the construct pGEM-MB16, which contains the lacUV5 promoter, a fragment of DNA bounded by the restriction enzyme sites BamHI and HpaII was excised and replaced with a compatible synthetic oligonucleotide duplex containing the -35 sequence of the trp promoter, Cf RUSS82. This converted the lacUV5 promoter to a tac promoter in a construct designated pGEM-MB22, Table 112.

MB16

5'- GATCC tctagagtcggc TTTACA ctttatgcttc(cg-gctcg...-3'
 3'- G agatctcagccg aaatgt gaaatacgaag gc(cgagc...-5'
 5 | | -35 | |
 | | |
 BamHI HpaII

MB22 insert

10 5'- GATCC actcccatcccccctg TTGACA attaatacat -3'
 3'- G tgaggggtagggggac AACTGT taattagtagc-5'
 | | -35 | |
 | | |
 BamHI (HpaII)

15

Promoter and RBS variants of the fusion protein gene were constructed by basic DNA manipulation techniques to generate the following:

20	Promoter	RBS	Encoded Protein.
pGEM-MB16	lac	old	VIIIIs.p.-BPTI-matureVIII
pGEM-MB20	lac	new	" "
pGEM-MB22	tac	old	" "
pGEM-MB26	tac	new	" "

25 The synthetic gene from variants pGEM-MB20 and pGEM-MB26 were recloned into the altered phage vector M13-MB1/2 to generate the phage constructs designated M13-MB27 and M13-MB28 respectively.

iii. Signal Peptide Sequence.

30 In vitro expression of the synthetic gene regulated by tac and the "new" RBS produced a novel protein of the expected size for the unprocessed protein (about 16 kd). In vivo expression also produced novel protein of full size; no processed protein could be seen on phage or in
 35 cell extracts by silver staining or by Western analysis with anti-BPTI antibody.

Thus we analyzed the signal sequence of the fusion. Table 106 shows a number of typical signal sequences. Charged residues are generally thought to be of great

FO6230" 55055350

importance and are shown bold and underscored. Each signal sequence contains a long stretch of uncharged residues that are mostly hydrophobic; these are shown in lower case. At the right, in parentheses, is the length of the stretch of uncharged residues. We note that the fusions of gene VIII signal to BPTI and gene III signal to BPTI have rather short uncharged segments. These short uncharged segments may reduce or prevent processing of the fusion peptides. We know that the gene III signal sequence is capable of directing: a) insertion of the peptide comprising (mature-BPTI)::(mature-gene-III-protein) into the lipid bilayer, and b) translocation of BPTI and most of the mature gene III protein across the lipid bilayer (vide infra). That the gene III remains anchored in the lipid bilayer until the phage is assembled is directed by the uncharged anchor region near the carboxy terminus of the mature gene III protein (see Table 116) and not by the secretion signal sequence. The phoA signal sequence can direct secretion of mature BPTI into the periplasm of E. coli (MARK86). Furthermore, there is controversy over the mechanism by which mature authentic gene VIII protein comes to be in the lipid bilayer prior to phage assembly:

Thus we decided to replace the DNA coding on expression for the gene-VIII-putative-signal-sequence by each of: 1) DNA coding on expression for the phoA signal sequence, 2) DNA coding on expression for the bla signal sequence, or 3) DNA coding on expression for the M13 gene III signal. Each of these replacements produces a tripartite gene encoding a fusion protein that comprises, in order: (a) a signal peptide that directs secretion into the periplasm of parts (b) and (c), derived from a first gene; (b) an initial potential binding domain (BPTI in this case), derived from a second gene (in this case, the second gene is an animal gene); and (c) a structural

00896095-062901
106290-5505660

packaging signal (the mature gene VIII coat protein), derived from a third gene.

The process by which the IPBD::packaging-signal fusion arrives on the phage surface is illustrated in Figure 1. In Figure 1a, we see that authentic gene VIII protein appears (by whatever process) in the lipid bilayer so that both the amino and carboxy termini are in the cytoplasm. Signal peptidase-I cleaves the gene VIII protein liberating the signal peptide (that is absorbed by the cell) and mature gene VIII coat protein that spans the lipid bilayer. Many copies of mature gene VIII coat protein accumulate in the lipid bilayer awaiting phage assembly (Figure 1c). Some signal sequences are able to direct the translocation of quite large proteins across the lipid bilayer. If additional codons are inserted after the codons that encode the cleavage site of the signal peptidase-I of such a potent signal sequence, the encoded amino acids will be translocated across the lipid bilayer as shown in Figure 1b. After cleavage by signal peptidase-I, the amino acids encoded by the added codons will be in the periplasm but anchored to the lipid bilayer by the mature gene VIII coat protein, Figure 1d. The circular single-stranded phage DNA is extruded through a part of the lipid bilayer containing a high concentration of mature gene VIII coat protein; the carboxy terminus of each coat protein molecule packs near the DNA while the amino terminus packs on the outside. Because the fusion protein is identical to mature gene VIII coat protein within the trans-bilayer domain, the fusion protein will co-assemble with authentic mature gene VIII coat protein as shown in Figure 1e.

In each case, the mature VIII coat protein moiety is intended to co-assemble with authentic mature VIII coat protein to produce phage particle having BPTI domains

0996095.062901

displayed on the surface. The source and character of the secretion signal sequence is not important because the signal sequence is cut away and degraded. The structural packaging signal, however, is quite important because it must co-assemble with the authentic coat protein to make a working virus sheath.

a) Bacterial Alkaline Phosphatase (phoA) Signal Peptide.

Construct pGEM-MB26 contains a fragment of DNA bounded by restriction enzyme sites SacI and AccIII which contains the new RBS and sequences encoding the initiating methionine and the signal peptide of M13 gene VIII pro-protein. This fragment was replaced with a synthetic duplex (constructed from four annealed oligonucleotides) containing the RBS and DNA coding for the initiating methionine and signal peptide of PhoA (INOUB2). The resulting construct was designated pGEM-MB42; the sequence of the fusion gene is shown in Table 113. M13MB48 is a derivative of GemMB42. A BamHI-SalI DNA fragment from GenMB42, containing the gene construct, was ligated into a similarly cleaved vector M13MB1/2 giving rise to M13MB48.

PhoA RBS and signal peptide sequence

5'-GAGCTCCATGGGAGAAAATAAA.ATG.AAA.CAA.AGC.ACG.-
|SacI| met lys gln ser thr
 25 .ATC.GCA.CTC.TTA.CCG.TTA.CTG.TTT.ACC.CCT.GTG.ACA.-
 ile ala leu leu pro leu leu phe thr pro val thr
 .AAA.GCC.CGT.CCG.GAT.-3'
 lys ala arg pro asp.....
 30 |AccIII|

b) beta-lactamase signal peptide.

To enable the introduction of the beta-lactamase (amp) promoter and DNA coding for the signal peptide into the gene encoding (mature-BPTI)::(mature-VIII-coat-protein) an initial manipulation of the amp gene (encoding beta-lactamase) was required. Starting with pGEM-3Zf an

AccIII recognition site (TCCGGA) was introduced into the amp gene adjacent to the DNA sequence encoding the amino acids at the beta-lactamase signal peptide cleavage site. Using standard methods of in vitro site-directed oligo-
 5 nucleotide mutagenesis bases C₂₅₀₄ and A₂₅₀₁ were converted to T and G respectively to generate the construct designated pGEM-MB40. Further manipulation of pGEM-MB40 entailed the insertion of a synthetic oligonucleotide linker (CGGATCCG) containing the BamHI recognition
 10 sequence (GGATCC) into the AatII site (GACGTC starting at nucleotide number 2260) to generate the construct designated pGEM-MB45. The DNA bounded by the restriction enzyme sites of BamHI and AccIII contains the amp promoter, amp RBS, initiating methionine and beta-lactamase
 15 signal peptide. This fragment was used to replace the corresponding fragment from pGEM-MB26 to generate construct pGEM-MB46.

amp gene promoter and signal peptide sequences
 20 5'-GGATCCGGTGGCACTTTTCGGGGAAATGTGCGCGGAACCCCTATTTGTT-
 TATTTTCTAAATACATTCAAATATGTATCCGCTCATGAGACAATAACC-
 25 CTGATAAATGCTTCAATAATATTGAAAAAGGAAGAGT-
 ATG.AGT.ATT.CAA.CAT.TTC.CGT.GTC.GCC.CTT.ATT.-
 met ser ile gln his phe arg val ala leu ile
 30 CCC.TTT.TTT.GCG.GCA.TTT.TGC.CTT.CCT.GTT.TTT.-
 pro phe phe ala ala phe cys leu pro val phe
 GCT.CAT.CCG.-3'
 ala his pro....

c) M13-gene-III-signal::bpti::mature-VIII-coat-protein

We may also construct, as depicted in Figure 5, M13-MB51 which would carry a gene encoding a fusion of M13-gene-III-signal-peptide to the previously described
 40 BPTI::mature VIII coat protein. First the BstEII site

FO6250" 55095550

that follows the stop codons of the synthetic gene VIII is changed to an AlwNI site as follows. DNA of pGEM-MB26 is cut with BstEII and the ends filled in by use of Klenow enzyme; a blunt AlwNI linker is ligated to this DNA. This construction is called pGEM-MB26Alw. The XhoI to AlwNI fragment (approximately 300 bp) of pGEM-MB26Alw is purified. RF DNA from phage MK-BPTI (vide infra) is cut with AlwNI and XhoI and the large fragment purified. These two fragments are ligated together; the resulting construction is named M13-MB51. Because M13-MB51 contains no gene III, the phage can not form plaques. M13-MB51 can, however, render cells Km^R. Infectious phage particles can be obtained by use of helper phage. As explained below, the gene III signal sequence is capable of directing (BPTI)::(mature-gene-III-protein) to the surface of phage. In M13-MB51, we have inserted DNA encoding gene VIII coat protein (50 amino acids) and three stop codons 5' to the DNA encoding the mature gene III protein.

Summary of signal peptide fusion protein variants.

20		Promoter	RBS	Signal sequence	Fusion protein
	pGEM-MB26	<u>tac</u>	new	<u>VIII</u>	BPTI/VIII-coat
	pGEM-MB42	<u>tac</u>	new	<u>phoA</u>	BPTI/VIII-coat
	pGEM-MB46	<u>amp</u>	<u>amp</u>	<u>amp</u>	BPTI/VIII-coat
25	pGEM-MB51	<u>III</u>	<u>III</u>	<u>III</u>	BPTI/VIII-coat
	M13 MB48	<u>tac</u>	new	<u>phoA</u>	BPTI/VIII-coat

2. Analysis of the Protein Products Encoded by the Synthetic (signal-peptide::mature-bpti::viii-coat-protein)

30 Genes

i) In vitro analysis

A coupled transcription/translation prokaryotic system (Amersham Corp., Arlington Heights, IL) was

utilized for the in vitro analysis of the protein products encoded by the BPTI/VIII synthetic gene and the variants derived from this.

Table 107 lists the protein products encoded by the 5 listed vectors which are visualized by the standard method of fluorography following in vitro synthesis in the presence of ^{35}S -methionine and separation of the products using SDS polyacrylamide gel electrophoresis. In each sample a pre-beta-lactamase product (approximately 31 kd) 10 can be seen. This is derived from the amp gene which is the common selection gene for each of the vectors. In addition, a (pre-BPTI/VIII) product encoded by the synthetic gene and variants can be seen as indicated. The migration of these species (approximately 14.5 kd) is 15 consistent with the expected size of the encoded proteins.

ii) In vivo analysis.

The vectors detailed in sections (B) and (C) were freshly transfected into the E. coli strain XL1-blue (TM) (Stratagene, La Jolla, CA) and in strain SEF'. E. coli 20 strain SE6004 (LISS85) carries the prlA4 mutation and is more permissive in secretion than strains that carry the wild-type prlA allele. SE6004 is F^- and is deleted for lacI; thus the cells can not be infected by M13 and lacUV5 and tac promoters can not be regulated with IPTG. Strain 25 SEF' is derived from strain SE6004 (LISS85) by crossing with XL1-Blue(TM); the F' in XL1-Blue(TM) carries Tc^{R} and lacI^q. SE6004 is streptomycin^R, Tc^{S} while XL1-Blue(TM) is streptomycin^S, Tc^{R} so that both parental strains can be killed with the combination of Tc and streptomycin. SEF' 30 retains the secretion-permissive phenotype of the parental strain, SE6004(prlA4).

The fresh transfectants were grown in NZYCM medium (SAMB89) for 1 hour after which IPTG was added over the range of concentrations 1.0 μM to 0.5 mM (to derepress the











106290" 55095360

lacUV5 and tac promoters) and grown for an additional 1.5 hours.

Aliquots of the bacterial cells expressing the synthetic insert encoded proteins together with the appropriate controls (no vector, vector with no insert and zero IPTG) were lysed in SDS gel loading buffer and electrophoresed in 20% polyacrylamide gels containing SDS and urea. Duplicate gels were either silver stained (Daiichi, Tokyo, Japan) or electrotransferred to a nylon matrix (Immobilon from Millipore, Bedford, MA) for western analysis by standard means using rabbit anti-BPTI polyclonal antibodies.

Table 108 lists the interesting proteins visualized on a silver stained gel and by western analysis of an identical gel. We can see clearly in the western analysis that protein species containing BPTI epitopes are present in the test strains which are absent from the control strains and which are also IPTG inducible. In XL1-Blue(TM), the migration of this species is predominantly that of the unprocessed form of the pro-protein although a small proportion of the encoded proteins appear to migrate at a size consistent with that of a fully processed form. In SEF', the processed form predominates, there being only a faint band corresponding to the unprocessed species.

Thus in strain SEF', we have produced a tripartite fusion protein that is specifically cleaved after the secretion signal sequence. We believe that the mature protein comprises BPTI followed by the gene VIII coat protein and that the coat protein moiety spans the membrane. We believe that it is highly likely that one or more copies, perhaps hundreds of copies, of this protein will co-assemble into M13 derived phage or M13-like phagemids. This construction will allow us to a) mutagenize the BPTI domain, b) display each of the variants on

15 carrying a gene that codes for the variant of BPTI
displayed on the phage surface.

i. Phage Production.

25 a phage stock for further infection, phage production and
display of the gene product of interest.

30 incubator at 37°C. To this culture is added a sufficient amount of the phage stock to give a MOI of 10 together with IPTG to give a final concentration of 0.5 mM. The culture is allowed to grow for a further 2 hrs.

ii. Phage Preparation and Purification.

The phage producing bacterial culture is centrifuged to separate the phage in the supernatant from the bacterial pellet. To the supernatant is added one quarter by
 5 volume of phage precipitation solution (20% PEG, 3.75 M ammonium acetate) and PMSF to a final concentration of 1mM. It is left on ice for 2 hours after which the precipitated phage is retrieved by centrifugation. The phage pellet is redissolved in TrisEDTA containing 0.1%
 10 Sarkosyl and left at 4°C for 1 hour after which any bacteria and bacterial debris is removed by centrifugation. The phage in the supernatant is reprecipitated with PEG overnight at 4°C. The phage pellet is resuspended in LB medium and reprecipitated another two times to remove
 15 the detergent. The phage is stored in LB medium at 4°C, titered and used for analysis and binding studies.

A more stringent phage purification scheme involves centrifugation in a CsCl gradient. 3.86 g of CsCl is dissolved in NET buffer (0.1 M NaCl, 1mM EDTA, 0.1M Tris
 20 pH 7.7) upto a volume of 10 ml. 10^{12} to 10^{13} phage in TE Sarkosyl buffer are mixed with 5 ml of CsCl NET buffer and transferred to a sealable ultracentrifuge tube. Centrifugation is performed overnight at 34K rpm in a Sorvall OTD-65B Ultracentrifuge. The tubes are opened and 400 μ l
 25 aliquots are carefully removed. 5 μ l aliquots are removed from the fractions and analysed by agarose gel electrophoresis after heating at 65°C for 15 minutes together with the gel loading buffer containing 0.1% SDS. Fractions containing phage are pooled, the phage reprecipitated and finally redissolved in LB medium to a concentration of 10^{12} to 10^{13} phage per ml.
 30

iii. Phage Analysis.

The display phage, together with appropriate controls are analyzed using standard methods of polyacrylamide gel

T06290" 5096860

- electrophoresis and either silver staining of the gel or electrotransfer to a nylon matrix followed by analysis with anti-BPTI antiserum (Western analysis). Quantitation of the display of heterologous proteins is achieved by
- 5 running a serial dilution of the starting protein, for example BPTI, together with the display phage samples in the electrophoresis and Western analyses described above. An alternative method involves running a 2 fold serial dilution of a phage in which both the major coat protein
 - 10 and the fusion protein are visualized by silver staining. A comparison of the relative ratios of the two protein species allows one to estimate the number of fusion proteins per phage since the number of VIII gene encoded proteins per phage (approximately 3000) is known.
 - 15 Incorporation of fusion protein into bacteriophage.

In vivo expression of the processed BPTI:VIII fusion protein, encoded by vectors GemMB42 (above and Table 113) and M13MB48 (above), implied that the processed fusion product was likely to be correctly located within the

- 20 bacterial cell membrane. This localization made it possible that it could be incorporated into the phage and that the BPTI moiety would be displayed at the bacteriophage surface.

- SEF' cells were infected with either M13MB48 (con-
- 25 sisting of the starting phage vector M13mp18, altered as described above, containing the synthetic gene consisting of a tac promoter, functional ribosome binding site, phoA signal peptide, mature BPTI and mature major coat protein) or M13mp18, as a control. Phage infections, preparation
 - 30 and purification was performed as described in Example VIII.

The resulting phage were electrophoresed (approximately 10^{11} phage per lane) in a 20% polyacrylamide gel containing urea followed by electrotransfer to a nylon

F06290" 5609660

matrix and western analysis using anti-BPTI rabbit serum. A single species of protein was observed in phage derived from infection with the M13MB48 stock phage which was not observed in the control infection. This protein had a migration of about 12 kd, consistent with that of the fully processed fusion protein.

Western analysis of SEF' bacterial lysate with or without phage infection demonstrated another species of protein of about 20kd. This species was also present, to a lesser degree, in phage preparations which were simply PEG precipitated without further purification (for example, using nonionic detergent or by CsCl gradient centrifugation). A comparison of M13MB48 phage preparations made in the presence or absence of detergent demonstrated that sarkosyl treatment and CsCl gradient purification did remove the bacterial contaminant while having no effect on the presence of the BPTI:VIII fusion protein. This indicates that the fusion protein has been incorporated and is a constituent of the phage body.

The time course of phage production and BPTI:VIII incorporation was followed post-infection and after IPTG induction. Phage production and fusion protein incorporation appeared to be maximal after two hours. This time course was utilized in further phage productions and analyses.

Polyacrylamide electrophoresis of the phage preparations, followed by silver staining, demonstrated that the preparations were essentially free of contaminating protein species and that an extra protein band was present in M13MB48 derived phage which was not present in the control phage. The size of the new protein was consistent with that seen by western analysis. A similar analysis of a serially diluted BPTI:VIII incorporated phage demonstrated that the ratio of fusion protein to major coat

protein was typically in the range of 1:150. Since the phage is known to contain in the order of 3000 copies of the gene VIII product, this means that the phage population contains, on average, 10's of copies of the fusion protein per phage.

Altering the initiating methionine of the natural gene VIII.

The OCV M13MB48 contains the synthetic gene encoding the BPTI:VIII fusion protein in the intergenic region of the modified M13mp18 phage vector. The remainder of the vector consists of the M13 genome which contains the genes necessary for various bacteriophage functions, such as DNA replication and phage formation etc. In an attempt to increase the phage incorporation of the fusion protein, we decided to try to diminish the production of the natural gene VIII product, the major coat protein, by altering the codon for the initiating methionine of this gene to one encoding leucine. In such cases, methionine is actually incorporated, but the rate of initiation is reduced. The change was achieved by standard methods of site-specific oligonucleotide mutagenesis as follows.

 M K K S -rest of VIII
 ACT.TCC.TC.ATG.AAA.AAG.TCT.
25 rest of XI - T S S stop

Site-specific mutagenesis.

 (L) K K S -rest of VIII
 ACT.TCC.AG.CTG.AAA.AAG.TCT.
30 rest of XI - T S S stop

Note that the 3' end of the XI gene overlaps with the 5' end of the VIII gene. Changes in DNA sequence were designed such that the desired change in the VIII gene product could be achieved without alterations to the predicted amino acid sequence of the gene XI product. A diagnostic PvuII recognition site was introduced at this

site.

It was anticipated that initiation of the natural gene VIII product would be hindered, enabling a higher proportion of the fusion protein to be incorporated into the resulting phage.

Analyses of the phage derived from this modified vector indicated that there was a significant increase in the ratio of fusion protein to major coat protein. Quantitative estimates indicated that within a phage population as much as 100 copies of the BPTI:VIII fusion were incorporated per phage.

Incorporation of interdomain extension fusion proteins into phage.

A phage pool containing a variegated pentapeptide extension at the BPTI:coat protein interface (see Example VII) was used to infect SEF' cells. IPTG induction, phage production and preparation were as described in Example VIII. Using the criteria detailed in the previous section, it was determined that extended fusion proteins were incorporated into phage. Gel electrophoresis of the generated phage, followed by either silver staining or western analysis with anti-BPTI rabbit serum, demonstrated fusion proteins that migrated similarly to but discernably slower than that of the starting fusion protein.

With regard to the 'EGGS linker' extensions of the domain interface, individual phage stocks predicted to contain one or more 5-amino-acid unit extensions were analyzed in a similar fashion. The migration of the extended fusion proteins were readily distinguishable from the parent fusion protein when viewed by western analysis or silver staining. Those clones analyzed in more detail included M13.3X4 (which contains a single inverted EGGS linker with a predicted amino acid sequence of GGGS),

- M13.3X7 (which contains a correctly orientated linker with a predicted amino acid sequence of EGGGS), M13.3X11 (which contains 3 linkers with an inversion and a predicted amino acid sequence for the extension of EGGGSGSSSLGSSSL) and
- 5 M13.3Xd which contains an extension consisting of at least 5 linkers or 25 amino acids.

The extended fusion proteins were all incorporated into phage at high levels (on average 10's of copies per phage were present and when analyzed by gel electrophoresis migrated rates consistent with the predicted size of the extension. Clones M13.3X4 and M13.3X7 migrated at a position very similar to but discernably different from the parent fusion protein, while M13.3X11 and M13.3Xd were markedly larger.

- 15 Display of BPTI:VIII fusion protein by bacteriophage.

The BPTI:VIII fusion protein had been shown to be incorporated into the body of the phage. This phage was analyzed further to demonstrate that the BPTI moiety was accessible to specific antibodies and hence displayed at

20 the phage surface.

The assay is detailed in section EE, but principally involves the addition of purified anti-BPTI IgG (from the serum of BPTI injected rabbits) to a known titer of phage. Following incubation, protein A-agarose beads are added to

25 bind the IgG and left to incubate overnight. The IgG-protein A beads and any bound phage are removed by centrifugation followed by a retitering of the supernatant to determine any loss of phage. The phage bound to the beads can be acid eluted and titered also. Appropriate

30 controls are included in the assay, such as a wild type phage stock (M13mp18) and IgG purified from normal rabbit pre-immune serum.

Table 140 shows that while the titer of the wild type

phage is unaltered by the presence of anti-BPTI IgG, BPTI-IIIIMK (the positive control for the assay), demonstrated a significant drop in titer with or without the extra addition of protein A beads. (Note that since the BPTI moiety is part of the III gene product which is involved in the binding of phage to bacterial pili, such a phenomenon is entirely expected.) Two batches of M13MB48 phage (containing the BPTI:VIII fusion protein) demonstrated a significant reduction in titer, as judged by plaque forming units, when anti-BPTI antibodies and protein A beads were added to the phage. The initial drop in titer with the antibody alone, differs somewhat between the two batches of phage. This may be a result of experimental or batch variation. Retrieval of the immunoprecipitated phage, while not quantitative, was significant when compared to the wild type phage control.

Further control experiments relating to this section are shown in Table 141 and Table 142. The data demonstrated that the loss in titer observed for the BPTI:VIII containing phage is a result of the display of BPTI epitopes by these phage and the specific interaction with anti-BPTI antibodies. No significant interaction with either protein A agarose beads or IgG purified from normal rabbit serum could be demonstrated. The larger drop in titer for M13MB48 batch five reflects the higher level incorporation of the fusion protein in this preparation.

Functionality of the BPTI moiety in the BPTI-VIII display phage.

The previous two sections demonstrated that the BPTI:VIII fusion protein has been incorporated into the phage body and that the BPTI moiety is displayed at the phage surface. To demonstrate that the displayed molecule is functional, binding experiments were performed in a manner almost identical to that described in the previous

section except that proteases were used in place of antibodies. The display phage, together with appropriate controls, are allowed to interact with immobilized proteases or immobilized inactivated proteases. Binding
5 can be assessed by monitoring the loss in titer of the display phage or by determining the number of phage bound to the respective beads.

Table 143 shows the results of an experiment in which BPTI.VIII display phage, M13MB48, were allowed to bind to
10 anhydrotrypsin-agarose beads. There was a significant drop in titer when compared to wild type phage, which do not display BPTI. A pool of phage (5AA Pool), each contain a variegated 5 amino acid extension at the BPTI:major coat protein interface, demonstrated a similar
15 decline in titer. In a control experiment (table 143) very little non-specific binding of the above display phage was observed with agarose beads to which an unrelated protein (streptavidin) is attached.

Actual binding of the display phage is demonstrated
20 by the data shown for two experiments in Table 144. The negative control is wild type M13mp18 and the positive control is BPTI-IIIIMK, a phage in which the BPTI moiety, attached to the gene III protein, has been shown to be displayed and functional. M13MB48 and M13MB56 both bind
25 to anhydrotrypsin beads in a manner comparable to that of the positive control, being 40 to 60 times better than the negative control (non-display phage). Hence functionality of the BPTI moiety, in the major coat fusion protein, was established.

30 To take this analysis one step further, a comparison of phage binding to active and inactivated trypsin is shown in Table 145. The control phage, M13mp18 and BPTI-III MK, demonstrated binding similar to that detailed in Example III. Note that the relative binding is enhanced

with trypsin due to the apparent marked reduction in the non-specific binding of the wild type phage to the active protease. M13.3X7 and M13.3X11, which both contain 'EGGGS' linker extensions at the domain interface, bound to anhydrotrypsin and trypsin in a manner similar to BPTI-IIIMK phage. The binding, relative to non-display phage, was approximately 100 fold higher in the anhydrotrypsin binding assay and at least 1000 fold higher in the trypsin binding assay. The binding of another 'EGGGS' linker variant (M13.3Xd) was similar to that of M13.3X7.

To demonstrate the specificity of binding the assays were repeated with human neutrophil elastase (HNE) beads and compared to that seen with trypsin beads Table 146. BPTI has a very high affinity for trypsin and a low affinity for HNE, hence the BPTI display phage should reflect these affinities when used in binding assays with these beads. The negative and positive controls for trypsin binding were as already described above while an additional positive control for the HNE beads, BPTI(K15L,-MGNG)-III MA (see Example III) was included. The results, shown in Table 146, confirmed this prediction. M13MB48, M13.3X7 and M13.3X11 phage demonstrated good binding to trypsin, relative to wild type phage and the HNE control (BPTI(K15L,MGNG)-III MA), being comparable to BPTI-IIIMK phage. Conversely poor binding occurred when HNE beads were used, with the exception of the HNE positive control phage.

Taken together the accumulated data demonstrated that when BPTI is part of a fusion protein with the major coat protein of M13 phage, the molecule is both displayed at the surface of the phage and a significant proportion of it is functional in a specific protease binding manner.

--- *** ---

EXAMPLE II

CONSTRUCTION OF BPTI/GENE-III DISPLAY VECTOR

DNA manipulations were conducted according to standard procedures as described in Maniatis *et al.* (MANI82). First the unwanted lacZ gene of M13-MB1/2 was removed. M13-MB1/2 RF was cut with BamHI and SalI and the large fragment was isolated by agarose gel electrophoresis. The recovered 6819 bp fragment was filled in with Klenow fragment of E. coli DNA polymerase and ligated to a synthetic HindIII 8mer linker (CAAGCTTG). The ligation sample was used to transfect competent XL1-Blue^(TM) (Stratagene, La Jolla, CA) cells which were subsequently plated for plaque formation. RF DNA was prepared from chosen plaques and a clone, M13-MB1/2-delta, containing regenerated BamHI and SalI sites as well as a new HindIII site, all 500 bp upstream of the BglIII site (6935) was picked.

A unique NarI site was introduced into codons 17 and 18 of gene III (changing the amino acids from H-S to G-A, *Cf.* Table 110). 10⁶ phage produced from bacterial cells harboring the M13-MB1/2-delta RF DNA were used to infect a culture of CJ236 cells (relevant genotype: F', dut1, ung1, Cm^R) (OD595=0.35). Following overnight incubation at 37°C, phage were recovered and uracil-containing ss DNA was extracted from phage in accord with the instructions for the MUTA-GENE^(R) M13 *in vitro* Mutagenesis Kit (Catalogue Number 170-3571, Bio-Rad, Richmond, CA). Two hundred nanograms of the purified single stranded DNA was annealed to 3 picomoles of a phosphorylated 25mer mutagenic oligonucleotide,

5'-gtttcagcggCgCCagaatagaaag-3',

where upper case indicates the changes). Following filling in with T4 DNA polymerase and ligation with T4 DNA

T06290-5609660

ligase, the reaction sample was used to transfect competent XL1-Blue(TM) cells which were subsequently plated to permit the formation of plaques.

RF DNA, isolated from phage-infected cells which had
 5 been allowed to propagate in liquid culture for 8 hours,
 was denatured, spotted on a Nytran membrane, baked and
 hybridized to the 25mer mutagenic oligonucleotide which
 had previously been phosphorylated with ³²P-ATP. Clones
 exhibiting strong hybridization signals at 70°C (6°C less
 10 than the theoretical T_m of the mutagenic oligonucleotide)
 were chosen for large scale RF preparation. The presence
 of a unique NarI site at nucleotide 1630 was confirmed by
 restriction enzyme analysis. The resultant RF DNA, M13-
 MB1/2-delta-NarI was cut with BamHI, dephosphorylated with
 15 calf intestinal phosphatase, and ligated to a 1.3 Kb BamHI
 fragment, encoding the kanamycin-resistance gene (kan),
 derived from plasmid pUC4K (Pharmacia, Piscataway, NJ).
 The ligation sample was used to transfect competent
 XL1-Blue(TM) cells which were subsequently plated onto LB
 20 plates containing kanamycin (Km). RF DNA prepared from
 Km^R colonies was prepared and subjected to restriction
 enzyme analysis to confirm the insertion of kan into M13-
 MB1/2-delta-NarI DNA thereby creating the phage MK. Phage
 MK grows as well as wild-type M13, indicating that the
 25 changes at the cleavage site of gene III protein are not
 detectably deleterious to the phage.

INSERTION OF SYNTHETIC BPTI GENE

The construction of the BPTI-III expression vector is
 shown in Figure 6. The synthetic bpti-VIII fusion
 30 contains a NarI site that comprises the last two codons of
 the BPTI-encoding region. A second NarI site was intro-
 duced upstream of the BPTI-encoding region as follows. RF
 DNA of phage M13-MB26 was cut with AccIII and ligated to
 the dsDNA adaptor:

5'-TATTCTGGCGCCCGT -3'
 3'-ATAAGACCGCGGGCAGGCC-5'
|NarI| |AccIII|

- 5 The ligation sample was subsequently restricted with NarI and a 180 bp DNA fragment encoding BPTI was isolated by agarose gel electrophoresis. RF DNA of phage MK was digested with NarI, dephosphorylated with calf intestinal phosphatase and ligated to the 180 bp fragment. Ligation
- 10 samples were used to transfect competent XL1-Blue(TM) cells which were plated to enable the formation of plaques. DNA, isolated from phage derived from plaques, was denatured, applied to a Nytran membrane, baked and hybridized to a ³²P-phosphorylated double stranded DNA
- 15 probe corresponding to the BPTI gene. Large scale RF preparations were made for clones exhibiting a strong hybridization signal. Restriction enzyme digestion analysis confirmed the insertion of a single copy of the synthetic BPTI gene into gene III of MK to generate phage
- 20 MK-BPTI. Subsequent DNA sequencing confirmed that the sequence of the bpti-III fusion gene is correct and that the correct reading frame is maintained (Table 111). Table 116 shows the entire coding region, the translation into protein sequence, and the functional parts of the
- 25 polypeptide chain.

EXPRESSION OF THE BPTI-III FUSION GENE IN VITRO

- MK-BPTI RF DNA was added to a coupled prokaryotic transcription-translation extract (Amersham). Newly synthesized radiolabelled proteins were produced and
- 30 subsequently separated by electrophoresis on a 15% SDS-polyacrylamide gel subjected to fluorography. The MK-BPTI DNA directs the synthesis of an unprocessed gene III fusion protein which is 7 Kd larger than the gene III product encoded by MK. This is consistent with the
- 35 insertion of 58 amino acids of BPTI into the gene III

106290-56096860

protein. Immunoprecipitation of radiolabelled proteins generated by the cell-free prokaryotic extract was conducted. Neither rabbit anti(M13-gene-VIII-protein) IgG nor normal rabbit IgG were able to immunoprecipitate the gene III protein encoded by either MK or MK-BPTI. However, rabbit anti-BPTI IgG is able to immunoprecipitate the gene III protein encoded by MK-BPTI but not by MK. This confirms that the increase in size of the III protein encoded by MK-BPTI is attributable to the insertion of the BPTI protein.

WESTERN ANALYSIS

Phage were recovered from bacterial cultures by PEG precipitation. To remove residual bacterial cells, recovered phage were resuspended in a high salt buffer and subjected to centrifugation, in accord with the instructions for the MUTA-GENE(R) M13 in vitro Mutagenesis Kit (Catalogue Number 170-3571, Bio-Rad, Richmond, CA). Aliquots of phage (containing up to 40 µg of protein) were subjected to electrophoresis on a 12.5% SDS-urea-polyacrylamide gel and proteins were transferred to a sheet of Immobilon by electro-transfer. Western blots were developed using rabbit anti-BPTI serum, which had previously been incubated with an E. coli extract, followed by goat anti-rabbit antibody conjugated to alkaline phosphatase. An immunoreactive protein of 67 Kd is detected in preparations of the MK-BPTI but not the MK phage. The size of the immunoreactive protein is consistent with the predicted size of a processed BPTI-III fusion protein (6.4 Kd plus 60 Kd). These data indicate that BPTI-specific epitopes are presented on the surface of the MK-BPTI phage but not the MK phage.

NEUTRALIZATION OF PHAGE TITER WITH AGAROSE-IMMOBILIZED ANHYDRO-TRYPSIN

Anhydro-trypsin is a derivative of trypsin in which the active site serine has been converted to dehydroalanine. Anhydro-trypsin retains the specific binding of trypsin but not the protease activity. Unlike polyclonal-
 5 antibodies, anhydro-trypsin is not expected to bind unfolded BPTI or incomplete fragments.

Phage MK-BPTI and MK were diluted to a concentration $1.4 \cdot 10^{12}$ particles per ml. in TBS buffer (PARM88) containing 1.0 mg/ml BSA. Thirty microliters of diluted phage
 10 were added to 2, 5, or 10 microliters of a 50% slurry of agarose-immobilized anhydro-trypsin (Pierce Chemical Co., Rockford, IL) in TBS/BSA buffer. Following incubation at 25°C, aliquots were removed, diluted in ice cold LB broth and titered for plaque-forming units on a lawn of XL1-
 15 Blue(TM) cells. Table 114 illustrates that incubation of the MK-BPTI phage with immobilized anhydro-trypsin results in a very significant loss in titer over a four hour period while no such effect is observed with the MK (control) phage. The reduction in phage titer is also
 20 proportional to the amount of immobilized anhydro-trypsin added to the MK-BPTI phage. Incubation with five microliters of a 50% slurry of agarose-immobilized streptavidin (Sigma, St. Louis, MO) in TBS/BSA buffer does not reduce the titer of either the MK-BPTI or MK phage. These data
 25 are consistent with the presentation of a correctly-folded, functional BPTI protein on the surface of the MK-BPTI phage but not on the MK phage. Unfolded or incomplete BPTI domains are not expected to bind anhydro-trypsin. Furthermore, unfolded BPTI domains are expected
 30 to be non-specifically sticky.

NEUTRALIZATION OF PHAGE TITER WITH ANTI-BPTI ANTIBODY

MK-BPTI and MK phage were diluted to a concentration of $4 \cdot 10^8$ plaque-forming units per ml in LB broth. Fifteen microliters of diluted phage were added to an

T05250-56096850

equivalent volume of either rabbit anti-BPTI serum or normal rabbit serum (both diluted 10 fold in LB broth). Following incubation at 37°C, aliquots were removed, diluted by 10^4 in ice-cold LB broth and titered for plaque-forming units on a lawn of XL1-Blue^(TM) cells. Incubation of the MK-BPTI phage with anti-BPTI serum results in a steady loss in titer over a two hour period while no such effect is observed with the MK phage. As expected, normal rabbit serum does not reduce the titer of either the MK-BPTI or the MK phage. Prior incubation of the anti-BPTI serum with authentic BPTI protein but not with an equivalent amount of E. coli protein, blocks the ability of the serum to reduce the titer of the MK-BPTI phage. This data is consistent with the presentation of BPTI-specific epitopes on the surface of the MK-BPTI phage but not the MK phage. More specifically, the data indicates that these BPTI epitopes are associated with the gene III protein and that association of this fusion protein with an anti-BPTI antibody blocks its ability to mediate the infection of bacterial cells.

NEUTRALIZATION OF PHAGE TITER WITH TRYPSIN

MK-BPTI and MK phage were diluted to a concentration of $4 \cdot 10^8$ plaque-forming units per ml in LB broth. Diluted phage were added to an equivalent volume of trypsin diluted to various concentrations in LB broth. Following incubation at 37°C, aliquots were removed, diluted by 10^4 in ice cold LB broth and titered for plaque-forming units on a lawn of XL1-Blue^(TM) cells. Incubation of the MK-BPTI phage with 0.15 μ g of trypsin results in a 70% loss in titer after a two hour period while only a 15% loss in titer is observed for the MK phage. A reduction in the amount of trypsin added to phage results in a reduction in the loss of titer. However, at all trypsin concentrations investigated, the

MK-BPTI phage are more sensitive to incubation with trypsin than the MK phage. An interpretation of this data is that association of the BPTI-III fusion protein displayed on the surface of the MK-BPTI phage with trypsin blocks its ability to mediate the infection of bacterial cells.

The reduction in titer of phage MK by trypsin is an example of a phenomenon that is likely to be general: proteases, if present in sufficient quantity, will degrade proteins on the phage and reduce infectivity. The present application lists several means that can be used to overcome this problem.

AFFINITY SELECTION SYSTEM

Affinity Selection with Immobilized Anhydro-Trypsin

MK-BPTI and MK phage were diluted to a concentration of $1.4 \cdot 10^{12}$ particles per ml in TBS buffer (PARM88) containing 1.0 mg/ml BSA. We added $4.0 \cdot 10^{10}$ phage to 5 microliters of a 50% slurry of either agarose-immobilized anhydro-trypsin beads (Pierce Chemical Co.) or agarose-immobilized streptavidin beads (Sigma) in TBS/BSA. Following a 3 hour incubation at room temperature, the beads were pelleted by centrifugation for 30 seconds at 5000 rpm in a microfuge and the supernatant fraction was collected. The beads were washed 5 times with TBS/Tween buffer (PARM88) and after each wash the beads were pelleted by centrifugation and the supernatant was removed. Finally, beads were resuspended in elution buffer (0.1 N HCl containing 1.0 mg/ml BSA adjusted to pH 2.2 with glycine) and following a 5 minute incubation at room temperature, the beads were pelleted by centrifugation. The supernatant was removed and neutralized by the addition of 1.0 M Tris-HCl buffer, pH 8.0.

Aliquots of phage samples were applied to a Nytran

T05290-50036860

membrane using a Schleicher and Schuell (Keene, NH) filtration minifold and phage DNA was immobilized onto the Nytran by baking at 80°C for 2 hours. The baked filter was incubated at 42°C for 1 hour in pre-wash solution
 5 (MANI82) and pre-hybridization solution (5Prime-3Prime, West Chester, PA). The 1.0 Kb NarI (base 1630)/XmnI (base 2646) DNA fragment from MK RF was radioactively labelled with ³²P-dCTP using an oligolabelling kit (Pharmacia, Piscataway, NJ). The radioactive probe was added to the
 10 Nytran filter in hybridization solution (5Prime-3Prime) and, following overnight incubation at 42°C, the filter was washed and subjected to autoradiography.

The efficiency of this affinity selection system can be semi-quantitatively determined using the dot-blot
 15 procedure described elsewhere in the present application. Exposure of MK-BPTI-phage-treated anhydro-trypsin beads to elution buffer releases bound MK-BPTI phage. Streptavidin beads do not retain phage MK-BPTI. Anhydro-trypsin beads do not retain phage MK. In the experiment depicted in
 20 Table 115, we estimate that 20% of the total MK-BPTI phage were bound to 5 microliters of the immobilized anhydro-trypsin and were subsequently recovered by washing the beads with elution buffer (pH 2.2 HCl/glycine). Under the same conditions, no detectable MK-BPTI phage were bound
 25 and subsequently recovered from the streptavidin beads. The amount of MK-BPTI phage recovered in the elution fraction is proportional to the amount of immobilized anhydro-trypsin added to the phage. No detectable MK phage were bound to either the immobilized anhydro-
 30 trypsin or streptavidin beads and no phage were recovered with elution buffer. These data indicate that the affinity selection system described above can be utilized to select for phage displaying a specific folded protein (in this case, BPTI). Unfolded or incomplete BPTI domains
 35 are not expected to bind anhydro-trypsin.

FD-250 (Rev. 5-22-64)

Affinity Selection with Anti-BPTI antibodies

MK-BPTI and MK phage were diluted to a concentration of $1 \cdot 10^{10}$ particles per ml in Tris buffered saline solution (PARM88) containing 1.0 mg/ml BSA. Two $\cdot 10^8$ phage were added to 2.5 μ g of either biotinylated rabbit anti-BPTI IgG in TBS/BSA or biotinylated rabbit anti-mouse antibody IgG (Sigma) in TBS/BSA, and incubated overnight at 4°C. A 50% slurry of streptavidin-agarose (Sigma), washed three times with TBS buffer prior to incubation with 30 mg/ml BSA in TBS buffer for 60 minutes at room temperature, was washed three times with TBS/Tween buffer (PARM88) and resuspended to a final concentration of 50% in this buffer. Samples containing phage and biotinylated IgG were diluted with TBS/Tween prior to the addition of streptavidin-agarose in TBS/Tween buffer. Following a 60 minute incubation at room temperature, streptavidin-agarose beads were pelleted by centrifugation for 30 seconds and the supernatant fraction was collected. The beads were washed 5 times with TBS/Tween buffer and after each wash, the beads were pelleted by centrifugation and the supernatant was removed. Finally, the streptavidin-agarose beads were resuspended in elution buffer (0.1 N HCl containing 1.0 mg/ml BSA adjusted to pH 2.2 with glycine), incubated 5 minute at room temperature, and pelleted by centrifugation. The supernatant was removed and neutralized by the addition of 1.0 M Tris-HCl buffer, pH 8.0.

Aliquots of phage samples were applied to a Nytran membrane using a Schleicher and Schuell minifold apparatus. Phage DNA was immobilized onto the Nytran by baking at 80°C for 2 hours. Filters were washed for 60 minutes in pre-wash solution (MANI82) at 42°C then incubated at 42°C for 60 minutes in Southern pre-hybridization solution (5Prime-3Prime). The 1.0 Kb NarI (1630bp)/XmnI (2646 bp)

DNA fragment from MK RF was radioactively labelled with ^{32}P - αdCTP using an oligolabelling kit (Pharmacia, Piscataway, NJ). Nytran membranes were transferred from pre-hybridization solution to Southern hybridization solution (5Prime-3Prime) at 42°C . The radioactive probe was added to the hybridization solution and following overnight incubation at 42°C , the filter was washed 3 times with $2 \times \text{SSC}$, 0.1% SDS at room temperature and once at 65°C in $2 \times \text{SSC}$, 0.1% SDS. Nytran membranes were subjected to autoradiography. The efficiency of the affinity selection system can be semi-quantitatively determined using the above dot blot procedure. Comparison of dots A1 and B1 or C1 and D1 indicates that the majority of phage did not stick to the streptavidin-agarose beads. Washing with TBS/Tween buffer removes the majority of phage which are non-specifically associated with streptavidin beads. Exposure of the streptavidin beads to elution buffer releases bound phage only in the case of MK-BPTI phage which have previously been incubated with biotinylated rabbit anti-BPTI IgG. This data indicates that the affinity selection system described above can be utilized to select for phage displaying a specific antigen (in this case BPTI). We estimate an enrichment factor of at least 40 fold based on the calculation

$$\text{Enrichment Factor} = \frac{\text{Percent MK-BPTI phage recovered}}{\text{Percent MK phage recovered}}$$

EXAMPLE III

CHARACTERIZATION AND FRACTIONATION OF CLONALLY PURE POPULATIONS OF PHAGE, EACH DISPLAYING A SINGLE CHIMERIC APROTININ HOMOLOGUE/M13 GENE III PROTEIN:

This Example demonstrates that chimeric phage

DNA fragment from MK RF was radioactively labelled with ^{32}P - αdCTP using an oligolabelling kit (Pharmacia, Piscataway, NJ). Nytran membranes were transferred from pre-hybridization solution to Southern hybridization solution (5Prime-3Prime) at 42°C . The radioactive probe was added to the hybridization solution and following overnight incubation at 42°C , the filter was washed 3 times with $2 \times \text{SSC}$, 0.1% SDS at room temperature and once at 65°C in $2 \times \text{SSC}$, 0.1% SDS. Nytran membranes were subjected to autoradiography. The efficiency of the affinity selection system can be semi-quantitatively determined using the above dot blot procedure. Comparison of dots A1 and B1 or C1 and D1 indicates that the majority of phage did not stick to the streptavidin-agarose beads. Washing with TBS/Tween buffer removes the majority of phage which are non-specifically associated with streptavidin beads. Exposure of the streptavidin beads to elution buffer releases bound phage only in the case of MK-BPTI phage which have previously been incubated with biotinylated rabbit anti-BPTI IgG. This data indicates that the affinity selection system described above can be utilized to select for phage displaying a specific antigen (in this case BPTI). We estimate an enrichment factor of at least 40 fold based on the calculation

$$\text{Enrichment Factor} = \frac{\text{Percent MK-BPTI phage recovered}}{\text{Percent MK phage recovered}}$$

EXAMPLE III

CHARACTERIZATION AND FRACTIONATION OF CLONALLY PURE POPULATIONS OF PHAGE, EACH DISPLAYING A SINGLE CHIMERIC APROTININ HOMOLOGUE/M13 GENE III PROTEIN:

This Example demonstrates that chimeric phage

At position 34, all 20 amino acids were allowed. Fourteen have been seen. LYS appeared seven times, GLU five times, THR four times, LEU three times, GLY, ASP, GLN, MET, ASN, and HIS twice each, and ARG, PRO, VAL, and TYR once each. There were no instances of ALA, CYS, PHE, ILE, SER, or TRP. No homologue of aprotinin with GLU, GLY, or MET at 34 has been reported heretofore. Here, as

At position 34, all 20 amino acids were allowed. Fourteen have been seen. LYS appeared seven times, GLU five times, THR four times, LEU three times, GLY, ASP, GLN, MET, ASN, and HIS twice each, and ARG, PRO, VAL, and TYR once each. There were no instances of ALA, CYS, PHE, ILE, SER, or TRP. No homologue of aprotinin with GLU, GLY, or MET at 34 has been reported heretofore. Here, as

5

40 and 41. Alanine is somewhat preferred at 40; ALA:GLY::19:16. Both ALA and GLY have been reported in aprotinin homologues.

15

25

30

We tested EpiNE7.23 (the sequence closest to consensus) against EpiNE7 on HNE beads. Figure 16 shows the fractionation of strains of phage that display these two

EpINEs. Phage that display EpINE7 are eluted at higher pH than are phage that display EpINE7.23. Furthermore, more of the EpINE7.23 phage are retained than of the EpINE7 phage. Note the peak at pH 2.25 in the EpINE7.23 elution.

- 5 This suggests that EpINE7.23 has a higher affinity for HNE than does EpINE7. In a similar way, we tested EpINE7.4 and found that it is not retained on HNE so well as EpINE7. This is consistent with the fractionation not being complete.

- 10 Further fractionation, characterization of clonally pure EpINE7.nn strains, and biochemical characterization of soluble EpINE7.nn derivatives will reveal which sequences in this collection have the highest affinity for HNE.

- 15 Fractionation of the library involves a number of factors. Differential binding allows phage that display PBDs having the desired binding properties to be enriched. Differences in infectivity, plaque size, and phage yield are related to differences in the sequence of the PBDs,
20 but are not directly correlated to affinity for the target. These factors may reduce the effectiveness of the desired fractionation. An additional factor that may be present is differential abundance of PBD sequences in the initial library. One step we employ to reduce the effect
25 of differential infectivity is to transduce cells with isolated phage rather than to infect them. In the first fractionation, we did not obtain sufficient material for transduction and so infected cells; this fractionation was successful. Because the parental sequence, EpINE7, was
30 selected for a sequence at residues 15 through 19 that confer high affinity for HNE, we believe that many, if not most, members of the KLMUT population have significant affinity for HNE. Thus the present fractionations must separate variants having very high affinity for HNE from

277a

aprotinin homologue to HNE. Different substitutions at these positions is likely to confer different specificity on those derivatives. One of the major advantages of the present invention is that many substitutions at several locations may be tested with an amount of effort not much greater than is required to test a single derivative by previously used methods.

There exist a number of proteases produced by lymphocytes. Neutrophil elastase is not the only lymphocytic protease that degrades elastin. The protease p29 is related to HNE. Screening the MYMUT and KLMUT libraries against immobilized p29 is likely to allow isolation of an aprotinin derivative having high affinity for p29.

EXAMPLE VII

BPTI:VIII BOUNDARY EXTENSIONS.

The aim of this work was to introduce peptide extensions between the C-terminus of the BPTI domain and the N-terminus of the M13 major coat protein within the

those merely having high affinity for HNE. It is perhaps relevant that BPTI-III MK phage are only partially eluted from immobilized trypsin at pH 2.2.; $K_d(\text{trypsin}, \text{BPTI}) = 6.0 \cdot 10^{-14}$ M. Elution of EpiNE7-III MA phage from immobilized HNE gives a peak at about pH 3.5 with some phage appearing at lower pH; $K_d(\text{HNE}, \text{EpiNE7}) \leq 1 \cdot 10^{-11}$ M. We recycled phage that either were eluted at pH 2.0 or that were retained after elution with pH 2.0 buffer. A large percentage of EpiNE7-III MA phage would have been washed away with the fractions at pHs less acid than 2.0. This, together with the marked preferences at positions 39, 36, and 34, strongly suggests that we have successfully fractionated the KLMUT library on the basis of affinity for HNE and that the EpiNE7.nn proteins have higher affinity for HNE than does EpiNE7 or any other reported aprotinin derivative.

Fractionation in a few stringent steps emphasizes the affinity of the PBD and allows isolation of variants that confer a small-plaque phenotype on cells (through low infectivity or by slowing cell growth). More gradual fractionation allows observation of a wider variety of variants that show high affinity and favors sequences that start at low abundance. Gradual fractionation also favors selection of variants that do not confer a small-plaque phenotype; such variants may be easier to work with and are preferred for some purposes. In either case, it is preferred to fractionate until there is a manageable number of distinct isolates and to characterize these isolates as pure clones. Thus, it is desirable, in most cases, to fractionate a library in more than one way.

None have identified positions 39 and 34 as key in determining the affinity and specificity of aprotinin homologues and derivatives for particular serine proteases. None have suggested the tryptophan at 39 or charged amino acids (LYS or GLU) at 34 will enhance binding of an

contained new sequences at this position.

A pool of phages, containing the novel interface pentapeptide extensions, was collected by combining the phage extracted from the plated plaques.

5 2. Adding multiple unit extensions to the fusion protein interface.

10 The M13 gene III product contains 'stalk-like' regions as implied by electron micrographic visualization of the bacteriophage (LOPE85). The predicted amino acid sequence of this protein contains repeating motifs, which include:

glu.gly.gly.gly.ser (EGGGS) seven times
gly.gly.gly.ser (GGGS) three times
glu.gly.gly.gly.thr (EGGGT) once.

15 The aim of this section was to insert, at the domain interface, multiple unit extensions which would mirror the repeating motifs observed in the III gene product.

20 Two synthetic oligonucleotides were designed and custom synthesized. GLY is encoded by four codons (GGN); when translated in the opposite direction, these codons give rise to THR, PRO, ALA, and SER. The third base of these codons was picked so that translation of the oligonucleotide in the opposite direction would encode SER. When annealed the synthetic oligonucleotides give
25 the following unit duplex sequence (an EGGGS linker):

```

          E   G   G   G   S
5' C.GAG.GGA.GGA.GGA.TC 3'
3'   TC.CCT.CCT.CCT.AGG.C 5'
    (L) (S) (S) (S) (G)

```

30

35 The duplex has a common two base pair 5' overhang (GC) at either end of the linker which allows for both the ligation of multiple units and the ability to clone into the unique NarI recognition sequence present in OCV's

TOP SECRET 50000000

fusion protein. The reasons for this were two fold; firstly to alter potential protease cleavage sites at the interdomain boundary (as evidenced by an apparent instability of the fusion protein) and secondly to increase interdomain flexibility.

1) Insertion of a variegated pentapeptide at the BPTI:VIII interface.

The gene shown in Table 113 was modified by insertion of five RVT codons between codon 81 and 82. Two synthetic oligonucleotides were designed and custom synthesized. The first consisted of, from 5' to 3': a) from base 2 of codon 77 to the end of codon 81, b) five copies of RVT, and c) from codon 82 to the second base of codon 94. The second comprised 20 bases complementary to the 3' end of the first oligonucleotide. Each RVT codon allows one of the amino acids [T, N, S, A, D, and G] to be encoded. This variegation codon was picked because: a) each amino acid occurs once, and b) all these amino acids are thought to foster a flexible linker. When annealed, the primed variegated oligonucleotide was converted to double-stranded DNA using standard methods.

The duplex was digested with restriction enzymes SfiI and NarI and the resulting 45 base-pair fragment was ligated into a similarly cleaved OCV, M13MB48 (Example I.1.iii.a). The ligated material was transfected into competent E. coli cells (strain XL1-Blue^(TM)) and plated onto a lawn of the same cells on normal bacterial growth plates to form plaques. The bacteriophage contained within the plaques were analyzed using standard methods of nitrocellulose lifts and probing using a ³²P-labeled oligonucleotide complementary to the DNA sequence encoding the fusion protein interface. Approximately 80% of the plaques probed poorly with this oligonucleotide and hence

M13MB48 and Gem MB42. This site is positioned within 1 codon of the DNA encoding the interface. The cloning of an EGGGS linker (or multiple linker) into the vector NarI site destroys this recognition sequence. Insertion of the
 5 EGGGS linker in reverse orientation leads to insertion of GSSSL into the fusion protein.

Addition of a single EGGGS linker at the NarI site of the gene shown in Table 113 leads to the following gene:

```

10      79  80  80a 80b 80c 80d 80e 81  82  83  84
          G   G   E   G   G   G   S   A   A   E   G
          -----
          GGT.GGC.GAG.GGA.GGA.GGA.TCC.GCC.GCT.GAA.GGT
          -----
  
```

15 Note that there is no preselection for the orientation of the linker(s) inserted into the OCV and that multiple linkers of either orientation (with the predicted EGGGS or GSSSL amino acid sequence) or a mixture of
 20 orientations (inverted repeats of DNA) could occur.

A ladder of increasingly large multiple linkers was established by annealing and ligating the two starting oligonucleotides containing different proportions of 5' phosphorylated and non-phosphorylated ends. The logic
 25 behind this is that ligation proceeds from the 3' unphosphorylated end of an oligonucleotide to the 5' phosphorylated end of another. The use of a mixture of phosphorylated and non-phosphorylated oligonucleotides allows for an element of control over the extent of multiple linker
 30 formation. A ladder showing a range of insert sizes was readily detected by agarose gel electrophoresis spanning 15 bp (1 unit duplex-5 amino acids) to greater than 600 base pairs (40 ligated linkers-200 amino acids).

Large inverted repeats can lead to genetic instability.
 35 bility. Thus we chose to remove them, prior to ligation into the OCV, by digesting the population of multiple

linkers with the restriction enzymes AccIII or XhoI, since the linkers, when ligated 'head-to-head' or 'tail-to--tail', generate these recognition sequences. Such a digestion significantly reduces the range in sizes of the multiple linkers to between 1 and 8 linker units (i.e. between 5 and 40 amino acids in steps of 5), as assessed by agarose gel electrophoresis.

The linkers were ligated (as a pool of different insert sizes or as gel-purified discrete fragments) into NarI cleaved OCVs M13MB48 or GemMB42 using standard methods. Following ligation the restriction enzyme NarI was added to remove the self-ligating starting OCV (since linker insertion destroys the NarI recognition sequence). This mixture was used to transform competent XL-1 blue cells and appropriately plated for plaques (OCV M13MB48) or ampicillin resistant colonies (OCV GemMB42).

The transformants were screened using dot blot DNA analysis with one of two ³²p labeled oligonucleotide probes. One probe consisted of a sequence complementary to the DNA encoding the P1 loop of BPTI while the second had a sequence complementary to the DNA encoding the domain interface region. Suitable linker candidates would probe positively with the first probe and negatively or poorly with the second. Plaque purified clones were used to generate phage stocks for binding analyses and BPTI display while the Rf DNA derived from phage infected bacterial cells was used for restriction enzyme analysis and sequencing. Representative insert sequences of selected clones analyzed are as follows:

30	M13.3X4	(GG)C.GGA.TCC.TCC.TCC.CT(C.GCC) gly ser ser ser leu
35	M13.3X7	(G C.GAG.GGA.GGA.GGA.TC(C.GCC) glu gly gly gly ser
	M13.3X11	(GG)C.GAG.GGA.GGA.GGA.TCC.GGA.TCC.TCC.

glu gly gly gly ser gly ser ser

TCC.CTC.GGA.TCC.TCC.TCC.CT(C.GCCC)
ser leu gly ser ser ser leu

5

These highly flexible oligomeric linkers are believed to be useful in joining a binding domain to the major coat (gene VIII) protein of filamentous phage to facilitate the display of the binding domain on the phage surface. They
10 may also be useful in the construction of chimeric OSPs for other genetic packages as well.

EXAMPLE VIII

BACTERIAL EXPRESSION VECTORS.

The expression vectors were designed for the bacterial production of BPTI analogues resulting from the
5 mutagenesis and screening for variants with specific binding properties. The expression vectors used are derivatives of the OCV's M13MB48 and GemMB42. The conversion was achieved by replacing the first codon of the mature VIII gene (codon 82 as shown in Table 113) with
10 a translational stop codon by site specific mutagenesis.

The salient points of the expression vector composition are identical to that of the parent OCV's, namely a lacUV5 promoter (hence IPTG induction), ribosome binding site, initiating methionine, pho A signal peptide and
15 transcriptional termination signal (see Table 113). The placement of the stop codon allows for the expression of only the first half the fusion protein. The Gem-based expression system, containing the genes encoding BPTI analogues, is stored as plasmid DNA, being freshly
20 transfected into cells for expression of the analogue protein. The M13-based expression system is stored as both RF DNA and as phage stocks. The phage stocks are used to infect fresh bacterial cells for expression of the protein of interest.

T05230-56096860

Bacterial Expression of BPTI and Analogues.

i. Gem-based expression vector and protocol.

The gem-based expression vector is a derivative of the OCV GemMB42 (Example I and Table 113). This vector, at least when it contains the BPTI or analogue genes, has demonstrated a degree of insert instability on prolonged growth in liquid culture. To reduce the risk of this the following protocol is used.

Expression vector DNA (containing the BPTI or analogue gene) is transfected into the *E. coli* strain, XL1-Blue^(TM), which is plated on bacterial plates containing ampicillin and allowed to incubate overnight at 37°C to give a dense population of colonies. The colonies are scraped from the plate with a glass spreader in 1ml of NZCYM medium and combined with the scraped cells from other duplicate plates. This stock of cells is diluted approximately one hundred fold into NZCYM liquid medium containing ampicillin (100µg per ml) and allowed to grow in a shaking incubator to a cell density of approximately half log (absorbance of 0.3 at 600nm). IPTG is added to a final concentration of 0.5 mM and the induced culture allowed to grow for a further two hours when it is processed as described below.

ii. M13-based expression vector and protocol.

The M13-based expression vector is derived from OCV M13MB48 (Example I). The BPTI gene (or analogue) is contained within the intergenic region and its transcription is under the control of a lacUV5 promoter, hence IPTG inducible. The expression vector, containing the gene of interest, is maintained and utilized as a phage stock. This method enables a potentially lethal or deleterious gene to be supplied to a bacterial culture and gene induction to occur only when the bacterial culture has

Regardless of the reason, the error rate is extremely low considering only 1 unexpected alteration was observed after sequencing 20 codons in 19 different clones. Furthermore, the value of such a mutation is not diminished by its accidental nature.

Some of the EpINE clones are identical. The sequences of EpINE1, EpINE3, and EpINE7 appear a total of 4, 6 and 5 times respectively. Assuming the 1745 potentially different DNA sequences encoded by the MYMUT oligonucleotide were present at equal frequency in the fusion phage library, the frequent appearance of the sequences for clones EpINE1, EpINE3, and EpINE7 may have important implications. EpINE1, EpINE3, and EpINE7 fusion phage may display BPTI variants with the highest affinity for HNE of all the 1000 potentially different BPTI variants in the MYMUT library.

An examination of the sequences of the EpINE clones is illuminating. A strong preference for either VAL or ILE at the P1 position (residue 15) is indicated with VAL being favored over ILE by 14 to 6. In the MYMUT library, VAL at position 15 is approximately twice as prevalent as ILE. No examples of LEU, PHE, or MET at the P1 position were observed although the MYMUT oligonucleotide has the potential to encode these residues at P1. This is consistent with the observation that BPTI variants with single amino acid substitutions of LEU, PHE, or MET for LYS₁₅ exhibit a significantly lower affinity for HNE than their counterparts containing either VAL or ILE (BECK88b).

PHE is strongly favored at position 17, appearing in 12 of 20 codons. MET is the second most prominent residue at this position but it only appears when VAL is present at position 15. At position 18 PHE was observed in all 20 clones sequenced even though the MYMUT oligonucleotide is capable of encoding other residues at this position. This

result is quite surprising and could not be predicted from previous mutational analysis of BPTI, model building, or on any theoretical grounds. We infer that the presence of PHE at position 18 significantly enhances the ability each of the EpiNEs to bind to HNE. Finally at position 19, PRO appears in 10 of 20 codons while SER, the second most prominent residue, appears at 6 of 20 codons. Of the residues targeted for mutagenesis in the present study, residue 19 is the nearest to the edge of the interaction surface of a PEPI with HNE. Nevertheless, a preponderance of PRO is observed and may indicate that PRO at 19, like PHE at 18, enhances the binding of these proteins to HNE. Interestingly, EpiNE5 appears only once and differs from EpiNE1 only at position 19; similarly, EpiNE6 differs from EpiNE3 only at position 19. These alterations may have only a minor effect on the ability of these proteins to interact with HNE. This is supported by the fact that the pH elution profiles for EpiNE5 and EpiNE6 are very similar to those of EpiNE1 and EpiNE3 respectively.

Only EpiNE2 and EpiNE8 exhibit pH profiles which differ from those of the other selected clones. Both clones contain LYS at position 19 which may restrict the interaction of BPTI with HNE. However, we can not exclude the possibility that other alterations within EpiNE2 and EpiNE8 (R15L and Y21S respectively) influence their affinity for HNE.

EpiNE7 was expressed as a soluble protein and analyzed for HNE inhibition activity by the fluorometric assay of Castillo et al. (CAST79); the data were analyzed by the method of Green and Work (GREE53). Preliminary results indicate that $K_d(\text{HNE}, \text{EpiNE7}) \leq 8 \cdot 10^{-12} \text{ M}$, i.e. at least 7.5-fold lower than the lowest K_d reported for a BPTI derivative with respect to HNE.

C. Summary

Taken together, these data show that the alterations which appear in the P1 region of the EPI mutants confer the ability to bind to HNE and hence be selected through the fractionation process. That the sequences of EpiNE1, 5 EpiNE3, and EpiNE7 appear frequently in the population of selected clones suggests that these clones display BPTI variants with the highest affinity for HNE of any of the 1000 potentially different variants in the MYMUT library. Furthermore, that pH conditions less than 4.0 are required 10 to elute these fusion phage from immobilized HNE suggests that they display BPTI variants having a higher affinity for HNE than BPTI(K15V,R17L). EpiNE7 exhibits a lower K_d toward HNE than does BPTI(K15V,R17L); EpiNE1 and EpiNE3 should also be expected to exhibit lower K_d s for HNE than 15 BPTI(K15V,R17L). It is possible that all of the listed EpiNEs have lower K_d s than BPRI(K15V,R17L).

Position 18 has not previously been identified as a key position in determining specificity or affinity of aprotinin homologues or derivatives for particular serine 20 proteases. None have reported or suggested that phenylalanine at position 18 will confer specificity and high affinity for HNE. One of the powerful advantages of the present invention is that many diverse amino-acid sequences may be tested simultaneously.

25

EXAMPLE V

SCREENING OF THE MYMUT LIBRARY FOR BINDING TO CATHEPSIN G BEADS.

We fractionated the MYMUT library over immobilized 5 human Cathepsin G to find an engineered protease inhibitor having high affinity for Cathepsin G, hereafter designated as an EpiC. The details of phage binding, elution of bound phage with buffers of decreasing pH (pH profile), titrating of the phage contained in these fractions,

composition of the MYMUT library, and the preparation of cathepsin G (Cat G) beads are essentially the same as detailed in Example IV.

- A pH profile for the binding of two starting controls, BPTI-III MK and EpiNE1, are shown in Figure 10. BPTI-III MK phage, which contains wild type BPTI fused to the III gene product, shows no apparent binding to Cat G beads in this assay. EpiNE1 phage was obtained by enrichment with HNE beads (Example IV and Table 208).
- 10 EpiNE1-III MK demonstrated little binding to Cat G beads in the assay, although a small peak or shoulder is visible in the pH 5 eluted fraction.

- Figure 11 shows the pH profiles of the MYMUT library phage when bound to Cat G beads. Library-Cat G interaction was monitored using three cycles of binding, pH elution, transduction of the pH 2 eluted phage, growth of the transduced phage and rebinding of any selected phage to Cat G beads, in an exact copy of that used to find variants of BPTI which bound to HNE. In contrast to the
- 15 pH profiles elicited with HNE beads, little enhancement of binding was observed for the same phage library when cycled with Cat G beads (with the exception of a possible 'shoulder' developing in the pH5 elutions).

- To investigate the elution profile around the pH 5 point in more detail, the binding of phage taken from the
- 25 pH 4 eluted fraction (bound to Cat G beads) rather than the previously used pH 2 fraction was examined. Figure 12 demonstrates a marked enhancement of phage binding to the Cat G beads with an apparent elution peak of pH 5. The
- 30 binding, as a fraction of the input phage population, increased with subsequent binding and elution cycles.

Individual phage clones were picked, grown and analyzed for binding to Cat G beads. Figure 13 shows the binding and pH profiles for the individual Cat G binding

clones (designated Epic variants). All clones exhibited minor peaks, superimposed upon a gradual fall in bound phage, at pH elutions of 5 (clones 1, 8, 10 and 11) or pH 4.5 (clone 7).

- 5 DNA sequencing of the Epic clones, shown in Table 209, demonstrated that the clones selected for binding to Cat G beads represented a distinct subset of the available sequences in the MYMUT library and a cluster of sequences different from that obtained when enriched with HNE beads.
- 10 The P1 residue in the Epic mutants is predominantly MET, with one example of PHE, while in BPTI it is LYS and in the EpINE variants it is either VAL or LEU. In the Epic mutants residue 16 is predominantly ALA with one example of GLY and residue 17 is PHE, ILE or LEU. Interestingly
- 15 residues 16 and 17 appear to pair off by complementary size, at least in this small sample. The small GLY residue pairs with the bulky PHE while the relatively larger ALA residue pairs with the less bulky LEU and ILE. The majority of the available residues in the MYMUT
- 20 library for positions 18 and 19 are represented in the Epic variants.

Hence, a distinct subset of related sequences from the MYMUT library have been selected for and demonstrated to bind to Cat G. A comparison of the pH profiles

25 elicited for the Epic variants with Cat G and the EpINE variants for HNE indicates that the EpINE variants have a high affinity for HNE while the Epic variants have a moderate affinity for Cat G. Nonetheless, the starting molecule, BPTI, has virtually no detectable affinity for

30 Cat G and the selection of clones with a moderate affinity is a significant finding.

EXAMPLE VI

SECOND ROUND OF VARIEGATION OF EpINE7 TO ENHANCE BINDING

TO HNE

A. MUTAGENESIS OF EpINE7 PROTEIN IN THE LOOP COMPRISING
RESIDUES 34-41

In Example IV, we described engineered protease
5 inhibitors EpINE1 through EpINE8 that were obtained by
affinity selection. Modeling of the structure of the
BPTI-Trypsin complex (Brookhaven Protein Data Bank entry
1TPA) indicates that the EpINE protein surface that
interacts with HNE is formed not only by residues 15-19
10 but also by residues 34-40 that are brought close to this
primary loop when the protein folds (HUBE74, HUBE75,
OAST88). Acting upon this assumption, we changed amino
acid residues in a second loop of the EpINE7 protein to
find EpINE7 derivatives having higher affinity for HNE.

15 In the complex of BPTI and trypsin found in Brook-
haven Protein Data Bank entry 1TPA ("1TPA complex"),
VAL₃₄ contacts TYR₁₅₁ and GLN₁₉₂. (Residues in trypsin or
HNE are underscored to distinguish them from the inhib-
itor.) In HNE, the corresponding residues are ILE₁₅₁ and
20 PHE₁₉₂. ILE is smaller and more hydrophobic than TYR.
PHE is larger and more hydrophobic than GLN. Neither of
the HNE side groups have the possibility to form hydrogen
bonds. When side groups larger than that of VAL are
substituted at position 34, interactions with residues
25 other than 151 and 192 may be possible. In particular, an
acidic residue at 34 might interact with ARG₁₄₇ of HNE
that corresponds to SER₁₄₇ of trypsin in 1TPA. Table 15
shows that, in 59 homologues of BPTI, 13 different amino
acids have been seen at position 34. Thus we allow all
30 twenty amino acids at 34.

Position 36 is not highly varied; only GLY, SER, and
ARG have been observed with GLY by far the most prevalent.
In the 1TPA complex, GLY₃₆ contacts HIS₅₇ and GLN₁₉₂.

09806095-062001

HIS₅₇ is conserved and GLN₁₉₂ corresponds to PHE₁₉₂ of HNE. Adding a methyl group to GLY₃₆ could increase hydrophobic interactions with PHE₁₉₂ of HNE. GLY₃₆ is in a conformation that most amino acids can achieve: $\phi = -79^\circ$ and $\psi = -9^\circ$ (Deisenhoffer cited in CREI84, p.222.).

In the 1TPA complex, ARG₃₉ contacts SER₉₆, ASN₉₇, THR₉₈, LEU₉₉, GLN₁₇₅, and TRP₂₁₅. In HNE, all of the corresponding residues are different! SER₉₆ is deleted; ASN₉₇ corresponds to ASP₉₇ (bearing a negative charge); THR₉₈ corresponds to PRO₉₈; LEU₉₉ corresponds to the residues VAL₉₉, ASN_{99a}, and LEU_{99b}; GLN₁₇₅ is deleted; and TRP₂₁₅ corresponds to PHE₂₁₅. Position 39 shows a moderately high degree of variability with 7 different amino acids observed, *viz.* ARG, GLY, LYS, GLN, ASP, PRO, and MET. Having seen PRO (the most rigid amino acid), GLY (the most flexible amino acid), LYS and ASP (basic and acidic amino acids), we assume that all amino acids are structurally compatible with the aprotinin backbone. Because the context of residue 39 has changed so much, we allow all 20 amino acids.

Position 40 is not highly variable; only GLY and ALA have been observed (with similar frequency, 24:16). Position 41 is moderately varied, showing ASN, LYS, ASP, GLN, HIS, GLU, and TYR. The side groups of residues 40 and 41 are not thought to contact trypsin in the 1TPA complex. Nevertheless, these residues can exert electrostatic effects and can influence the dynamic properties of residues 39, 38, and others. The choice of residues 34, 36, 39, 40, and 41 to be varied simultaneously illustrates the rule that the varied residues should be able to touch one molecule of the target material at one time or be able to influence residues that touch the target. These residues are not contiguous in sequence, nor are they contiguous on the surface of EpINE7. They can, nonethe-

less, all influence the contacts between the EpINE and HNE.

Amino acid residues VAL₃₄, GLY₃₆, MET₃₉, GLY₄₀, and ASN₄₁ were variegated as follows: any of 20 genetically encodable amino acids at positions 34 and 39 (NNS codons in which N is approximately equimolar A,C,T,G and S is approximately equimolar C and G), GLY or ALA at position 36 and 40 (GST codon), and [ASP, GLU, HIS, LYS, ASN, GLN, TYR, or stop] at position 41 (NAS codon). Because the PEPIs are displayed fused to gIII protein, DNA containing stop codons will not give rise to infectious phage in non-suppressor hosts.

For cassette mutagenesis, a 61 base long oligonucleotide DNA population was synthesized that contained 32,768 different DNA sequences coding on expression for a total of 11,200 amino acid sequences. This oligonucleotide extends from the third base of codon 51 in Table 113 (the middle of the StuI site) to base 2 of codon 70 (the EagI site (identified as XmaIII in Table 113)).

20 We used a mutagenesis method similar to that described by Cwirla et al. (CWIR90) and other standard DNA manipulations -described in Maniatis et al. (MANI82) and Sambrook et al. (SAMB89). EpiNE7 RF DNA was restricted with EagI and StuI, agarose gel purified, and dephosphorylated using HK^(TM) phosphatase (Epicentre Technologies). We prepared insert by annealing two small, 16 base and 17 base, phosphorylated synthetic DNA primers to the phosphorylated 61 base long oligonucleotide population described above. The resulting insert DNA population had the following features: double stranded DNA ends capable of regenerating upon ligation the EagI (5' overhang) and StuI (blunt) restricted sites of the EpiNE7 RF DNA, and single stranded DNA in the central mutagenic region. Insert and EpiNE7 vector DNA were ligated. Ligation

samples were used to transfect competent XL1-Blue^(TM) cells which were subsequently plated for formation of ampicillin resistant (Ap^R) colonies. The resulting phage-producing, Ap^R colonies were harvested and recombinant
 5 phage was isolated. By following these procedures, a phage library of $1.2 \cdot 10^5$ independent transformants was assembled. We estimated that 97.4% of the approximately $3.3 \cdot 10^4$ possible DNA sequences were represented:

$$0.974 = (1 - \exp\{-1.2 \cdot 10^5 / 32768\})$$

10 The probability of observing the parental sequence is higher than .974 because VAL occurs twice in the NNS codon:

Probability of seeing (V₃₄, G₃₆, M₃₉, G₄₀, N₄₁) =

$$(1 - \exp\{- (1.2 \cdot 10^5 \times 2 / 32768)\})$$

15 = $(1 - \exp\{-7.32\})$

$$= (1 - 6.5 \cdot 10^{-4})$$

$$= 0.99934$$

Furthermore, we expect that a small amount (for example, 1 part in 1000) of uncut or once-cut and religated parental
 20 vector would come through the procedures used. Thus the parental sequence is almost certainly present in the library. This library is designated the KLMUT library.

B. AFFINITY SELECTION WITH IMMOBILIZED HUMAN NEUTROPHIL ELASTASE

25 1) First Fractionation

We added $1.1 \cdot 10^8$ plaque forming units of the KLMUT library to 10 μ l of a 50% slurry of agarose-immobilized human neutrophil elastase beads (HNE from Calbiochem cross-linked to Reacti-Gel^(TM) agarose beads from Pierce
 30 Chemical Co. following manufacturer's directions) in TBS/BSA. Following 3 hours incubation at room temperature, the beads were washed and phage was eluted as done in the selection of EpiNE phage isolates (Example IV). The progression in lowering pH during the elution was: pH

7.0, 6.0, 5.0, 4.5, 4.0, 3.5, 3.0, 2.5, and 2.0. Beads carrying phage remaining after pH 2.0 elution were used to infect XL1-Blue^(TM) cells that were plated to allow plaque formation. The 348 resulting plaques were pooled to form
 5 a phage population for further affinity selection. A population of phage particles containing $6.0 \cdot 10^8$ plaque forming units was added to 10 μ l of a 50% slurry of agarose-immobilized HNE beads in TBS/BSA and the above selection procedure was repeated.

10 Following this second round of affinity selection, a portion of the beads was mixed with XL1-Blue^(TM) cells and plated to allow plaque formation. Of the resulting plaques, 480 were pooled to form a phage population for a third affinity selection. We repeated the selection
 15 procedure described above using a population of phage particles containing $3.0 \cdot 10^9$ plaque forming units. Portions of the pH 2.0 eluate and of the beads were plated with XL1-Blue^(TM) cells to allow formation of plaques. Individual plaques were picked for preparation of RF DNA.
 20 From DNA sequencing, we determined the amino acid sequence in the mutated secondary loop of 15 EpiNE7-homolog clones. The sequences are given in Table 210 as EpiNE7.1 through EpiNE7.20. Three sequences were observed twice: EpiNE7.4 and EpiNE7.14; EpiNE7.8 and EpiNE7.9; and EpiNE7.10 and
 25 EpiNE7.20. EpiNE7.4 was eluted at pH 2 while EpiNE7.14 was obtained by culturing HNE beads that had been washed with pH 2 buffer. Similarly, EpiNE7.10 came from pH 2 elution but EpiNE7.20 came from beads. EpiNE7.8 and EpiNE7.9 both came from pH 2 elution. Interestingly,
 30 EpiNE7.8 is found in both the first and second fractionations (EpiNE7.31 (vide infra)).

2) Second Fractionation

The purpose of affinity fractionation is to reduce diversity on the basis of affinity for the target. The

0986095-062901
 106290-56096860

first enrichment step of the first fractionation reduced the population from $3 \cdot 10^4$ possible DNA sequences to no more than 348. This might be too severe and some of the loss of diversity might not be related to affinity. Thus we carried out a second fractionation of the entire KLMUT library seeking to reduce the diversity more gradually.

We added $2.0 \cdot 10^{11}$ plaque forming units of the KLMUT library to 10 μ l of a 50% slurry of agarose-immobilized HNE beads in TBS/BSA. Following 3 hours incubation at room temperature, phage were eluted as described above. We then transduced XL1-Blue^(TM) cells with portions of the pH 2.0 eluate and plated for Ap^R colonies.

The resulting phage-producing colonies were harvested to obtain amplified phage for further affinity selection. A population of these phage particles containing $2.0 \cdot 10^{10}$ plaque forming units was added to 10 μ l of a 50% slurry of agarose-immobilized HNE beads in TBS/BSA and incubated for 90 minutes at room temperature. Phage were eluted as described above and portions of the pH 2.0 eluate were used to transduce XL1-Blue^(TM) cells. We plated the transductants for Ap^R colonies and obtained amplified phage from the harvested colonies.

In a third round of affinity selection, a population of phage particles containing $3.0 \cdot 10^{10}$ plaque forming units was added to 20 μ l of 50% slurry of agarose-immobilized HNE beads and incubated for 2 hours at room temperature. We eluted the phage with the following pH washes: pH 7.0, 6.0, 5.0, 4.5, 4.0, 3.5, 3.25, 3.0, 2.75, 2.5, 2.25, and 2.0. After plating a portion of the pH 2.0 eluate fraction for plaque formation, we picked individual plaques for preparation of RF DNA. DNA sequencing yielded the amino acid sequence in the mutated secondary loop for 20 EpiNE7 homolog clones. These sequences, together with EpiNE7, are given in Table 210 as EpiNE7.21 through

EpINE7.40. The plaques observed when EpINEs are plated display a variety of sizes. EpINE7.21 through EpINE7.30 were picked with attention to plaque size: 7.21, 7.22, and 7.23 from small plaques, 7.24 through 7.30 from plaques of increasing size, with 7.30 coming from a large plaque. TRP occurs at position 39 in EpINE7.21, 7.22, 7.23, 7.25, and 7.30. Thus plaque size does not correlate with the appearance of TRP at 39. One sequence, EpINE7.31, from this fractionation is identical to sequences EpINE7.8 and EpINE7.9 obtained in the first fractionation. EpINE7.30, EpINE7.34, and EpINE7.35 are identical, indicating that the diversity of the library has been greatly reduced. It is believed that these sequences have an affinity for HNE that is at least comparable to that of EpINE7 and probably higher. Because the parental EpINE7 sequence did not recur, it is quite likely that some or all of the EpINE7-.nn derivatives have higher affinity for HNE than does EpINE7.

3) Conclusions

One can draw some conclusions. First, because some sequences have been isolated repeatedly, the fractionation is nearly complete. The diversity has been reduced from $\geq 10^4$ to a few tens of sequences.

Second, the parental sequence has not recurred. At 39, MET did not occur! At position 34 VAL occurred only once in 35 sequences. At 41, ASN occurred only 4 of 35 times. At 40, GLY occurred 17 of 35 times. At position 36, GLY occurred 34 of 35 times, indicating that ALA is undesirable here. EpINE7.24 and EpINE7.36 are most like EpINE7, having three of the varied residues identical to EpINE7.

Third, the results of the first and second fractionation are similar. In the second fractionation, the prevalence of TRP at position 39 is more marked (5/15 in

FO6290-56095850

fractionation #1, 14/20 in #2). It is possible that the first fractionation lost some high-affinity EPIs through under-sampling. Nevertheless, the first fractionation was clearly quite successful.

- 5 Fourth, there are strong preferences at positions 39 and 36 and lesser but significant preferences at positions 34 and 41 with little preference at 40.

Heretofore, no homologues of aprotinin have been reported having ALA at 36. In the selected EpiNE7.nn
 10 sequences, the preference for GLY over ALA at position 36 is 34:1. This preference is probably not due to differences in protein stability. The process of the present invention, as applied in the present example, does not select against proteins on the basis of stability so long
 15 as the protein does fold and function at the temperature used in the procedure. ALA is probably tolerated at position 36 well enough to allow those proteins having ALA₃₆ to fold and function; one example was found having ALA₃₆. It may be relevant that the sole sequence having
 20 ALA₃₆ also has GLY₃₄. The flexibility of GLY at 34 may allow the methyl of ALA at 36 to fit into HNE in a way that is not possible when other amino acids occupy position 34.

At position 39, all 20 amino acids were allowed, but
 25 only seven were seen. TRP is strongly preferred with 19 occurrences, HIS second with six occurrences, and LEU third with 5 occurrences. No homologues of aprotinin have been reported having either TRP or HIS at position 39 as are now disclosed. Although LEU is represented in the NNS
 30 codon thrice, TRP and HIS have but one codon each and their prevalence is surprising. We constructed a model having HNE (Brookhaven Protein Data Bank entry 1HNE) and EpiNE7.9 spatially related as in the 1TPA complex. (The α carbons of HNE of conserved internal residues were

FOUO230"56096860

that they might facilitate binding to HNE, it was not and is not possible to predict which combination of these amino acids will lead to high affinity for HNE. The mutagenic oligonucleotide MYMUT was synthesized by Genetic Design Inc. (Houston, Texas).

2) Construction of Library of Fusion Phage Displaying Potential Engineered Protease Inhibitors

The single-stranded mutagenic MYMUT DNA was converted to the double stranded form with compatible XhoI and StuI ends and dephosphorylated with HK^(TM) phosphatase as described above for the VAL1 oligonucleotide. BPTI(MGNG)-III MA Rf DNA was digested with XhoI and StuI for 3 hours at 37°C to ensure complete digestion. The 8.0 kb DNA fragment was purified by agarose gel electrophoresis and Ultrafree-MC unit filtration. One µl of the dephosphorylated MYMUT DNA (5 ng) was ligated to 50 ng of the 8.0 kb fragment derived from BPTI(MGNG)-III MA Rf DNA. Under these conditions, the 10:1 molar ratio of insert to vector was found to be optimal for the generation of transformants. Ligation samples were extracted with phenol, phenol/chloroform/IAA (25:24:1, v:v:v) and chloroform/IAA (24:1, v:v) and DNA was ethanol precipitated prior to electroporation. One µl of the recovered ligation DNA was added to 40 µl of electro-competent cells. Cells were shocked using a Bio-Rad Gene Pulser device as described above. Immediately following electroshock, 1.0 ml of SOC media was added to the cells which were allowed to recover at 37°C for 60 minutes with shaking. The electroporated cells were plated onto LB plates containing Ap to permit the formation of colonies.

To assess the efficiency of the cassette mutagenesis procedure, 39 transformants were picked at random and phage present in culture supernatants were applied to a Nytran membrane and probed using the Dot Blot Procedure.

Two Nytran membranes were prepared in this manner. The first filter was allowed to hybridize to the CYSB oligonucleotide which had previously been radiolabelled. The second membrane was allowed to hybridize to the PRP1 oligonucleotide which had also been radiolabelled. Filters were subjected to autoradiography following washing under high stringency conditions. Of the 39 phage samples applied to the membrane, all 39 hybridized to the CYSB probe. This indicated that there was fusion phage in the culture supernatants and that at least the DNA encoding residues 35-47 appeared to be present in the phage genomes. Only 11 of the 39 samples hybridized to the PRP1 oligonucleotide indicating that 28% of the transformants were probably the parental phage BPTI(MGNG)-III MA used to generate the library. The remaining 28 clones failed to hybridize to the PRP1 probe indicating that substantial alterations were introduced into the P1 region by cassette mutagenesis using the MYMUT oligonucleotide. Of these 28 samples, all were found to contain infectious phage indicating that mutagenesis did not result in frame shift mutations which would lead to the generation of defective gene III products and non-infectious phage. (These 28 PEPI-displaying phage constitute a mini-library, the fractionation of which is discussed below.) Hence the overall efficiency of mutagenesis was estimated to be 72% in those cases where ligation DNA was not subjected to ApaI digestion prior to electroporation.

Bacterial colonies were harvested by overlaying chilled LB plates containing Ap with 5 ml of ice cold LB broth and scraping off cells using a sterile glass rod. A total of 4899 transformants were harvested in this manner of which 3299 were obtained by electroporation of ligation samples which were not digested with ApaI. Hence we estimate that 72% of these transformants (i.e. 2375)

represent mutants of the parental BPTI(MGNG)-III MA phage derived by cassette mutagenesis of the P1 position. An additional 1600 transformants were obtained by electroporation of ligation samples which had been digested with ApaI. If we assume that all of these clones contain new sequences at the P1 position then the total number of mutants in the pool of 4899 transformants is estimated to be $2375 + 1600 = 3975$. The total number of potentially different DNA sequences in the MYMUT library is 1728. We calculate that the library should display about 90% of the potential engineered protease inhibitor sequences as follows:

$$\begin{aligned} N_{\text{displayed}} &= N_{\text{possible}} \cdot (1 - \exp(-\text{Libsize}/N(\text{DNA}))) \\ &= 1000 \cdot (1 - \exp(-3975/1728)) = 900 \\ \% \text{ of possible sequences displayed} &= 100 \cdot (900 \div 1000) \\ &= 90\% \end{aligned}$$

3) Fractionation of a Mini-Library of Fusion Phage

We studied the fractionation of the mini library of 28 PEPIs to establish the appropriate parameters for fractionation of the entire MYMUT PEPI library. We anticipated that fractionation could be easier when the library of fusion phage was much less diverse than the entire MYMUT library. Fewer cycles of fractionation might be required to affinity purify a fusion phage exhibiting a high affinity for HNE. Secondly, since the sequences of all the fusion phage in the mini-library can be determined, one can determine the probability of selecting a given fusion phage from the initial population.

Two ml of the culture supernatants of the 28 PEPIs described above were pooled. Fusion phage were recovered, resuspended in 300 mM NaCl, 100 mM Tris, pH 8.0, 1 mM EDTA and stored on ice for 15 minutes. Insoluble material was

removed by centrifugation for 3 minutes in a microfuge at 4°C. The supernatant fraction was collected and PEPI phage were precipitated with PEG-8000. The final phage pellet was resuspended in TBS/BSA. Aliquots of the recovered phage were titrated for plaque-forming units on a lawn of cells. The final stock solution consisted of 200 μ l of fusion phage at a concentration of $5.6 \cdot 10^{12}$ pfu/ml.

a) First Enrichment Cycle

Forty μ l of the above phage stock was added to 10 μ l of a 50% slurry of HNE beads in TBS/BSA. The sample was allowed to mix on a Labquake shaker for 1.5 hours. Five hundred μ l of TBS/BSA was added to the sample and after an additional 5 minutes of mixing, the HNE beads were collected by centrifugation. The supernatant fraction was removed and the beads were resuspended in 0.5 ml of TBS/0.5% Tween-20. Beads were washed for 5 minutes on the shaker and recovered by centrifugation as above. The supernatant fraction was removed and the beads were subjected to 4 additional washes with TBS/Tween-20 as described above to reduce non-specific binding of fusion phage to HNE beads. Beads were washed twice as above with 0.5 ml of 50 mM sodium citrate pH 7.0, 150 mM NaCl containing 1.0 mg/ml BSA. The supernatants from the two washes were pooled. Subsequently, the HNE beads were washed sequentially with a series of 50 mM sodium citrate, 150 mM NaCl, 1.0 mg/ml BSA buffers of pH 6.0, 5.0, 4.5, 4.0, 3.5, 3.0, 2.5 and 2.0. Two washes were performed at each pH and the supernatants were pooled and neutralized by the addition of 260 μ l of 1 M Tris, pH 8.0. Aliquots of each pH fraction were diluted in LB broth and titrated for plaque-forming units on a lawn of cells. The total amount of fusion phage (as judged by pfu) appearing in each pH wash fraction was determined.

Figure 7 illustrates that the largest percentage of

T06290"56096860

input phage which bound to the HNE beads was recovered in the pH 5.0 fraction. The elution peak exhibits a trailing edge on the low pH side suggesting that a small proportion of the total bound fusion phage might elute from the HNE beads at a pH < 5. BPTI(K15L)-III phage display a BPTI variant with a moderate affinity for HNE ($K_d = 2.9 \cdot 10^{-9}$ M) (BECK88b). Since BPTI(K15L)-III phage elute from HNE beads as a peak centered on pH 4.75 and the highest peak in the first passage of the mini-library over HNE beads is centered on pH 5.0, we infer that many members of the MYMUT PEPI mini-library display PEPIs having moderate to high affinity for HNE.

To enrich for fusion phage displaying the highest affinity for HNE, phage contained in the lowest pH fraction (pH 2.0) from the first enrichment cycle were amplified and subjected to a second round of fractionation. Amplification involved the Transduction Procedure described above. Fusion phage (2000 pfu) were incubated with 100 μ l of cells for 15 minutes at 37°C in 200 μ l of 1 X Minimal A salts. Two hundred μ l of 2 X LB broth was added to the sample and cells were allowed to recover for 15 minutes at 37°C with shaking. One hundred μ l portions of the above sample were plated onto LB plates containing Ap. Five such transduction reactions were performed yielding a total of 20 plates, each containing approximately 350 colonies (7000 transformants in total). Bacterial cells were harvested as described for the preparation of the MYMUT library and fusion phage were collected as described for the preparation of the mini-library. A total of 200 μ l of fusion phage ($4.3 \cdot 10^{12}$ pfu/ml in TBS/BSA) derived from the pH 2.0 fraction from the first passage of the mini-library was obtained in this manner.

b) Second Enrichment Cycle

Forty μ l of the above phage stock was added to 10 μ l of a 50% slurry of HNE beads in TBS/BSA. The sample was allowed to mix for 1.5 hours and the HNE beads were washed with TBS/BSA, TBS/0.5% Tween and sodium citrate buffers as described above. Aliquots of neutralized pH fractions were diluted and titered as described above.

The elution profile for the second passage of the mini-library over HNE beads is shown in Figure 7. The largest percentage of the input phage which bound to the HNE beads was recovered in the pH 3.5 wash. A smaller peak centered on pH 4.5 may represent residual fusion phage from the first passage of the mini-library which eluted at pH 5.0. The percentage of total input phage which eluted at pH 3.5 in the second cycle exceeds the percentage of input phage which eluted at pH 5.0 in the first cycle. This is indicative of more avid binding of fusion phage to the HNE matrix. Taken together, the significant shift in the pH elution profile suggests that selection for fusion phage displaying BPTI variants with higher affinity for HNE occurred.

c) Third Cycle

Phage obtained in the pH 2.0 fraction from the second passage of the mini-library were amplified as above and subjected to a third round of fractionation. The pH elution profile is shown in Figure 7. The largest percentage of input phage was recovered in the pH 3.5 wash as is the case with the second passage of the mini-library. However, the minor peak centered on pH 4.5 is diminished in the third passage relative to the second passage. Furthermore, the percentage of input phage which eluted at pH 3.5 is greater in the third passage than in the second passage. In comparison, the BPTI(K15V,R17L)-III fusion phage elute from HNE beads as a peak centered on pH 4.25. Taken together, the data suggests that a

5

10

15

20

25

22

three passes of the mini-library over HNE beads, 1 out of 28 input fusion phage was selected for and appears as a pure population in the lowest pH fraction from the third passage of the library. That the EpiNE α phage elute at pH 3.5 while BPTI(K15V,R17L)-III MA phage elute at a higher pH strongly suggests that the EpiNE α protein has a significantly higher affinity than BPTI(K15V,R17L) for HNE.

4) Fractionation of the MYMUT Library

a) Three cycles of enrichment

The same procedure used above to fractionate the mini-library was used to fractionate the entire MYMUT PEPI library consisting of fusion phage displaying 1000 different proteins. The phage inputs for the first, second and third rounds of fractionation were $4.0 \cdot 10^{11}$, $5.8 \cdot 10^{10}$, and $1.1 \cdot 10^{11}$ pfu respectively. Figure 8 illustrates that the largest percentage of input phage which bound to the HNE matrix was recovered in the pH 5.0 wash in the first enrichment cycle. The pH elution profile is very similar to that seen for the first passage of the mini-library over HNE beads. A trailing edge is also observed on the low pH side of the pH 5.0 peak however this is not as prominent as that observed for the mini-library. The percentage of input phage which eluted in the pH 7.0 wash was greater than that eluted in the pH 6.0 wash. This is in contrast to the result obtained for the first passage of the mini library and may reflect the presence of $\approx 20\%$ parental BPTI(MGNG)-III MA phage in the MYMUT library pool. These phage adhere to the HNE beads weakly (if at all) and elute in the pH 7.0 fraction. That no parent phage were present in the mini-library is consistent with the absence of a peak at pH 7.0 in the first passage of the mini-library.

Phage present in the pH 2.0 fraction from the first passage of the MYMUT library were amplified as described

T06290"56056350

previously and subjected to a second round of fractionation. The largest percentage of input phage which bound to the HNE beads was recovered in the pH 3.5 wash (Figure 8). A minor peak centered on pH 4.5 was also evident.

5 The fact that more extreme pH conditions were required to elute the majority of bound fusion phage suggested that selection of fusion phage displaying PEPs with higher affinity for HNE had occurred. This was also indicated by the fact that the total percentage of input phage which

10 appeared in the pH 3.5 wash in the second enrichment cycle was 10 times greater than the percentage of input which appeared in the pH 5.0 wash in the first cycle.

Fusion phage from the pH 2.0 fraction of the second pass of the MYMUT library were amplified and subjected to

15 a third passage over HNE beads. The proportion of fusion phage appearing in the pH 3.5 fraction relative to that in the 4.5 fraction was greater in the third passage than in the second passage (Figure 8). Also the amount of fusion phage appearing in the pH 3.5 fraction was higher in the

20 third passage than in the second passage. The fact that wash conditions less than pH 4.25 were required to elute bound fusion phage derived from the MYMUT library suggests that the EpiNEs displayed by these phage possess a higher affinity for HNE than the BPTI(K15V,R17L) variant.

25 b) Characterization of Selected Clones

The pH 2.0 fraction from the third enrichment cycle of the MYMUT library was titered on a lawn of cells. Twenty plaques were picked at random. Rf DNA was prepared for each of the clones and fusion phage were collected by

30 PEG precipitation. Clonally pure populations of fusion phage in TBS/BSA were prepared and characterized with respect to their affinity for immobilized HNE. pH elution profiles were obtained to determine the stringency of the conditions required to elute bound fusion phage from the

HNE matrix. Figure 9 illustrates the pH profiles obtained for EpiNE clones 1, 3, and 7. The pH profiles for all 3 clones exhibit a peak centered on pH 3.5. Unlike the pH profile obtained for the third passage of the MYMUT library, no minor peak centered on pH 4.5 is evident. This is consistent with the clonal purity of the selected EpiNE phage utilized to generate the profiles. The elution peaks are not symmetrical and a prominent trailing edge on the low pH side. In all probability, the 10 minute elution period employed is inadequate to remove bound fusion phage at the low pH conditions. EpiNE clones 1 through 8 have the following characteristics: five clones (identified as EpiNE1, EpiNE3, EpiNE5, EpiNE6, and EpiNE7) display very similar pH profiles centered on pH 3.5. The remaining 3 clones elute in the pH 3.5 to 4.0 range. There remains some diversity amongst the 20 randomly chosen clones obtained from the pH 2.0 fraction of the third passage of the MYMUT library and these clones might exhibit different affinities for HNE.

20 c) Sequences of the EpiNE Clones

The DNA sequences encoding the P1 regions of the different EpiNE clones were determined by dideoxy sequencing of Rf DNA. The sequences are shown in Table 208. Essentially, only the codons targeted for mutagenesis (i.e. 15 to 19) were altered as a consequence of cassette mutagenesis using the MYMUT oligonucleotide. Only 1 codon outside the target region was found to contain an unexpected alteration. In this case, codon 21 of EpiNE8 was altered from a tyrosine codon (TAT) to a SER codon (TCT) by a single nucleotide substitution. This error could have been introduced into the MYMUT oligonucleotide during its synthesis. Alternatively, an error could have been introduced when the single-stranded MYMUT oligonucleotide was converted to the double-stranded form by Sequenase.

7) Effect of pH on the Dissociation of Bound BPTI-III MK and BPTI(K15L)-III MA Phage from Immobilized Neutrophil Elastase

The affinity of a given fusion phage for an immobilized serine protease can be characterized on the basis of the amount of bound fusion phage which elutes from the beads by washing with a pH 2.2 buffer. This represents rather extreme conditions for the dissociation of fusion phage from beads. Since the affinity of the BPTI variants described above for HNE is not high ($K_d > 1 \cdot 10^{-9}$ M) it was anticipated that fusion phage displaying these variants might dissociate from HNE beads under less severe pH conditions. Furthermore fusion phage might dissociate from HNE beads under specific pH conditions characteristic of the particular BPTI variant displayed by the phage. Low pH buffers providing stringent wash conditions might be required to dissociate fusion phage displaying a BPTI variant with a high affinity for HNE whereas neutral pH conditions might be sufficient to dislodge a fusion phage displaying a BPTI variant with a weak affinity for HNE.

Thirty μ l of BPTI(K15L)-III MA phage ($1.7 \cdot 10^{10}$ pfu/ml in TBS/BSA) were added to 5 μ l of a 50% slurry of immobilized HNE also in TBS/BSA. Similarly, 30 μ l of BPTI-III MA phage ($8.6 \cdot 10^{10}$ pfu/ml in TBS/BSA) were added to 5 μ l of
25 immobilized HNE. The above conditions were chosen to ensure that an approximately equivalent number of phage particles were added to the beads. The samples were incubated for 3 hours on a Labquake shaker. The beads were washed with 0.5 ml of TBS/BSA for 5 min on the
30 shaker, recovered by centrifugation and the supernatant was removed. The beads were washed with 0.5 ml of TBS/0.1% Tween-20 for 5 minutes and recovered by centrifugation. Four additional washes with TBS/0.1% Tween-20 were performed as described above. The beads were washed

as above with 0.5 ml of 100 mM sodium citrate, pH 7.0 containing 1.0 mg/ml BSA. The beads were recovered by centrifugation and the supernatant was removed. Subsequently, the HNE beads were washed sequentially with a series of 100 mM sodium citrate, 1.0 mg/ml BSA buffers of pH 6.0, 5.0, 4.0 and 3.0 and finally with the 2.2 elution buffer described above. The pH washes were neutralized by the addition of 1 M Tris, pH 8.0, diluted in LB broth and titered for plaque-forming units on a lawn of cells.

Table 203 illustrates that a low percentage of the input BPTI-III MK fusion phage adhered to the HNE beads and was recovered in the pH 7.0 and 6.0 washes predominantly. By contrast, a significantly higher percentage of the BPTI(K15L)-III MA phage bound to the HNE beads and was recovered predominantly in the pH 5.0 and 4.0 washes. Hence lower pH conditions (*i.e.* more stringent) are required to dissociate BPTI(K15L)-III MA than BPTI-MK phage from immobilized HNE. The affinity of BPTI(K15L) is over 1000 times greater than that of BPTI for HNE (based on reported K_d values (BECK88b)). Hence this suggests that lower pH conditions are indeed required to dissociate fusion phage displaying a BPTI variant with a higher affinity for HNE.

8) Construction of BPTI(MGNG)-III MA Phage

The light chain of bovine inter- α -trypsin inhibitor contains 2 domains highly homologous to BPTI. The amino terminal proximal domain (called BI-8e) has been generated by proteolysis and shown to be a potent inhibitor of HNE ($K_d = 4.4 \cdot 10^{-11}$ M) (ALBR83). By contrast a BPTI variant with the single substitution of LEU for LYS₁₅ exhibits a moderate affinity for HNE ($K_d = 2.9 \cdot 10^{-9}$ M) (BECK88b). It has been proposed that the P1 residue is the primary determinant of the specificity and potency of BPTI-like molecules (BECK88b, LASK80 and works cited therein).

Although both BI-8e and BPTI(K15L) feature LEU at their respective P1 positions, there is a 66 fold difference in the affinities of these molecules for HNE. Structural features, other than the P1 residue, must contribute to the affinity of BPTI-like molecules for HNE.

A comparison of the structures of BI-8e and BPTI-(K15L) reveals the presence of three positively charged residues at positions 39, 41, and 42 of BPTI which are absent in BI-8e. These hydrophilic and highly charged residues of BPTI are displayed on a loop which underlies the loop containing the P1 residue and is connected to it via a disulfide bridge. Residues within the underlying loop (in particular residue 39) participate in the interaction of BPTI with the surface of trypsin near the catalytic pocket (BLOW72) and may contribute significantly to the tenacious binding of BPTI to trypsin. However, these hydrophilic residues might hamper the docking of BPTI variants with HNE. In support of this hypothesis, BI-8e displays a high affinity for HNE and contains no charged residues in the region spanning residues 39-42. Hence residues 39 through 42 of wild type BPTI were replaced with the corresponding residues of the human homologue of BI-8e. We anticipated that a BPTI derivative containing the MET-GLY-ASN-GLY (MGNG) sequence would exhibit a higher affinity for HNE than corresponding derivatives which retain the sequence of wild type BPTI at residues 39-42.

A double stranded oligonucleotide with AccI and EagI compatible ends was designed to introduce the desired alteration of residues 39 to 42 via cassette mutagenesis. Codon 45 was altered to create a new XmnI site, unique in the structure of the BPTI gene, which could be used to screen for mutants. This alteration at codon 45 does not alter the encoded amino-acid sequence. BPTI-III MA Rf DNA

was digested with AccI. Two oligonucleotides (CYSB and CYST) corresponding to the bottom and top strands of the mutagenic DNA were annealed and ligated to the AccI digested BPTI-III MA Rf DNA. The sample was digested with

5 BglII and the 2.1 kb BglII/EagI fragment was purified. BPTI-III MA Rf was also digested with BglII and EagI and the 6.0 kb fragment was isolated and ligated to the 2.1 kb BglII/EagI fragment described above. Ligation samples were used to transfect competent cells which were plated

10 to permit the formation of plaques on a lawn of cells. Phage derived from plaques were probed with a radioactively labelled oligonucleotide (CYSB) using the Dot Blot Procedure. Positive clones were identified by autoradiography of the Nytran membrane after washing at high

15 stringency conditions. Rf DNA was prepared from Ap^R cultures containing fusion phage which hybridized to the CYSB probe. Restriction enzyme analysis and DNA sequencing confirmed that codons 39-42 of BPTI had been altered. The Rf DNA was designated BPTI(MGNG)-III MA.

20 9) Construction of BPTI(K15L, MGNG)-III MA

BPTI(MGNG)-III MA Rf DNA was digested with AccI and the 5.6 kb fragment was purified. BPTI(K15L)-III MA was digested with AccI and the 2.5 kb DNA fragment was purified. The two fragments above were ligated together

25 and ligation samples were used to transfect competent cells which were plated for plaque production. Large and small plaques were observed on the plate. Representative plaques of each type were picked and phage were probed with the LEU1 oligonucleotide via the Dot Blot Procedure.

30 After the Nytran filter had been washed under high stringency conditions, positive clones were identified by autoradiography. Only the phage which hybridized to the LEU1 oligonucleotide gave rise to the small plaques confirming an earlier observation that substitution of LEU

T05290 569560

for LYS₁₅ substantially reduces phage infectivity. Appropriate cultures containing phage which hybridized to the LEU1 oligonucleotide were used to prepare Rf DNA. Restriction enzyme analysis and DNA sequencing confirmed
 5 that the K15L mutation had been introduced into BPTI-(MGNG)-III MA. This Rf DNA was designated BPTI(K15L,-MGNG)-III MA.

10) Effect of Mutation of Residues 39-42 of BPTI(K15L) on its Affinity for Immobilized HNE

10 Thirty μ l of BPTI(K15L,MGNG)-III MA phage ($9.2 \cdot 10^9$ pfu/ml in TBS/BSA) were added to 5 μ l of a 50% slurry of immobilized HNE also in TBS/BSA. Similarly 30 μ l of BPTI(K15L)-III MA phage ($1.2 \cdot 10^{10}$ pfu/ml in TBS/BSA) were added to immobilized HNE. The samples were incubated for
 15 3 hours on a Labquake shaker. The beads were washed for 5 min with 0.5 ml of TBS/BSA and recovered by centrifugation. The beads were washed 5 times with 0.5 ml of TBS/0.1% Tween-20 as described above. Finally, the beads were washed sequentially with a series of 100 mM sodium
 20 citrate buffers of pH 7.0, 6.0, 5.5, 5.0, 4.75, 4.5, 4.25, 4.0 and 3.5 as described above. pH washes were neutralized, diluted in LB broth and titered for plaque-forming units on a lawn of cells.

Table 204 illustrates that almost twice as much of
 25 the BPTI(K15L,MGNG)-III MA as BPTI(K15L)-III MA phage bound to HNE beads. In both cases the pH 4.75 fraction contained the largest proportion of the recovered phage. This confirms that replacement of residues 39-42 of wild type BPTI with the corresponding residues of BI-8e
 30 enhances the binding of the BPTI(K15L) variant to HNE.

11) Fractionation of a Mixture of BPTI-III MK and BPTI(K15L,MGNG)-III MA Fusion Phage

The observations described above indicate that

0096095-062901

BPTI(K15L,MGNG)-III MA and BPTI-III MK phage exhibit different pH elution profiles from immobilized HNE. It seemed plausible that this property could be exploited to fractionate a mixture of different fusion phage.

- 5 Fifteen μ l of BPTI-III MK phage ($3.92 \cdot 10^{10}$ pfu/ml in TBS/BSA), equivalent to $8.91 \cdot 10^7$ Km^R transducing units, were added to 15 μ l of BPTI(K15L,MGNG)-III MA phage ($9.85 \cdot 10^9$ pfu/ml in TBS/BSA), equivalent to $4.44 \cdot 10^7$ Ap^R transducing units. Five μ l of a 50% slurry of immobilized
- 10 HNE in TBS/BSA was added to the phage and the sample was incubated for 3 hours on a Labquake mixer. The beads were washed for 5 minutes with 0.5 ml of TBS/BSA prior to being washed 5 times with 0.5 ml of TBS/2.0% Tween-20 as described above. Beads were washed for 5 minutes with 0.5
- 15 ml of 100 mM sodium citrate, pH 7.0 containing 1.0 mg/ml BSA. The beads were recovered by centrifugation and the supernatant was removed. Subsequently, the HNE beads were washed sequentially with a series of 100 mM citrate buffers of pH 6.0, 5.0 and 4.0. The pH washes were
- 20 neutralized by the addition of 130 μ l of 1 M Tris, pH 8.0.

- The relative proportion of BPTI-III MK and BPTI(K15L-,MGNG)-III MA phage in each pH fraction was evaluated by determining the number of phage able to transduce cells to Km^R as opposed to Ap^R. Fusion phage diluted in 1 X
- 25 Minimal A salts were added to 100 μ l of cells (O.D.600 = 0.8 concentrated to 1/20 original culture volume) also in Minimal salts in a final volume of 200 μ l. The sample was incubated for 15 min at 37°C prior to the addition of 200 μ l of 2 X LB broth. After an additional 15 min incubation
 - 30 at 37°C, duplicate aliquots of cells were plated on LB plates containing either Ap or Km to permit the formation of colonies. Bacterial colonies on each type of plate were counted and the data was used to calculate the number of Ap^R and Km^R transducing units in each pH fraction. The

FO6290 56096860

number of Ap^R transducing units is indicative of the amount of BPTI(K15L, MGNG)-III MA phage in each pH fraction while the total number of Km^R transducing units is indicative of the amount of BPTI-III MK phage.

- 5 Table 205 illustrates that a low percentage of the BPTI-III MK input phage (as judged by Km^R transducing units) adhered to the HNE beads and was recovered predominantly in the pH 7.0 fraction. By contrast, a significantly higher percentage of the BPTI(K15L, MGNG)-III MA
- 10 phage (as judged by Ap^R transducing units) adhered to the HNE beads and was recovered predominantly in the pH 4.0 fraction. A comparison of the total number of Ap^R and Km^R transducing units in the pH 4.0 fraction shows that a 984-fold enrichment of BPTI(K15L, MGNG)-III MA phage over BPTI-
- 15 III MK phage was achieved. Hence, the above procedure can be utilized to fractionate mixtures of fusion phage on the basis of their relative affinities for immobilized HNE.

12) Construction of BPTI(K15V, R17L)-III MA

- A BPTI variant containing the alterations K15V and
- 20 R17L demonstrates the highest affinity for HNE of any BPTI variant described to date ($K_d = 6 \cdot 10^{-11}$ M) (AUER89). As a means of testing the selection system described herein, a fusion phage displaying this variant of BPTI was generated and used as a "reference" phage to characterize the
- 25 affinity for immobilized HNE of fusion phage displaying a BPTI variant with a known affinity for free HNE. A 76 bp mutagenic oligonucleotide (VAL1) was designed to convert the LYS₁₅ codon (AAA) to a VAL codon (GTT) and the ARG₁₇ codon (CGA) to a LEU codon (CTG). At the same time codons
- 30 11, 12 and 13 were altered to destroy the ApaI site resident in the wild type BPTI gene while creating a new RsrII site, which could be used to screen for correct clones.

The single stranded VAL1 oligonucleotide was convert-

0956095 062901

ed to the double stranded form following the procedure described in Current Protocols in Molecular Biology (AUSU87). One μg of the VAL1 oligonucleotide was annealed to one μg of a 20 bp primer (MB8). The sample was heated to 80°C, cooled to 62°C and incubated at this temperature for 30 minutes before being allowed to cool to 37°C. Two μl of a 2.5 mM mixture of dNTPs and 10 units of Sequenase (U.S.B., Cleveland, Ohio) were added to the sample and second strand synthesis was allowed to proceed for 45 minutes at 37°C. One hundred units of XhoI was added to the sample and digestion was allowed to proceed for 2 hours at 37°C in 100 μl of 1 X XhoI digestion buffer. The digested DNA was subjected to electrophoreses on a 4% GTG NuSieve agarose (FMC Bioproducts, Rockland, ME) gel and the 65 bp fragment was excised and purified from melted agarose by phenol extraction and ethanol precipitation. A portion of the recovered 65 bp fragment was subjected to electrophoresis on a 4% GTG NuSieve agarose gel for quantitation. One hundred nanograms of the recovered fragment was dephosphorylated with 1.9 μl of HK^(TM) phosphatase (Epicentre Technologies, Madison, WI) at 37°C for 60 minutes. The reaction was stopped by heating at 65°C for 15 minutes. BPTI-MA Rf DNA was digested with XhoI and StuI and the 8.0 kb fragment was isolated. One μl of the dephosphorylation reaction (5 ng of double-stranded VAL1 oligonucleotide) was ligated to 50 ng of the 8.0 kb XhoI/StuI fragment derived from BPTI-III MA Rf. Ligation samples were subjected to phenol extraction and DNA was recovered by ethanol precipitation. Portions of the recovered ligation DNA were added to 40 μl of electro-competent cells which were shocked using a Bio-Rad Gene Pulser device set at 1.7 kv, 25 μF and 800 Ω . One ml of SOC media was immediately added to the cells which were allowed to recover at 37°C for one hour. Aliquots of the electroporated cells were plated onto LB plates containing

T06290" 56096860

Ap to permit the formation of colonies.

Phage contained within cultures derived from picked Ap^R colonies were probed with two radiolabelled oligonucleotides (PRP1 and ESP1) via the Dot Blot Procedure. Rf DNA was prepared from cultures containing phage which exhibited a strong hybridization signal with the ESP1 oligonucleotide but not with the PRP1 oligonucleotide. Restriction enzyme analysis verified loss of the ApaI site and acquisition of a new RsrII site diagnostic for the changes in the P1 region. Fusion phage were also probed with a radiolabelled oligonucleotide (VLP1) via the Dot Blot Procedure. Autoradiography confirmed that fusion phage which previously failed to hybridize to the PRP1 probe, hybridized to the VLP1 probe. DNA sequencing confirmed that the LYS₁₅ and ARG₁₇ codons had been converted to VAL and LEU codons respectively. The Rf DNA was designated BPTI(K15V,R17L)-III MA.

13) Affinity of BPTI(K15V,R17L)-III MA Phage for Immobilized HNE

Forty μ l of BPTI(K15,R17L)-III MA phage ($9.8 \cdot 10^{10}$ pfu/ml) in TBS/BSA were added to 10 μ l of a 50% slurry of immobilized HNE also in TBS/BSA. Similarly, 40 μ l of BPTI(K15L,MGNG)-III MA phage ($5.13 \cdot 10^9$ pfu/ml) in TBS/BSA were added to immobilized HNE. The samples were mixed for 1.5 hours on a Labquake shaker. Beads were washed once for 5 min with 0.5 ml of TBS/BSA and then 5 times with 0.5 ml of TBS/1.0% Tween-20 as described previously. Subsequently the beads were washed sequentially with a series of 50 mM sodium citrate buffers containing 150 mM NaCl, 1.0 mg/ml BSA of pH 7.0, 6.0, 5.0, 4.5, 4.0, 3.75, 3.5 and 3.0. In the case of the BPTI(K15L,MGNG)-III MA phage, the pH 3.75 and 3.0 washes were omitted. Two washes were performed at each pH and the supernatants were pooled, neutralized with 1 M Tris pH 8.0, diluted in LB broth and

titered for plaque-forming units on a lawn of cells.

Table 206 illustrates that the pH 4.5 and 4.0 fractions contained the largest proportion of the recovered BPTI(K15V,R17L)-III MA phage. By contrast, the BPTI(K15L,MCNG)-III MA phage, like BPTI(K15L)-III MA phage, were recovered predominantly in the pH 5.0 and 4.5 fractions, as shown above. The affinity of BPTI(K15V,-R17L) is 48 times greater than that of BPTI(K15L) for HNE (based on reported K_d values, AUER89 for BPTI(K15V,R17L) and BECK88b for BPTI(K15L)). That the pH elution profile for BPTI(K15V,R17L)-III MA phage exhibits a peak at pH 4.0 while the profile for BPTI(K15L)-III MA phage displays a peak at pH 4.5 supports the contention that lower pH conditions are required to dissociate, from immobilized HNE, fusion phage displaying a BPTI variant with a higher affinity for free HNE.

--- *** ---

EXAMPLE IV

CONSTRUCTION OF A VARIEGATED POPULATION OF PHAGE DISPLAY-
ING BPTI DERIVATES AND FRACTIONATION FOR MEMBERS THAT
DISPLAY BINDING DOMAINS HAVING HIGH AFFINITY FOR HUMAN
NEUTROPHIL ELASTASE:

We here describe generation of a library of 1000 different potential engineered protease inhibitors (PEPIs) and the fractionation with immobilized HNE to obtain an engineered protease inhibitor (Epi) having high affinity for HNE. Successful Epis that bind HNE are designated EpiNEs.

1) Design of a Mutagenic Oligonucleotide to Create a
Library of Fusion Phage

A 76 bp variegated oligonucleotide (MYMUT) was

designed to construct a library of fusion phage displaying 1000 different PEPIs derived from BPTI. The oligonucleotide contains 1728 different DNA sequences but due to the degeneracy of the genetic code, it encodes 1000 different protein sequences. The oligonucleotide was designed so as to destroy an ApaI site (shown in Table 113) encompassing codons 12 and 13. ApaI digestion could be used to select against the parental Rf DNA used to construct the library.

The MYMUT oligonucleotide permits the substitution of 5 hydrophobic residues (PHE, LEU, ILE, VAL, and MET via a DTS codon (D = approximately equimolar A, T, and G; S = approximately equimolar C and G)) for LYS₁₅. Replacement of LYS₁₅ in BPTI with aliphatic hydrophobic residues via semi-synthesis has provided proteins having higher affinity for HNE than BPTI (TANN77, JERI74a,b, WENZ80, TSCH86, BECK88b). At position 16, either GLY or ALA are permitted (GST codon). This is in keeping with the predominance of these two residues at the corresponding positions in a variety of BPTI homologues (CREI87). The variegation scheme at position 17 is identical to that at 15. Limited data is available on the relative contribution of this residue to the interaction of BPTI homologues with HNE. A variety of hydrophobic residues at position 17 was included with the anticipation that they would enhance the docking of a BPTI variant with HNE. Finally at positions 18 and 19, 4 (PHE, SER, THR, and ILE via a WYC codon (W = approximately equimolar A and T; Y = approximately equimolar T and C)) and 5 (SER, PRO, THR, LYS, GLN, and stop via an HMA codon (H = approximately equimolar A, C, and T; M = approximately equimolar A and C)) different amino acids respectively are encoded. These different amino acid residues are found in the corresponding positions of BPTI homologues that are known to bind to HNE (CREI87). Although the amino acids included in the PEPI library were chosen because there was some indication

DNA fragment from MK RF was radioactively labelled with ^{32}P - αdCTP using an oligolabelling kit (Pharmacia, Piscataway, NJ). Nytran membranes were transferred from pre-hybridization solution to Southern hybridization solution (5Prime-3Prime) at 42°C . The radioactive probe was added to the hybridization solution and following overnight incubation at 42°C , the filter was washed 3 times with $2 \times \text{SSC}$, 0.1% SDS at room temperature and once at 65°C in $2 \times \text{SSC}$, 0.1% SDS. Nytran membranes were subjected to autoradiography. The efficiency of the affinity selection system can be semi-quantitatively determined using the above dot blot procedure. Comparison of dots A1 and B1 or C1 and D1 indicates that the majority of phage did not stick to the streptavidin-agarose beads. Washing with TBS/Tween buffer removes the majority of phage which are non-specifically associated with streptavidin beads. Exposure of the streptavidin beads to elution buffer releases bound phage only in the case of MK-BPTI phage which have previously been incubated with biotinylated rabbit anti-BPTI IgG. This data indicates that the affinity selection system described above can be utilized to select for phage displaying a specific antigen (in this case BPTI). We estimate an enrichment factor of at least 40 fold based on the calculation

$$\text{Enrichment Factor} = \frac{\text{Percent MK-BPTI phage recovered}}{\text{Percent MK phage recovered}}$$

EXAMPLE III

CHARACTERIZATION AND FRACTIONATION OF CLONALLY PURE POPULATIONS OF PHAGE, EACH DISPLAYING A SINGLE CHIMERIC APROTININ HOMOLOGUE/M13 GENE III PROTEIN:

This Example demonstrates that chimeric phage

proteins displaying a target-binding domain can be eluted from immobilized target by decreasing pH, and the pH at which the protein is eluted is dependent on the binding affinity of the domain for the target.

5 Standard Procedures:

Unless otherwise noted, all manipulations were carried out at room temperature. Unless otherwise noted, all cells are XL1-Blue^(TM) (Stratagene, La Jolla, CA).

1) Demonstration of the Binding of BPTI-III MK Phage to

10 Active Trypsin Beads

Previous experiments designed to verify that BPTI displayed by fusion phage is functional relied on the use of immobilized anhydro-trypsin, a catalytically inactive form of trypsin. Although anhydro-trypsin is essentially
 15 identical to trypsin structurally (HUBE75, YOKO77) and in binding properties (VINC74, AKOH72), we demonstrated that BPTI-III fusion phage also bind immobilized active trypsin. Demonstration of the binding of fusion phage to immobilized active protease and subsequent recovery of
 20 infectious phage facilitates subsequent experiments where the preparation of inactive forms of serine proteases by protein modification is laborious or not feasible.

Fifty μ l of BPTI-III MK phage (identified as MK-BPTI
 25 in USSN 07/487,063) ($3.7 \cdot 10^{11}$ pfu/ml) in either 50 mM Tris, pH 7.5, 150 mM NaCl, 1.0 mg/ml BSA (TBS/BSA) buffer or 50 mM sodium citrate, pH 6.5, 150 mM NaCl, 1.0 mg/ml BSA (CBS/BSA) buffer were added to 10 μ l of a 25% slurry of immobilized trypsin (Pierce Chemical Co., Rockford, IL)
 30 also in TBS/BSA or CBS/BSA. As a control, 50 μ l MK phage ($9.3 \cdot 10^{12}$ pfu/ml) were added to 10 μ l of a 25% slurry of immobilized trypsin in either TBS/BSA or CBS/BSA buffer. The infectivity of BPTI-III MK phage is 25-fold lower than

that of MK phage; thus the conditions chosen above ensure that an approximately equivalent number of phage particles are added to the trypsin beads. After 3 hours of mixing on a Labquake shaker (Labindustries Inc., Berkeley, CA)

5 0.5 ml of either TBS/BSA or CBS/BSA was added where appropriate to the samples. Beads were washed for 5 min and recovered by centrifugation for 30 sec. The supernatant was removed and 0.5 ml of TBS/0.1% Tween-20 was added. The beads were mixed for 5 minutes on the shaker

10 and recovered by centrifugation as above. The supernatant was removed and the beads were washed an additional five times with TBS/0.1% Tween-20 as described above. Finally, the beads were resuspended in 0.5 ml of elution buffer (0.1 M HCl containing 1.0 mg/ml BSA adjusted to pH 2.2

15 with glycine), mixed for 5 minutes and recovered by centrifugation. The supernatant fraction was removed and neutralized by the addition of 130 μ l of 1 M Tris, pH 8.0. Aliquots of the neutralized elution sample were diluted in LB broth and titered for plaque-forming units on a lawn of

20 cells.

Table 201 illustrates that a significant percentage of the input BPTI-III MK phage bound to immobilized trypsin and was recovered by washing with elution buffer. The amount of fusion phage which bound to the beads was

25 greater in TBS buffer (pH 7.5) than in CBS buffer (pH 6.5). This is consistent with the observation that the affinity of BPTI for trypsin is greater at pH 7.5 than at pH 6.5 (VINC72, VINC74). A much lower percentage of the MK control phage (which do not display BPTI) bound to

30 immobilized trypsin and this binding was independent of the pH conditions. At pH 6.5, 1675 times more of the BPTI-III MK phage than of the MK phage bound to trypsin beads while at pH 7.5, a 2103-fold difference was observed. Hence fusion phage displaying BPTI adhere not

35 only to anhydro-trypsin beads but also to active trypsin

beads and can be recovered as infectious phage. These data, in conjunction with earlier findings, strongly suggest that BPTI displayed on the surface of fusion phage is appropriately folded and functional.

5 2) Generation of P1 Mutants of BPTI

To demonstrate the specificity of interaction of BPTI-III fusion phage with immobilized serine proteases, single amino acid substitutions were introduced at the P1 position (residue 15 of mature BPTI) of the BPTI-III fusion protein by site-directed mutagenesis. A 25mer mutagenic oligonucleotide (P1) was designed to substitute a LEU codon for the LYS₁₅ codon. This alteration is desired because BPTI(K15L) is a moderately good inhibitor of human neutrophil elastase (HNE) ($K_d = 2.9 \cdot 10^{-9}$ M) (BECK88b) and a poor inhibitor of trypsin. A fusion phage displaying BPTI(K15L) should bind to immobilized HNE but not to immobilized trypsin. BPTI-III MK fusion phage would be expected to display the opposite phenotype (bind to trypsin, fail to bind to HNE). These observations would illustrate the binding specificity of BPTI-III fusion phage for immobilized serine proteases.

Mutagenesis of the P1 region of the BPTI-VIII gene contained within the intergenic region of recombinant phage MB46 was carried out using the Muta-Gene M13 In Vitro Mutagenesis Kit (Bio-Rad, Richmond, CA). MB46 phage ($7.5 \cdot 10^6$ pfu) were used to infect a 50 ml culture of CJ236 cells (O.D.600 = 0.5). Following overnight incubation at 37°C, phage were recovered and uracil-containing single-stranded DNA was extracted from the phage. The single-stranded DNA was further purified by NACS chromatography as recommended by the manufacturer (B.R.L., Gaithersburg, MD).

Two hundred nanograms of the purified single-stranded DNA were annealed to 3 picomoles of the phosphorylated

T05290-5596850

25mer mutagenic oligonucleotide (P1). Following filling in with T4 DNA polymerase and ligation with T4 DNA ligase, the sample was used to transfect competent cells which were subsequently plated on LB plates to permit the formation of plaques. Phage derived from picked plaques were applied to a Nytran membrane using a Schleicher and Schuell (Keene, NH) minifold I apparatus (Dot Blot Procedure). Phage DNA was immobilized onto the filter by baking at 80°C for 2 hours. The filter was bathed in 1 X Southern pre-hybridization buffer (5Prime-3Prime, West Chester, PA) for 2 hours. Subsequently, the filter was incubated in 1 X Southern hybridization solution (5Prime-3Prime) containing a 21mer probing oligonucleotide (LEU1) which had been radioactively labelled with gamma-³²P-ATP (N.E.N./DuPont, Boston, MA) by T4 polynucleotide kinase (New England BioLabs (NEB), Beverly, MA). Following overnight hybridization, the filter was washed 3 times with 6 X SSC at room temperature and once at 60°C in 6 X SSC prior to autoradiography. Clones exhibiting strong hybridization signals were chosen for large scale Rf preparation using the PZ523 spin column protocol (5Prime-3Prime). Restriction enzyme analysis confirmed that the structure of the Rf was correct and DNA sequencing confirmed the substitution of a LEU codon (TTG) for the LYS₁₅ codon (AAA). This Rf DNA was designated MB46(K15L).

3) Generation of the BPTI-III MA Vector

The original gene III fusion phage MK can be detected on the basis of its ability to transduce cells to kanamycin resistance (Km^R). It was deemed advantageous to generate a second gene III fusion vector which can confer resistance to a different antibiotic, namely ampicillin (Ap). One could then mix a fusion phage conferring Ap^R while displaying engineered protease inhibitor A (EPI-A) with a second fusion phage conferring Km^R while displaying

EPI-B. The mixture could be added to an immobilized serine protease and, following elution of bound fusion phage, one could evaluate the relative affinity of the two EPIs for the immobilized protease from the relative abundance of phage that transduce cells to Km^R or Ap^R .

The ap^R gene is contained in the vector pGem3Zf (Promega Corp., Madison, WI) which can be packaged as single stranded DNA contained in bacteriophage when helper phage are added to bacteria containing this vector. The recognition sites for restriction enzymes SmaI and SnaBI were engineered into the 3' non-coding region of the Ap^R (β -lactamase) gene using the technique of synthetic oligonucleotide directed site specific mutagenesis. The single stranded DNA was used as the template for in vitro mutagenesis leading to the following DNA sequence alterations (numbering as supplied by Promega): a) to create a SmaI (or XmaI) site, bases $T_{1115} \rightarrow C$ and $A_{1116} \rightarrow C$, and b) to create a SnaBI site, $G_{1125} \rightarrow T$, $C_{1129} \rightarrow T$, and $T_{1130} \rightarrow A$. The alterations were confirmed by radiolabelled probe analysis with the mutating oligonucleotide and restriction enzyme analysis; this plasmid is named pSGK3.

Plasmid SGK3 was cut with AatII and SmaI and treated with T4 DNA polymerase (NEB) to remove overhanging 3' ends (MANI82, SAMB89). Phosphorylated HindIII linkers (NEB) were ligated to the blunt ends of the DNA and following HindIII digestion, the 1.1 kb fragment was isolated by agarose gel electrophoresis followed by purification on an Ultrafree-MC filter unit as recommended by the manufacturer (Millipore, Bedford, MA). M13-MB1/2-delta Rf DNA was cut with HindIII and the linearized Rf was purified and ligated to the 1.1 kb fragment derived from pSGK3. Ligation samples were used to transfect competent cells which were plated on LB plates containing Ap. Colonies were picked and grown in LB broth containing Ap overnight

T06290"56096360

at 37°C. Aliquots of the culture supernatants were assayed for the presence of infectious phage. Rf DNA was prepared from cultures which were both Ap^R and contained infectious phage. Restriction enzyme analysis confirmed
 5 that the Rf contained a single copy of the Ap^R gene inserted into the intergenic region of the M13 genome in the same transcriptional orientation as the phage genes. This Rf DNA was designated MA.

The 5.9 kb BglIII/BsmI fragment from MA Rf DNA and the
 10 2.2 kb BglIII/BsmI fragment from BPTI-III MK Rf DNA were ligated together and a portion of the ligation mixture was used to transfect competent cells which were subsequently plated to permit plaque formation on a lawn of cells. Large and small size plaques were observed on the plates.
 15 Small size plaques were picked for further analysis since BPTI-III fusion phage give rise to small plaques due to impairment of gene III protein function. Small plaques were added to LB broth containing Ap and cultures were incubated overnight at 37°C. An Ap^R culture which
 20 contained phage which gave rise to small plaques when plated on a lawn of cells was used as a source of Rf DNA. Restriction enzyme analysis confirmed that the BPTI-III fusion gene had been inserted into the MA vector. This Rf was designated BPTI-III MA.

25 4) Construction of BPTI(K15L)-III MA

MB46(K15L) Rf DNA was digested with XhoI and EagI and the 125 bp DNA fragment was isolated by electrophoresis on a 2% agarose gel followed by extraction from an agarose slice by centrifugation through an Ultrafree-MC filter
 30 unit. The 8.0 kb XhoI/EagI fragment derived from BPTI-III MA Rf was also prepared. The above two fragments were ligated and the ligation sample was used to transfect competent cells which were plated on LB plates containing Ap. Colonies were picked and used to inoculate LB broth

09896095.062901
 106290"55055860

containing Ap. Cultures were incubated overnight at 37°C and phage within the culture supernatants was probed using the Dot Blot Procedure. Filters were hybridized to a radioactively labelled oligonucleotide (LEU1). Positive clones were identified by autoradiography after washing filters under high stringency conditions. Rf DNA was prepared from Ap^R cultures which contained phage carrying the K15L mutation. Restriction enzyme analysis and DNA sequencing confirmed that the K15L mutation had been introduced into the BPTI-III MA Rf. This Rf was designated BPTI(K15L)-III MA. Interestingly, BPTI(K15L)-III MA phage gave rise to extremely small plaques on a lawn of cells and the infectivity of the phage is 4 to 5 fold less than that of BPTI-III MK phage. This suggests that the substitution of LEU for LYS₁₅ impairs the ability of the BPTI:gene III fusion protein to mediate phage infection of bacterial cells.

5) Preparation of Immobilized Human Neutrophil Elastase

One ml of Reacti-Gel 6 x CDI activated agarose (Pierce Chemical Co.) in acetone (200 µl packed beads) was introduced into an empty Select-D spin column (5Prime-3Prime). The acetone was drained out and the beads were washed twice rapidly with 1.0 ml of ice cold water and 1.0 ml of ice cold 100 mM boric acid, pH 8.5, 0.9% NaCl. Two hundred µl of 2.0 mg/ml human neutrophil elastase (HNE) (CalBiochem, San Diego, CA) in borate buffer were added to the beads. The column was sealed and mixed end over end on a Labquake Shaker at 4°C for 36 hours. The HNE solution was drained off and the beads were washed with ice cold 2.0 M Tris, pH 8.0 over a 2 hour period at 4°C to block remaining reactive groups. A 50% slurry of the beads in TBS/BSA was prepared. To this was added an equal volume of sterile 100% glycerol and the beads were stored as a 25% slurry at -20°C. Prior to use, the beads were

washed 3 times with TBS/BSA and a 50% slurry in TBS/BSA was prepared.

6) Characterization of the Affinity of BPTI-III MK and BPTI(K15L)-III MA Phage for Immobilized Trypsin and Human

5 Neutrophil Elastase

Thirty μ l of BPTI-III MK phage in TBS/BSA ($1.7 \cdot 10^{11}$ pfu/ml) was added to 5 μ l of a 50% slurry of either immobilized human neutrophil elastase or immobilized trypsin (Pierce Chemical Co.) also in TBS/BSA. Similarly
 10 30 μ l of BPTI(K15L)-III MA phage in TBS/BSA ($3.2 \cdot 10^{10}$ pfu/ml) was added to either immobilized HNE or trypsin. Samples were mixed on a Labquake shaker for 3 hours. The beads were washed with 0.5 ml of TBS/BSA for 5 minutes and recovered by centrifugation. The supernatant was removed
 15 and the beads were washed 5 times with 0.5 ml of TBS/0.1% Tween-20. Finally, the beads were resuspended in 0.5 ml of elution buffer (0.1 M HCl containing 1.0 mg/ml BSA adjusted to pH 2.2 with glycine), mixed for 5 minutes and recovered by centrifugation. The supernatant fraction was
 20 removed, neutralized with 130 μ l of 1 M Tris, pH 8.0, diluted in LB broth, and titered for plaque-forming units on a lawn of cells.

Table 202 illustrates that 82 times more of the BPTI-III MK input phage bound to the trypsin beads than to the
 25 HNE beads. By contrast, the BPTI(K15L)-III MA phage bound preferentially to HNE beads by a factor of 36. These results are consistent with the known affinities of wild type and the K15L variant of BPTI for trypsin and HNE. Hence BPTI-III fusion phage bind selectively to immobil-
 30 ized proteases and the nature of the BPTI variant displayed on the surface of the fusion phage dictates which particular protease is the optimum receptor for the fusion phage.

TOP SECRET 56096860

achieved sufficient mass. Poor growth and insert instability can be circumvented to a large extent, giving this system an advantage over the Gem-based vector described above.

5 An overnight bacterial culture of XL1-Blue^(TM) or SEF' is grown in LB medium containing tetracycline (50 μ g per ml) to ensure the presence of pili as sites for bacteriophage binding and infection. This culture is diluted 100-fold into NZCYM medium containing tetracycline
10 and bacterial growth allowed to proceed in an incubator shaker until a cell density of 1.0 (Ab 600nm) has been achieved. Phage, containing the expression vector and gene of interest, are added to the bacterial culture at a multiplicity of infection (MOI) of 10 and allowed to
15 infect the cells for 30 minutes. Gene expression is then induced by the addition of IPTG to a final concentration of 0.5 mM and the culture allowed to grow overnight. Media collection and cell fractionation is as described elsewhere.

20 Bacterial Cell Fractionation.

After heterologous gene expression the bacterial cell culture can be separated into the following fractions: conditioned medium, periplasmic fraction and post-periplasmic cell lysate. This is achieved using the following
25 procedures.

The culture is centrifuged to pellet the bacteria, allowing the supernatant to be stored as conditioned medium. This fraction contains any exported proteins. The pellet is taken up in 20% sucrose, 30mM Tris pH 8 and
30 1mM EDTA (80 ml of buffer per gram of fresh weight pellet) and allowed to sit at room temperature for 10 minutes. The cells are repelleted and taken up in the same volume of ice cold 5mM MgSO₄ and left on ice for 10 minutes. Following centrifugation, to pellet the cells, the

106250-56096660

supernatant (periplasmic fraction) is stored. A second round of osmotic shock fractionation can be undertaken if desired.

The post-periplasmic pellet can be further lysed as follows. The pellet is resuspended in 1.5 ml of 20% sucrose, 40 mM Tris pH 8, 50mM EDTA and 2.5 mg of lysozyme (per gram fresh weight of starting pellet). After 15 minutes at room temperature 1.15 ml of 0.1% Triton X is added together with 300 μ l of 5M NaCl and incubated for a further 15 minutes. 2.5 ml of 0.2 M triethanolamine (pH 7.8), 150 μ l of 1M CaCl_2 , 100 μ l of 1M MgCl_2 and 5 μ g of DNA'se are added and allowed to incubate, with end-over-end mixing, for 20 minutes to reduce viscosity. This is followed by centrifugation with the supernatant being retained as the post-periplasmic lysate.

The present invention is not, of course, limited to any particular expression system, whether bacterial or not.

EXAMPLE IX

CONSTRUCTION OF AN ITI-DOMAIN I/GENE III DISPLAY VECTOR

1. ITI domain I as an IPBD

Inter- α -trypsin inhibitor (ITI) is a large (M_r ca 240,000) circulating protease inhibitor found in the plasma of many mammalian species (for recent reviews see ODOM90, SALI90, GEBH90, GEBH86). The intact inhibitor is a glycoprotein and is currently believed to consist of three glycosylated subunits that interact through a strong glycosaminoglycan linkage (ODOM90, SALI90, ENGH89, SELL87). The anti-trypsin activity of ITI is located on the smallest subunit (ITI light chain, unglycosylated M_r ca 15,000) which is identical in amino acid sequence to an acid stable inhibitor found in urine (UTI) and serum (STI)

(GEBH86, GEBH90). The mature light chain consists of a 21 residue N-terminal sequence, glycosylated at SER₁₀, followed by two tandem Kunitz-type domains the first of which is glycosylated at ASN₄₅ (ODOM90). In the human
 5 protein, the second Kunitz-type domain has been shown to inhibit trypsin, chymotrypsin, and plasmin (ALBR83a, ALBR83b, SELL87, SWAI88). The first domain lacks these activities but has been reported to inhibit leukocyte elastase ($10^{-6} > K_i > 10^{-9}$) (ALBR83a,b, ODOM90). cDNA
 10 encoding the ITI light chain also codes for α -1-microglobulin (TRAB86, KAUM86, DIAR90); the proteins are separated post-translationally by proteolysis.

The N-terminal Kunitz-type of the ITI light chain (ITI-D1, comprising residues 22 to 76 of the UTI sequence
 15 shown in Fig. 1 of GEBH86) possesses a number of characteristics that make it useful as an IPBD. The domain is highly homologous to both BPTI and the EpINE series of proteins described elsewhere in the present application. Although an x-ray structure of the isolated domain is not
 20 available, crystallographic studies of the related Kunitz-type domain isolated from the Alzheimer's amyloid β -protein (AA β P) precursor show that this polypeptide assumes a crystal structure almost identical to that of BPTI (HYNE90). Thus, it is likely that the solution
 25 structure of the isolated ITI-D1 polypeptide will be highly similar to the structures of BPTI and AA β P. In this case, the advantages described previously for use of BPTI as an IPBD apply to ITI-D1. ITI-D1 provides additional advantages as an IDBP for the development of
 30 specific anti-elastase inhibitory activity. First, this domain has been reported to inhibit both leukocyte elastase (ALBR83a,b, ODOM90) and Cathepsin-G (SWAI88, ODOM90); activities which BPTI lacks. Second, ITI-D1 lacks affinity for the related serine proteases trypsin,
 35 chymotrypsin, and plasmin (ALBR83a,b, SWAI88), an advan-

106630 55036850

tage for the development of specificity in inhibition. Finally, ITI-D1 is a human-derived polypeptide so derivatives are anticipated to show minimal antigenicity in clinical applications.

5 2. Construction of the display vector.

For purposes of this discussion, numbering of the nucleic acid sequence for the ITI light chain gene is that of TRAB86 and of the amino acid sequence is that shown for UTI in Fig. 1 of GEBH86. DNA manipulations were conducted according to standard methods as described in SAMB89 and AUSU87.

The protein sequence of human ITI-D1 consists of 56 amino acid residues extending from LYS₂₂ to ARG₇₇ of the complete ITI light chain sequence. This sequence is encoded by the 168 bases between positions 750 and 917 in the cDNA sequence presented in TRAB86. The majority of the domain is contained between a BglI site spanning bases 663 to 773 and a PstI site spanning bases 903 to 908. The insertion of the ITI-D1 sequence into M13 gene III was conducted in two steps. First a linker containing the appropriate ITI sequences outside the central BglI to PstI region was ligated into the NarI site of phage MA RF DNA. In the second step, the remainder of the ITI-D1 sequence was incorporated into the linker-bearing phage RF DNA.

The linker DNA consisted of two synthetic oligonucleotides (top and bottom strands) which, when annealed, produced a 54 bp double-stranded fragment with the following structure (5' to 3'):

30 NAR I OVERHANG/ITI-5'/BGLI/STUFFER/PSTI/ITI-3'/NAR I
OVERHANG

The NarI OVERHANG sequences provide compatible ends for ligation into a cut NarI site. The ITI-5' sequence

consists of ds DNA corresponding to the thirteen positions from A750 to T662 immediately 5' adjacent to the BglI site in the ITI-D1 sequence. Two changes, both silent, are introduced in this sequence: T to C at position 658 (changes codon for ASP₂₄ from GAT to GAC) and G to T at position 661 (changes codon for SER₂₅ from TCG to TCT). The sequences BGLI and PSTI are identical to the BglI and PstI sites, respectively, in the ITI-D1 sequence. The ITI-3' sequence consists of dsDNA corresponding to the nine positions from A909 to T917 immediately 3' adjacent to the PstI site in the ITI-D1 sequence. The one base change included in this sequence, A to T at position 917, is silent and changes the codon for ARG₇₇ from CGA to CGT. The STUFFER sequence consists of dsDNA encoding three residues (5' to 3'): LEU (TTA), TRP(TGG), and SER(TCA). The reverse complement of the STUFFER sequence encodes two translation termination codons (TGA and TAA). Phage expressing gene III containing the linker in opposite orientation to that shown above will not produce a functional gene III product.

Phage MA RF DNA was digested with NarI and the linear ca. 8.2 kb fragment was gel purified and subsequently dephosphorylated using HK phosphatase (Epicentre). The linker oligonucleotides were annealed to form the linker fragment described above, which was then kinased using T4 Polynucleotide Kinase. The kinased linker was ligated to the NarI-digested MA RF DNA in a 10:1 (linker:-RF) molar ratio. After 18 hrs at 16°C, the ligation was stopped by incubation at 65°C for 10 min and the ligation products were ethanol precipitated in the presence of 10 µg of yeast tRNA. The dried precipitate was dissolved in 5 µl of water and used to transform D1210 cells by electroporation. After 60 min of growth in SOC at 37°C, transformed cells were plated onto LB plates supplemented with ampicillin (Ap, 200 µg/ml). RF DNA prepared from

AP^r isolates was subjected to restriction enzyme analysis. The DNA sequences of the linker insert and the immediately surrounding regions were confirmed by DNA sequencing. Phage strains containing the ITI Linker sequence inserted
5 into the NarI site in gene III are called MA-IL.

Phage MA-IL RF DNA was partially digested with BglI and the ca. 8.2 kb linear fragment was gel purified. This fragment was digested with PstI and the large linear fragment was gel purified. The BglI to PstI fragment of
10 ITI-D1 was isolated from pMG1A (a plasmid carrying the sequence shown in TRAB86). pMG1A was digested to completion with BglI and the ca. 1.6 kb fragment was isolated by agarose gel electrophoresis and subsequent GeneClean (Bio101, La Jolla, CA) purification. The purified BglI
15 fragment was digested to completion with PstI and EcoRI and the resulting mixture of fragments was used in a ligation with the BglI and PstI cut MA-IL RF DNA described above. Ligation, transformation, and plating were as described above. After 18 hr. of growth on LB Ap plates
20 at 37°C, Ap^r colonies were harvested with LB broth supplemented with Ap (200 µg/ml) and the resulting cell suspension was grown for two hours at 37°C. Cells were pelleted by centrifugation (10 min at 5000xg, 4°C). The supernatant fluid was transferred to sterile centrifuga-
25 tion tubes and recentrifuged as above. The supernatant fluid from the second centrifugation step was retained as the phage stock POP1.

PCR was used to demonstrate the presence of phage containing the complete ITI-D1-III fusion gene. Upstream
30 PCR primers, 1UP and 2UP, are located spanning nucleotides 1470 to 1494 and 1593 to 1618 of the phage M13 DNA sequence, respectively. A downstream PCR primer 3DN spans nucleotides 1779 to 1804. Two ITI-D1-specific primers, IAI-1 and IAI-2, are located spanning positions 789 to 810

and 894 to 914, respectively, in the ITI light chain sequence of TRAB86. IAI-1 and IAI-2 are used as downstream primers in PCR reactions with 1UP or 2UP. IAI-1 is entirely contained within the BglI to PstI region of the ITI-D1 sequence, while IAI-2 spans the PstI site in the ITI-D1 sequence. When aliquots of POP1 phage were used as substrates for PCR, template-specific products of characteristic size were produced in reactions containing 1UP or 2UP plus IAI-1 or IAI-2 primer pairs. No such products are obtained using MA-IL phage as template. No PCR products with sizes corresponding to complete ITI-D1-gene III templates were obtained using POP1 phage and the 1UP or 2UP plus 3DN primer pairs. This last result reflects the low abundance (<1%) of phage containing the complete ITI-D1 sequence in POP1.

Preparative PCR was used to generate substrate amounts of the 330 bp PCR product of a reaction using the 1UP and IAI-2 primer pair to amplify the POP1 template. The 330 bp PCR product was gel purified and then cut to completion with BglI and PstI. The 138 bp BglI to PstI fragment from ITI-D1 was isolated by agarose gel electrophoresis followed by Qiaex extraction (Qiagen, Studio City, CA). MA-IL phage RF DNA was digested to completion with PstI. The ca. 8.2 kb linear fragment was gel purified and subsequently digested to completion with BglI. The BglI digest was extracted once with phenol:chloroform (1:1), the aqueous phase was ethanol precipitated, and the pellet was dissolved in TE (pH8.0). An aliquot of this solution was used in a ligation reaction with the 138 bp BglI to PstI fragment as described above. The ethanol precipitated ligation products were used to transform XL1-Blue^(TM) cells by electroporation and after 1 hr growth in SOC at 37°C, cells were plated on LB Ap plates. A phage population, POP2, was prepared from Ap^r colonies as described previously.

Phage stocks obtained from individual plaques produced on titration of POP2 were tested by PCR for the presence of the complete ITI-D1-III gene fusion. PCR results indicate the entire fusion gene was present in seven of nine isolates tested. RF DNA from the seven isolates testing positive was subjected to restriction enzyme analysis. The complete sequence of the ITI-D1 insertion into gene III was confirmed in four of the seven isolates by DNA sequence analysis. Phage isolates containing the ITI-D1-III fusion gene are called MA-ITI.

3. Expression and display of ITI-DI.

Expression of the ITI domain I-Gene III fusion protein and its display on the surface of phage were demonstrated by Western analysis and phage titer neutralization experiments.

For Western analysis, aliquots of PEG-purified phage preparations containing up to $4 \cdot 10^{10}$ infective particles were subjected to electrophoresis on a 12.5% SDS-urea-polyacrylamide gel. Proteins were transferred to a sheet of Immobilon-P transfer membrane (Millipore, Bedford, MA) by electrotransfer. Western blots were developed using a rabbit anti-ITI serum (SALI87) which had previously been incubated with an *E. coli* extract, followed by goat anti-rabbit IgG conjugated to horse radish peroxidase (#401315, Calbiochem, La Jolla, Ca). An immunoreactive protein with an apparent size of ca. 65-69 kD is detected in preparations of MA-ITI phage but not with preparations of the parental MA phage. The size of the immunoreactive protein is consistent with the expected size of the processed ITI-D1-III fusion protein (ca. 67 kD, as previously observed for the BPTI-III fusion protein).

Rabbit anti-BPTI serum has been shown to block the ability of MK-BPTI phage to infect *E. coli* cells (Example II). To test for a similar effect of rabbit anti-ITI

serum on the infectivity of MA-ITI phage, 10 μ l aliquots of MA or MA-ITI phage were incubated in 100 μ l reactions containing 10 μ l aliquots of PBS, normal rabbit serum (NRS), or anti-ITI serum. After a three hour incubation at 37°C, phage suspensions were titered to determine residual plaque-forming activity. These data are summarized in Table 211. Incubation of MA-ITI phage with rabbit anti-ITI serum reduces titers 10- to 100-fold, depending on initial phage titer. A much smaller decrease in phage titer (10 to 40%) is observed when MA-ITI phage are incubated with NRS. In contrast, the titer of the parental MA phage is unaffected by either NRS or anti-ITI serum.

Taken together, the results of the Western analysis and the phage-titer neutralization experiments are consistent with the expression of an ITI-DI-III fusion protein in MA-ITI phage, but not in the parental MA phage, such that ITI-specific epitopes are present on the phage surface. The ITI-specific epitopes are located with respect to III such that antibody binding to these epitopes prevents phage from infecting *E. coli* cells.

4. Fractionation of MA-ITI phage bound to agarose-immobilized protease beads.

To test if phage displaying the ITI-DI-III fusion protein interact strongly with the proteases human neutrophil elastase (HNE) or cathepsin-G, aliquots of display phage were incubated with agarose-immobilized HNE or cathepsin-G beads (HNE beads or Cat-G beads, respectively). The beads were washed and bound phage eluted by pH fractionation as described in Examples II and III. The procession in lowering pH during the elution was: pH 7.0, 6.0, 5.5, 5.0, 4.5, 4.0, 3.5, 3.0, 2.5, and 2.0. Following elution and neutralization, the various input, wash, and pH elution fractions were titered.

The results of several fractionations are summarized in Table 212 (EpiNE-7 or MA-ITI phage bound to HNE beads) and Table 213 (EpiC-10 or MA-ITI phage bound to Cat-G beads). For the two types of beads (HNE or Cat-G), the pH elution profiles obtained using the control display phage (EpiNE-7 or EpiC-10, respectively) were similar to those seen previously (Examples II and III). About 0.3% of the EpiNE-7 display phage applied to the HNE beads were eluted during the fractionation procedure and the elution profile had a maximum for elution at about pH 4.0. A smaller fraction, 0.02%, of the EpiC-10 phage applied to the Cat-G beads were eluted and the elution profile displayed a maximum near pH 5.5.

The MA-ITI phage show no evidence of great affinity for either HNE or cathepsin-G immobilized on agarose beads. The pH elution profiles for MA-ITI phage bound to HNE or Cat-G beads show essentially monotonic decreases in phage recovered with decreasing pH. Further, the total fractions of the phage applied to the beads that were recovered during the fractionation procedures were quite low: 0.002% from HNE beads and 0.003% from Cat-G beads.

Published values of K_i for inhibition neutrophil elastase by the intact, large ($M_r=240,000$) ITI protein range between 60 and 150 nM and values between 20 and 6000 nM have been reported for the inhibition of Cathepsin G by ITI (SWAI88, ODOM90). Our own measurements of pH fraction of display phage bound to HNE beads show that phage displaying proteins with low affinity ($>\mu M$) for HNE are not bound by the beads while phage displaying proteins with greater affinity (nM) bind to the beads and are eluted at about pH 5. If the first Kunitz-type domain of the ITI light chain is entirely responsible for the inhibitory activity of ITI against HNE, and if this domain is correctly displayed on the MA-ITI phage, then it

appears that the minimum affinity of an inhibitor for HNE that allows binding and fractionation of display phage on HNE beads is 50 to 100 nM.

5. Alteration of the P1 region of ITI-DI.

5 If ITI-DI and EpiNE-7 assume the same configuration in solution as BPTI, then these two polypeptides have identical amino acid sequences in both the primary and secondary binding loops with the exception of four residues about the P1 position. For ITI-DI the sequence
10 for positions 15 to 20 is (position 15 in ITI-DI corresponds to position 36 in the UTI sequence of GEBH86): MET15, GLY16, MET17, THR18, SER19, ARG20. In EpiNE-7 the equivalent sequence is: VAL15, ALA16, MET17, PHE18, PRO19, ARG20. These two proteins appear to differ greatly in
15 their affinities for HNE. To improve the affinity of ITI-DI for HNE, the EpiNE-7 sequence shown above was incorporated into the ITI-DI sequence at positions 15 through 20.

The EpiNE-7 sequence was incorporated into the ITI-DI sequence in MA-ITI by cassette mutagenesis. The mutagenic
20 cassette consisted of two synthetic 51 base oligonucleotides (top and bottom stands) which were annealed to make double stranded DNA containing an Eag I overhang at the 5' end and a Sty I overhang at the 3' end. The DNA sequence between the Eag I and Sty I overhangs is identical to the
25 ITI-DI sequence between these sites except at four codons: the codon for position 15, AT (MET), was changed to GTC (VAL), the codon for position 16, GGA (GLY), was changed to GCT (ALA), the codon for position 18, ACC (THR) was changed to TTC (PHE), and the codon for position 19, AGC
30 (SER), was changed to CCA (PRO). MA-ITI RF DNA was digested with Eag I and Sty I. The large, linear fragment was gel purified and used in a ligation with the mutagenic cassette described above. Ligation products were used to transform XL1-Bluetm cells as described previously. Phage

stocks obtained from overnight cultures of Ap^r transduc-
tants were screened by PCR for incorporation of the
altered sequence and the changes in the codons for
positions 15, 16, 18, and 19 were confirmed by DNA
5 sequencing. Phage isolates containing the ITI-DI-III
fusion gene with the EpINE-7 changes around the P1
position are called MA-ITI-E7.

6. Fractionation of MA-ITI-E7 phage.

To test if the changes at positions 15, 16, 18, and
10 19 of the ITI-DI-III fusion protein influence binding of
display phage to HNE beads, abbreviated pH elution
profiles were measured. Aliquots of EpINE-7, MA-ITI, and
MA-ITI-E7 display phage were incubated with HNE beads for
three hours at room temperature. The beads were washed
15 and phage were eluted as described (Example III), except
that only three pH elutions were performed: pH 7.0, 3.5,
and 2.0. The results of these elutions are shown in Table
214.

Binding and elution of the EpINE-7 and MA-ITI display
20 phage were found to be as previously described. The total
fraction of input phages was high (0.4%) for EpINE-7
phage and low (0.001%) for MA-ITI phage. Further, the
EpINE-7 phage showed maximum phage elution in the pH 3.5
fraction while the MA-ITI phage showed only a monotonic
25 decrease in phage yields with decreasing pH, as seen
above.

The two strains of MA-ITI-E7 phage show increased
levels of binding to HNE beads relative to MA-ITI phage.
The total fraction of the input phage eluted from the
30 beads is 10-fold greater for both MA-ITI-E7 phage strains
than for MA-ITI phage (although still 40-fold lower than
EpINE-7 phage). Further, the pH elution profiles of the
MA-ITI-E7 phage strains show maximum elutions in the pH
3.5 fractions, similar to EpINE-7 phage.

005505.03401

To further define the binding properties of MA-ITI-E7 phage, the extended pH fractionation procedure described previously was performed using phage bound to HNE beads. These data are summarized in Table 215. The pH elution profile of EpiNE-7 display phage is as previously described. In this more resolved, pH elution profile, MA-ITI-E7 phage show a broad elution maximum centered around pH 5. Once again, the total fraction of MA-ITI-E7 phage obtained on pH elution from HNE beads was about 40-fold less than that obtained using EpiNE-7 display phage.

The pH elution behavior of MA-ITI-E7 phage bound to HNE beads is qualitatively similar to that seen using BPTI[K15L]-III-MA phage. BPTI with the K15L mutation has an affinity for HNE of $\approx 3 \cdot 10^{-9}$ M. Assuming all else remains the same, the pH elution profile for MA-ITI-E7 suggests that the affinity of the free ITI-DI-E7 domain for HNE might be in the nM range. If this is the case, the substitution of the EpiNE-7 sequence in place of the ITI-DI sequence around the P1 region has produced a 20- to 50-fold increase in affinity for HNE (assuming $K_i = 60$ to 150 nM for the unaltered ITI-DI).

If EpiNE-7 and ITI-DI-E7 have the same solution structure, these proteins present the identical amino acid sequences to HNE over the interaction surface. Despite this similarity, EpiNE-7 exhibits a roughly 1000-fold greater affinity for HNE than does ITI-DI-E7. Again assuming similar structure, this observation highlights the importance of non-contacting secondary residues in modulating interaction strengths.

Native ITI light chain is glycosylated at two positions, SER10 and ASN45 (GEBH86). Removal of the glycosaminoglycan chains has been shown to decrease the affinity of the inhibitor for HNE about 5-fold (SELL87). Another potentially important difference between EpiNE-7

and ITI-DI-E7 is that of net charge. The changes in BPTI that produce EpiNE-7 reduce the total charge on the molecule from +6 to +1. Sequence differences between EpiNE-7 and ITI-DI-E7 further reduce the charge on the latter to -1. Furthermore, the change in net charge between these two molecules arises from sequence differences occurring in the central portions of the molecules. Position 26 is LYS in EpiNE-7 and is THR in ITI-DI-E7, while at position 31 these residues are GLN and GLU, respectively. These changes in sequence not only alter the net charge on the molecules but also position negatively charged residue close to the interaction surface in ITI-DI-E7. It may be that the occurrence of a negative charge at position 31 (which is not found in any other of the HNE inhibitors described here) destabilized the inhibitor-protease interaction.

EXAMPLE X

GENERATION OF A VARIEGATED ITI-DI POPULATION

The following is a hypothetical example demonstrating how to obtain a derivative of ITI having high affinity for HNE.

The results of Example IX demonstrate that the nature of the protein sequence around the P1 position in ITI-DI can significantly influence the strength of the interaction between ITI-DI and HNE. While incorporation of the EpiNE-7 sequence increases the affinity of ITI-DI for HNE, it is unlikely that this particular sequence is optimal for binding.

We generate a large population of potential binding proteins having differing sequences in the P1 region of ITI-DI using the oligonucleotide ITIMUT. ITIMUT is designed to incorporate variegation in ITI-DI at the six

positions about and including the P1 residue: 13, 15, 16, 17, 18, and 19. ITIMUT is synthesized as one long (top strand) 73 base oligonucleotide and one shorter (24 base) bottom strand oligonucleotide. The top strand sequence
 5 extends from position 770 (G) to position 842 (G) in the sequence of TREB86. This sequence includes the codons for the positions of variegation as well as the recognition sequences for the flanking restriction enzymes Eag I (778 to 783) and Sty I (829 to 834). The bottom strand
 10 oligonucleotide comprises the complement of the sequence from positions 819 to 842.

To generate the mutagenic cassette, the top and bottom strand oligonucleotides are annealed and the resulting duplex is completed in an extension reaction
 15 using DNA polymerase. Following digestion of the 73 bp dsDNA with Eag I and Sty I, the purified 51 bp mutagenic cassette is ligated with the large linear fragment obtained from a similar digestion of MA-ITI RF DNA. Ligation products are used to transform competent cells by
 20 electroporation and phage stocks produced from Ap^r transductants are analyzed for the presence and nature of novel sequences as described previously.

The variegation in the ITIMUT cassette is confined to the codons for the six positions in ITI-DI (13, 15, 16, 17, 18, and 19), and employs three different nucleotide
 25 mixes: N, R, and S. For this mutagenesis, the composition of the N-mix is 36%A, 17%C, 23%G, and 24%T, and corresponds to the N-mix composition in the optimized NNS codon described elsewhere. The R-mix composition is 50%A,
 30 50%G, and the S-mix composition is 50%C, 50%G.

The codon for ITI-DI position 13 (CCC, PRO) is changed to SNG in ITIMUT. This codon encodes the eight residues PRO, VAL, GLU, ALA, GLY, LEU, GLN, and ARG. The encoded group includes the parental residue (PRO) as well

as the more commonly observed variants at the position, ARG and LEU (see Table 15), and also provides for the occurrence of acidic (GLU), large polar (GLN) and nonpolar (VAL), and small (ALA, GLY) residues.

- 5 The codons for positions 15 and 17 (ATG, MET) are changed to the optimized NNS codon. All 20 natural amino acid residues and a translation termination are allowed.

10 The codon for position 16 (CGA, GLY) is changed to RNS in ITIMUT. This codon encodes the twelve amino acids GLY, ALA, ASP, GLU, VAL, MET, ILE, THR, SER, ARG, ASN, and LYS. The encoded group includes the most commonly observed residues at this position, ALA and GLY, and provides for the occurrence of both positively (ARG, LYS) and negatively (GLU, ASP) charged amino acids. Large
15 nonpolar residues are also included (ILE, MET, VAL).

20 Finally, at positions 18 and 19, the ITI-DI sequence is changed from ACC·AGC (THR·SER) to NNT·NNT. The NNT codon encodes the fifteen amino acid residues PHE, SER, TYR, CYS, LEU, PRO, HIS, ARG, ILE, THR, ASN, VAL, ALA, ASP, and GLY. This group includes the parental residues and the further advantages of the NNT codon have been discussed elsewhere.

The ITIMUT DNA sequence encodes a total of:

$$8 * 20 * 12 * 20 * 15 * 15 = 8,640,000$$

- 25 different protein sequences in a total of:

$$2^{25} = 33,554,422$$

different DNA sequences. The total number of protein sequences encoded by ITIMUT is only 7.4-fold fewer than the total possible number of natural sequences obtained
30 from variation at six positions ($= 20^6 = 6.4 \cdot 10^7$). However, this degree of variation in protein sequence is obtained from a minimum of 1.07×10^9 ($\text{NNS}^6 = 2^{30}$) DNA sequences, a 32-fold greater number than that comprising ITIMUT. Thus, ITIMUT is an efficient vehicle for the

generation of a large and diverse population of potential binding proteins.

EXAMPLE XI

DEVELOPMENT AND SELECTION OF BPTI MUTANTS FOR

5 BINDING TO HORSE HEART MYOGLOBIN (HHMB)

The following example is hypothetical and illustrates alternative embodiments of the invention not given in other examples.

HHMb is chosen as a typical protein target; any other
10 protein could be used. HHMb satisfies all of the criteria for a target: 1) it is large enough to be applied to an affinity matrix, 2) after attachment it is not reactive, and 3) after attachment there is sufficient unaltered surface to allow specific binding by PBDs.

15 The essential information for HHMb is known: 1) HHMb is stable at least up to 70°C, between pH 4.4 and 9.3, 2) HHMb is stable up to 1.6 M Guanidinium Cl, 3) the pI of HHMb is 7.0, 4) for HHMb, $M_r = 16,000$, 5) HHMb requires haem, 6) HHMb has no proteolytic activity.

20 In addition, the following information about HHMb and other myoglobins is available: 1) the sequence of HHMb is known, 2) the 3D structure of sperm whale myoglobin is known; HHMb has 19 amino acid differences and it is generally assumed that the 3D structures are almost
25 identical, 3) HHMb has no enzymatic activity, 4) HHMb is not toxic.

We set the specifications of an SBD as :

1) $T = 25^\circ\text{C}$; 2) $\text{pH} = 8.0$; 3) Acceptable solutes ((A) for binding: i) phosphate, as buffer, 0 to 20 mM, and ii)
30 KCl, 10 mM; (B) for column elution : i) phosphate, as buffer, 0 to 30 mM, ii) KCl, up to 5 M, and iii) Guanidinium Cl, up to 0.8 M.); 4) Acceptable $K_d < 1.0 \cdot 10^{-8}$ M.

As stated in Sec. III.B, the residues to be varied

TOP SECRET 56095550

are picked, in part, through the use of interactive computer graphics to visualize the structures. In this example, all residue numbers refer to BPTI. We pick a set of residues that forms a surface such that all
5 residues can contact one target molecule. Information that we refer to during the process of choosing residues to vary includes: 1) the 3D structure of BPTI, 2) solvent accessibility of each residue as computed by the method of Lee and Richards (LEEB71), 3) a compilation of se-
10 quences of other proteins homologous to BPTI, and 4) knowledge of the structural nature of different amino acid types.

Tables 16 and 34 indicate which residues of BPTI: a) have substantial surface exposure, and b) are known to
15 tolerate other amino acids in other closely related proteins. We use interactive computer graphics to pick sets of eight to twenty residues that are exposed and variable and such that all members of one set can touch a molecule of the target material at one time. If BPTI has
20 a small amino acid at a given residue, that amino acid may not be able to contact the target simultaneously with all the other residues in the interaction set, but a larger amino acid might well make contact. A charged amino acid might affect binding without making direct
25 contact. In such cases, the residue should be included in the interaction set, with a notation that larger residues might be useful. In a similar way, large amino acids near the geometric center of the interaction set may prevent residues on either side of the large central
30 residue from making simultaneous contact. If a small amino acid, however, were substituted for the large amino acid, then the surface would become flatter and residues on either side could make simultaneous contact. Such a residue should be included in the interaction set with a
35 notation that small amino acids may be useful.

Table 35 was prepared from standard model parts and shows the maximum span between C_β and the tip of each type of side group. C_β is used because it is rigidly attached to the protein main-chain; rotation about the $C_\alpha-C_\beta$ bond is the most important degree of freedom for determining the location of the side group.

Table 34 indicates five surfaces that meet the given criteria. The first surface comprises the set of residues that actually contacts trypsin in the complex of trypsin with BPTI as reported in the Brookhaven Protein Data Bank entry "1TPA". This set is indicated by the number "1". The exposed surface of the residues in this set (taken from Table 16) totals 1148 Å². Although this is not strictly the area of contact between BPTI and trypsin, it is approximately the same.

Other surfaces, numbered 2 to 5, were picked by first picking one exposed, variable residue and then picking neighboring residues until a surface was defined. The choice of sets of residues shown in Table 34 is in no way exhaustive or unique; other sets of variable, surface residues can be picked. Set #2 is shown in stereo view, Figure 14, including the α carbons of BPTI, the disulfide linkages, and the side groups of the set. We take the orientation of BPTI in Figure 14 as a standard orientation and hereinafter refer to K15 as being at the top of the molecule, while the carboxy and amino termini are at the bottom.

Solvent accessibilities are useful, easily tabulated indicators of a residue's exposure. Solvent accessibilities must be used with some caution; small amino acids are under-represented and large amino acids over-represented. The user must consider what the solvent accessibility of a different amino acid would be when substituted into the structure of BPTI.

To create specific binding between a derivative of BPTI and HHMb, we will vary the residues in set #2. This set includes the twelve principal residues 17(R), 19(I), 21(Y), 27(A), 28(G), 29(L), 31(Q), 32(T), 34(V), 48(A), 49(E), and 52(M) (Sec. III.B). None of the residues in set #2 is completely conserved in the sample of sequences reported in Table 34; thus we can vary them with a high probability of retaining the underlying structure. Independent substitution at each of these twelve residues of the amino acid types observed at that residue would produce approximately $4.4 \cdot 10^9$ amino acid sequences and the same number of surfaces.

BPTI is a very basic protein. This property has been used in isolating and purifying BPTI and its homologues so that the high frequency of arginine and lysine residues may reflect bias in isolation and is not necessarily required by the structure. Indeed, SCI-III from Bombyx mori contains seven more acidic than basic groups (SASA84).

Residue 17 is highly variable and fully exposed and can contain R, K, A, Y, H, F, L, M, T, G, Y, P, or S. All types of amino acids are seen: large, small, charged, neutral, and hydrophobic. That no acidic groups are observed may be due to bias in the sample.

Residue 19 is also variable and fully exposed, containing P, R, I, S, K, Q, and L.

Residue 21 is not very variable, containing F or Y in 31 of 33 cases and I and W in the remaining cases. The side group of Y21 fills the space between T32 and the main chain of residues 47 and 48. The OH at the tip of the Y side group projects into the solvent. Clearly one can vary the surface by substituting Y or F so that the surface is either hydrophobic or hydrophilic in that region. It is also possible that the other aromatic

amino acid (viz. H) or the other hydrophobics (L, M, or V) might be tolerated.

Residue 27 most often contains A, but S, K, L, and T are also observed. On structural grounds, this residue will probably tolerate any hydrophilic amino acid and perhaps any amino acid.

Residue 28 is G in BPTI. This residue is in a turn, but is not in a conformation peculiar to glycine. Six other types of amino acids have been observed at this residue: K, N, Q, R, H, and N. Small side groups at this residue might not contact HHMb simultaneously with residues 17 and 34. Large side groups could interact with HHMb at the same time as residues 17 and 34. Charged side groups at this residue could affect binding of HHMb on the surface defined by the other residues of the principal set. Any amino acid, except perhaps P, should be tolerated.

Residue 29 is highly variable, most often containing L. This fully exposed position will probably tolerate almost any amino acid except, perhaps, P.

Residues 31, 32, and 34 are highly variable, exposed, and in extended conformations; any amino acid should be tolerated.

Residues 48 and 49 are also highly variable and
25 fully exposed, any amino acid should be tolerated.

Residue 52 is in an α helix. Any amino acid, except perhaps P, might be tolerated.

Now we consider possible variation of the secondary set (Sec. 13.1.2) of residues that are in the neighborhood of the principal set. Neighboring residues that might be varied at later stages include 9(P), 11(T), 15(K), 16(A), 18(I), 20(R), 22(F), 24(N), 26(K), 35(Y), 47(S), 50(D), and 53(R).

5

10

15

20

25

30

this residue could affect binding at surface defined by residue set #2.

Residue 22 is only slightly varied, being Y, F, or H in 30 of 33 cases. Nevertheless, A, N, and S have been
5 observed at this residue. Amino acids such as L, M, I, or Q could be tried here. Alterations at residue 22 may affect the mobility of residue 21; changes in charge at residue 22 could affect binding at the surface defined by residue set #2.

10 Residue 24 shows some variation, but probably can not interact with one molecule of the target simultaneously with all the residues in the principal set. Variation in charge at this residue might have an effect on binding at the surface defined by the principal set.

15 Residue 26 is highly varied and exposed. Changes in charge may affect binding at the surface defined by residue set #2; substitutions may affect the mobility of residue 27 that is in the principal set.

Residue 35 is most often Y, W has been observed.
20 The side group of 35 is buried, but substitution of F or W could affect the mobility of residue 34.

Residue 47 is always T or S in the sequence sample used. The O_{gamma} probably accepts a hydrogen bond from the NH of residue 50 in the alpha helix. Nevertheless,
25 there is no overwhelming steric reason to preclude other amino acid types at this residue. In particular, other amino acids the side groups of which can accept hydrogen bonds, viz. N, D, Q, and E, may be acceptable here.

Residue 50 is often an acidic amino acid, but other
30 amino acids are possible.

Residue 53 is often R, but other amino acids have been observed at this residue. Changes of charge may affect binding to the amino acids in interaction set #2.

106250" 5505550

Stereo Figure 14 shows the residues in set #2, plus R39. From Figure 14, one can see that R39 is on the opposite side of BPTI from the surface defined by the residues in set #2. Therefore, variation at residue 39 at the same time as variation of some residues in set #2 is much less likely to improve binding that occurs along surface #2 than is variation of the other residues in set #2.

In addition to the twelve principal residues and 13 secondary residues, there are two other residues, 30(C) and 33(F), involved in surface #2 that we will probably not vary, at least not until late in the procedure. These residues have their side groups buried inside BPTI and are conserved. Changing these residues does not change the surface nearly so much as does changing residues in the principal set. These buried, conserved residues do, however, contribute to the surface area of surface #2. The surface of residue set #2 is comparable to the area of the trypsin-binding surface. Principal residues 17, 19, 21, 27, 28, 29, 31, 32, 34, 48, 49, and 52 have a combined solvent-accessible area of 946.9 A². Secondary residues 9, 11, 15, 16, 18, 20, 22, 24, 26, 35, 47, 50, and 53 have combined surface of 1041.7 A². Residues 30 and 33 have exposed surface totaling 38.2 A². Thus the three groups' combined surface is 2026.8 A².

Residue 30 is C in BPTI and is conserved in all homologous sequences. It should be noted, however, that C14/C38 is conserved in all natural sequences, yet Marks et al. (MARK87) showed that changing both C14 and C38 to A,A or T,T yields a functional trypsin inhibitor. Thus it is possible that BPTI-like molecules will fold if C30 is replaced.

Residue 33 is F in BPTI and in all homologous sequences. Visual inspection of the BPTI structure

106290 55095850

suggests that substitution of Y, M, H, or L might be tolerated.

Having identified twenty residues that define a possible binding surface, we must choose some to vary first. Assuming a hypothetical affinity separation sensitivity, C_{sensi} , of 1 in $4 \cdot 10^8$, we decide to vary six residues (leaving some margin for error in the actual base composition of variegated bases). To obtain maximal recognition, we choose residues from the principal set that are as far apart as possible. Table 36 shows the distances between the β carbons of residues in the principal and peripheral set. R17 and V34 are at one end of the principal surface. Residues A27, G28, L29, A48, E49, and M52 are at the other end, about twenty Angstroms away; of these, we will vary residues 17, 27, 29, 34, and 48. Residues 28, 49, and 52 will be varied at later rounds.

Of the remaining principal residues, 21 is left to later variations. Among residues 19, 31, and 32, we arbitrarily pick 19 to vary.

Unlimited variation of six residues produces $6.4 \cdot 10^7$ amino acid sequences. By hypothesis, C_{sensi} is 1 in $4 \cdot 10^8$. Table 37 shows the programmed variegation at the chosen residues. The parental sequence is present as 1 part in $5.5 \cdot 10^7$, but the least favored sequences are present at only 1 part in $4.2 \cdot 10^9$. Among single-amino-acid substitutions from the PPBD, the least favored is F17-I19-A27-L29-V34-A48 and has a calculated abundance of 1 part in $1.6 \cdot 10^8$. Using the optimal qfk codon, we can recover the parental sequence and all one-amino-acid substitutions to the PPBD if actual nt compositions come within 5% of programmed compositions. The number of transformants is $M_{\text{ntv}} = 1.0 \cdot 10^9$ (also by hypothesis), thus we will produce most of the programmed sequences.

T05290-56096850

The residue numbers of the preceding section are referred to mature BPTI (R1-P2-...-A58). Table 25 has residue numbers referring to the pre-M13CP-BPTI protein; all mature BPTI sequence numbers have been increased by 5 the length of the signal sequence, i.e. 23. Thus in terms of the pre-OSP-PBD residue numbers, we wish to vary residues 40, 42, 50, 52, 57, and 71. A DNA subsequence containing all these codons is found between the (ApaI/-DraII/PssI) sites at base 191 and the Sph I site at base 10 309 of the osp-pbd gene. Among ApaI, DraI, and PssI, ApaI is preferred because it recognizes six bases without any ambiguity. DraII and PssI, on the other hand, recognize six bases with two-fold ambiguity at two of the bases. The vgDNA will contain more DraII and PssI 15 recognition sites at the varied locations than it will contain ApaI recognition sites. The unwanted extraneous cutting of the vgDNA by ApaI and SphI will eliminate a few sequences from our population. This is a minor problem, but by using the more specific enzyme (ApaI), we 20 minimize the unwanted effects. The sequence shown in Table 37 illustrates an additional way in which gratuitous restriction sites can be avoided in some cases. The osp-ipbd gene had the codon GGC for g51; because we are varying both residue 50 and 52, it is possible to obtain 25 an ApaI site. If we change the glycine codon to GGT, the ApaI site can no longer arise. ApaI recognizes the DNA sequence (GGGCC/C).

Each piece of dsDNA to be synthesized needs six to eight bases added at either end to allow cutting with 30 restriction enzymes and is shown in Table 37. The first synthetic base (before cutting with ApaI and SphI) is 184 and the last is 322. There are 142 bases to be synthesized. The center of the piece to the synthesized lies between Q54 and V57. The overlap can not include 35 varied bases, so we choose bases 245 to 256 as the

overlap that is 12 bases long. Note that the codon for F56 has been changed to TTC to increase the GC content of the overlap. The amino acids that are being varied are marked as X with a plus over them. Codons 57 and 71 are synthesized on the sense (bottom) strand. The design calls for "qfk" in the antisense strand, so that the sense strand contains (from 5' to 3') a) equal part C and A (i.e. the complement of k), b) (0.40 T, 0.22 A, 0.22 C, and 0.16 G) (i.e. the complement of f), and c) (0.26 T, 0.26 A, 0.30 C, and 0.18 G).

Each residue that is encoded by "qfk" has 21 possible outcomes, each of the amino acids plus stop. Table 12 gives the distribution of amino acids encoded by "qfk", assuming 5% errors. The abundance of the parental sequence is the product of the abundances of R x I x A x L x V x A. The abundance of the least-favored sequence is 1 in $4.2 \cdot 10^9$.

Olig#27 and olig#28 are annealed and extended with Klenow fragment and all four (nt)TPs. Both the ds synthetic DNA and RF pLG7 DNA are cut with both ApaI and SphI. The cut DNA is purified and the appropriate pieces ligated (See Sec. 14.1) and used to transform competent PE383. (Sec. 14.2). In order to generate a sufficient number of transformants, V_C is set to 5000 ml.

- 1) culture E. coli in 5.0 l of LB broth at 37°C until cell density reaches $5 \cdot 10^7$ to $7 \cdot 10^7$ cells/ml,
- 2) chill on ice for 65 minutes, centrifuge the cell suspension at 4000g for 5 minutes at 4°C,
- 3) discard supernatant; resuspend the cells in 1667 ml of an ice-cold, sterile solution of 60 mM CaCl_2 ,
- 4) chill on ice for 15 minutes, and then centrifuge at 4000g for 5 minutes at 4°C,
- 5) discard supernatant; resuspend cells in 2 x 400 ml of ice-cold, sterile 60 mM CaCl_2 ; store cells at 4°C

for 24 hours,

- 6) add DNA in ligation or TE buffer; mix and store on ice for 30 minutes; 20 ml of solution containing 5 μ g/ml of DNA is used,
- 5 7) heat shock cells at 42°C for 90 seconds,
- 8) add 200 ml LB broth and incubate at 37°C for 1 hour,
- 9) add the culture to 2.0 l of LB broth containing ampicillin at 35-100 μ g/ml and culture for 2 hours at 37°C,
- 10 10) centrifuge at 8000 g for 20 minutes at 4°C,
- 11) discard supernatant, resuspend cells in 50 ml of LB broth plus ampicillin and incubate 1 hour at 37°C,
- 12) plate cells on LB agar containing ampicillin,
- 13) harvest virions by method of Salivar et al. (SALI64).
- 15 The heat shock of step (7) can be done by dividing the 200 ml into 100 200 μ l aliquots in 1.5 ml plastic Eppendorf tubes. It is possible to optimize the heat shock for other volumes and kinds of container. It is important to:
 - a) use all or nearly all the vgDNA synthesized in ligation,
 - 20 b) use all or nearly all the ligation mixture to transform cells, and c) culture all or nearly all the transformants at high density. These measures are directed at maintaining diversity.
- 25 IPTG is added to the growth medium at 2.0 mM (the optimal level) and virions are harvested in the usual way. It is important to collect virions in a way that samples all or nearly all the transformants. Because F⁻ cells are used in the transformation, multiple infections do not
- 30 pose a problem.

HHMb has a pI of 7.0 and we carry out chromatography at pH 8.0 so that HHMb is slightly negative while BPTI and most of its mutants are positive. HHMb is fixed (Sec. V.F) to a 2.0 ml column on Affi-Gel 10^(TM) or Affi-Gel

15(TM) at 4.0 mg/ml support matrix, the same density that is optimal for a column supporting trp.

We note that charge repulsion between BPTI and HHMb should not be a serious problem and does not impose any constraints on ions or solutes allowed as eluants. Neither BPTI nor HHMb have special requirements that constrain choice of eluants. The eluant of choice is KCl in varying concentrations.

To remove variants of BPTI with strong, indiscriminate binding for any protein or for the support matrix, we pass the variegated population of virions over a column that supports bovine serum albumin (BSA) before loading the population onto the {HHMb} column. Affi-Gel 10(TM) or Affi-Gel 15(TM) is used to immobilize BSA at the highest level the matrix will support. A 10.0 ml column is loaded with 5.0 ml of Affi-Gel-linked-BSA; this column, called {BSA}, has $V_V = 5.0$ ml. The variegated population of virions containing 10^{12} pfu in 1 ml ($0.2 \times V_V$) of 10 mM KCl, 1 mM phosphate, pH 8.0 buffer is applied to {BSA}. We wash {BSA} with 4.5 ml ($0.9 \times V_V$) of 50 mM KCl, 1 mM phosphate, pH 8.0 buffer. The wash with 50 mM salt will elute virions that adhere slightly to BSA but not virions with strong binding. The pooled effluent of the {BSA} column is 5.5 ml of approximately 13 mM KCl.

The column {HHMb} is first blocked by treatment with 10^{11} virions of M13(am429) in 100 μ l of 10 mM KCl buffered to pH 8.0 with phosphate; the column is washed with the same buffer until OD_{260} returns to base line or $2 \times V_V$ have passed through the column, whichever comes first. The pooled effluent from {BSA} is added to {HHMb} in 5.5 ml of 13 mM KCl, 1 mM phosphate, pH 8.0 buffer. The column is eluted in the following way:

- 1) 10 mM KCl buffered to pH 8.0 with phosphate, until optical density at 280nm falls to base line or $2 \times$

F06290" 56096B60

- V_V , whichever is first, (effluent discarded),
- 2) a gradient of 10 mM to 2 M KCl in $3 \times V_V$, pH held at 8.0 with phosphate, (30·100 μ l fractions),
 - 3) a gradient of 2 M to 5 M KCl in $3 \times V_V$, phosphate buffer to pH 8.0 (30·100 μ l fractions),
 - 4) constant 5 M KCl plus 0 to 0.8 M guanidinium Cl in $2 \times V_V$, with phosphate buffer to pH 8.0, (20·100 μ l fractions), and
 - 5) constant 5 M KCl plus 0.8 M guanidinium Cl in $1 \times V_V$, with phosphate buffer to pH 8.0, (10·100 μ l fractions).

In addition to the elution fractions, a sample is removed from the column and used as an inoculum for phage-sensitive Sup⁻ cells (Sec. V). A sample of 4 μ l from each fraction is plated on phage-sensitive Sup⁻ cells. Fractions that yield too many colonies to count are replated at lower dilution. An approximate titre of each fraction is calculated. Starting with the last fraction and working toward the first fraction that was titered, we pool fractions until approximately 10^9 phage are in the pool, i.e. about 1 part in 1000 of the phage applied to the column. This population is infected into $3 \cdot 10^{11}$ phage-sensitive PE384 in 300 ml of LB broth. The very low multiplicity of infection (moi) is chosen to reduce the possibility of multiple infection. After thirty minutes, viable phage have entered recipient cells but have not yet begun to produce new phage. Phage-born genes are expressed at this phase, and we can add ampicillin that will kill uninfected cells. These cells still carry F-pili and will absorb phage helping to prevent multiple infections.

If multiple infection should pose a problem that cannot be solved by growth at low multiple-of-infection on F⁺ cells, the following procedure can be employed to obviate the problem. Virions obtained from the affinity

5

10

20

25

primarily to allow different charges (+, 0, -) to appear. Residue 38 is varied through K, R, E, or G. Residue 41 is varied through I, V, K, or E. Residue 43 is varied through R, S, G, N, K, D, E, T, or A.


5 Now assume that in the most successful SBD of the second round of variegation (SBD-2!), residue 38 (K15 of BPTI) changed to E, 41 becomes V, 43 goes to N, 44 goes to F, 51 goes to F, 54 goes to S, 55 goes to A, and 72 goes to Q (see Table 40). A third round of variation is
10 illustrated in Table 41; eight amino acids are varied. Those in the principal set, residues 40, 55, and 57, are varied through all twenty amino acids. Residue 32 is varied through P, Q, T, K, A, or E. Residue 34 is varied through T, P, Q, K, A, or E. Residue 44 is varied through
15 F, L, Y, C, W, or stop. Residue 50 is varied through E, K, or Q. Residue 52 is varied through L, F, I, M, or V. The result of this variation is shown in Table 42.

 This example is hypothetical. It is anticipated that more variegation cycles will be needed to achieve
20 dissociation constants of 10^{-8} M. It is also possible that more than three separation cycles will be needed in some variegation cycles. Real DNA chemistry and DNA synthesizers may have larger errors than our hypothetical
25 5%. If $S_{err} > 0.05$, then we may not be able to vary six residues at once. Variation of 5 residues at once is certainly possible.

EXAMPLE XII

DESIGN AND MUTAGENESIS OF A CLASS 1 MINI-PROTEIN

 To obtain a library of binding domains that are
30 conformationally constrained by a single disulfide, we insert DNA coding for the following family of mini-proteins into the gene coding for a suitable OSP.



5 Where indicates disulfide bonding; this mini-protein is depicted in Figure 3. Disulfides normally do not form between cysteines that are consecutive on the polypeptide chain. One or more of the residues indicated above as X_n will be varied extensively to obtain novel
10 binding. There may be one or more amino acids that precede X_1 or follow X_8 , however, these additional residues will not be significantly constrained by the diagrammed disulfide bridge, and it is less advantageous to vary these remote, unbridged residues. The last X
15 residue is connected to the OSP of the genetic package.

X_1 , X_2 , X_3 , X_4 , X_5 , X_6 , X_7 , and X_8 can be varied independently; i.e. a different scheme of variegation could be used at each position. X_1 and X_8 are the least constrained residues and may be varied less than other
20 positions.

X_1 and X_8 can be, for example, one of the amino acids [E, K, T, and A]; this set of amino acids is preferred because: a) the possibility of positively charged, negatively charged, and neutral amino acids is provided,
25 b) these amino acids can be provided in 1:1:1:1 ratio via the codon RMG (R = equimolar A and G, M = equimolar A and C), and c) these amino acids allow proper processing by signal peptidases.

One option for variegation of X_2 , X_3 , X_4 , X_5 , X_6 , and
30 X_7 is to vary all of these in the same way. For example, each of X_2 , X_3 , X_4 , X_5 , X_6 , and X_7 can be chosen from the set [F, S, Y, C, L, P, H, R, I, T, N, V, A, D, and G] which is encoded by the mixed codon NNT. Tables 10 and 130 compares libraries in which six codons have been
35 varied either by NNT or NNK codons. NNT encodes 15

T05290-5605650

different amino acids and only 16 DNA sequences. Thus, there are $1.139 \cdot 10^7$ amino-acid sequences, no stops, and only $1.678 \cdot 10^7$ DNA sequences. A library of 10^8 independent transformants will contain 99% of all possible sequences. The NNK library contains $6.4 \cdot 10^7$ sequences, but complete sampling requires a much larger number of independent transformants.

EXAMPLE XIII

A CYS::HELIX::TURN::STRAND::CYS UNIT

10 The parental Class 2 mini-proteins may be a naturally-occurring Class 2 mini-protein. It may also be a domain of a larger protein whose structure satisfies or may be modified so as to satisfy the criteria of a class 2 mini-protein. The modification may be a simple one, such
15 as the introduction of a cysteine (or a pair of cysteines) into the base of a hairpin structure so that the hairpin may be closed off with a disulfide bond, or a more elaborate one, so as the modification of intermediate residues so as to achieve the hairpin structure. The
20 parental class 2 mini-protein may also be a composite of structures from two or more naturally-occurring proteins, e.g., an α helix of one protein and a β strand of a second protein.

One mini-protein motif of potential use comprises a
25 disulfide loop enclosing a helix, a turn, and a return strand. Such a structure could be designed or it could be obtained from a protein of known 3D structure. Scorpion neurotoxin, variant 3, (ALMA83a, ALMA83b) (hereafter ScorpTx) contains a structure diagrammed in
30 Figure 15 that comprises a helix (residues N22 through N33), a turn (residues 33 through 35), and a return strand (residues 36 through 41). ScorpTx contains disulfides that join residues 12-65, 16-41, 25-46, and 29-48. CYS₂₅ and CYS₄₁ are quite close and could be joined by a

F05230"56096850

disulfide without deranging the main chain. Figure 15 shows CYS₂₅ joined to CYS₄₁. In addition, CYS₂₉ has been changed to GLN. It is expected that a disulfide will form between 25 and 41 and that the helix shown will form; we know that the amino-acid sequence shown is highly compatible with this structure. The presence of GLY₃₅, GLY₃₆, and GLY₃₉ give the turn and extended strand sufficient flexibility to accommodate any changes needed around CYS₄₁ to form the disulfide.

From examination of this structure (as found in entry 1SN3 of the Brookhaven Protein Data Bank), we see that the following sets of residues would be preferred for variegation:

SET 1

Residue	Codon	Allowed amino acids	Naa/Ndna
1) T ₂₇	NNG	L ² R ² MVSPTAQKEWG.	13/15
2) E ₂₈	VHG	LMVPTAGKE	9/9
3) A ₃₁	VHG	LMVPTAGKE	9/9
4) K ₃₂	VHG	LMVPTAGKE	9/9
5) G ₂₄	NNG	L ² R ² MVSPTAQKEWG.	13/15
6) E ₂₃	VHG	LMVPTAGKE	9/9
7) Q ₃₄	VAS	HONKED	6/6

Note: Exponents on amino acids indicate multiplicity of codons.

Positions 27, 28, 31, 32, 24, and 23 comprise one face of the helix. At each of these locations we have picked a variegating codon that a) includes the parental amino acid, b) includes a set of residues having a predominance of helix favoring residues, c) provides for a wide variety of amino acids, and d) leads to as even a distribution as possible. Position 34 is part of a turn. The side group of residue 34 could interact with molecules that contact the side groups of residues 27, 28, 31, 32, 24, and 23. Thus we allow variegation here and provide amino acids that are compatible with turns. The variega-

tion shown leads to $6.65 \cdot 10^6$ amino acid sequences encoded by $8.85 \cdot 10^6$ DNA sequences.

SET 2

	Residue	Codon	Allowed amino acids	Naa/Ndna
5	1) D ₂₆	VHS	L ² IMV ² P ² T ² A ² HQNKDE	13/18
	2) T ₂₇	NNG	L ² R ² MVSPTAQKEWG.	13/15
	3) K ₃₀	VHG	KEQPTALMV	9/9
	4) A ₃₁	VHG	KEQPTALMV	9/9
	5) K ₃₂	VHG	LMVPTAGKE	9/9
10	6) S ₃₇	RRT	SNDG	4/4
	7) Y ₃₈	NHT	YSFHPLNTIDAV	9/9


Positions 26, 27, 30, 31, and 32 are variegated so as to enhance helix-favoring amino acids in the population. Residues 37 and 38 are in the return strand so that we
 15 pick different variegation codons. This variegation allows $4.43 \cdot 10^6$ amino-acid sequences and $7.08 \cdot 10^6$ DNA sequences. Thus a library that embodies this scheme can be sampled very efficiently.

EXAMPLE XIV

20 DESIGN AND MUTAGENESIS OF CLASS 3 MINI-PROTEIN

Two Disulfide Bond Parental Mini-Proteins

Mini-proteins with two disulfide bonds may be modelled after the α -conotoxins, e.g., GI, GIA, GII, MI, and SI. These have the following conserved structure:



Hashimoto et al. (HASH85) reported synthesis of twenty-four analogues of α conotoxins GI, GII, and MI. Using the numbering scheme for GI (CYS at positions 2, 3, 7, and 13), Hashimoto et al. reported alterations at 4, 8,

10, and 12 that allows the proteins to be toxic. Almquist et al. (ALMQ89) synthesized [des-GLU₁] α Conotoxin GI and twenty analogues. They found that substituting GLY for PRO₅ gave rise to two isomers, perhaps related to different disulfide bonding. They found a number of substitutions at residues 8 through 11 that allowed the protein to be toxic. Zafaralla et al. (ZAFA88) found that substituting PRO at position 9 gives an active protein. Each of the groups cited used only in vivo toxicity as an assay for the activity. From such studies, one can infer that an active protein has the parental 3D structure, but one can not infer that an inactive protein lacks the parental 3D structure.

Pardi et al. (PARD89) determined the 3D structure of α Conotoxin GI obtained from venom by NMR. Kobayashi et al. (KOBA89) have reported a 3D structure of synthetic α Conotoxin GI from NMR data which agrees with that of PARD89. We refer to Figure 5 of Pardi et al..

Residue GLU₁ is known to accomodate GLU, ARG, and ILE in known analogues or homologues. A preferred variegation codon is NNG that allows the set of amino acids [L'R'MVSPTAQKEWG<stop>]. From Figure 5 of Pardi et al. we see that the side group of GLU₁ projects into the same region as the strand comprising residues 9 through 12. Residues 2 and 3 are cysteines and are not to be varied. The side group of residue 4 points away from residues 9 through 12; thus we defer varying this residue until a later round. PRO₅ may be needed to cause the correct disulfides to form; when GLY was substituted here the peptide folded into two forms, neither of which is toxic. It is allowed to vary PRO₅, but not preferred in the first round.

No substitutions at ALA₆ have been reported. A preferred variegation codon is RMG which gives rise to

FOIA b7 - D

ALA, THR, LYS, and GLU (small hydrophobic, small hydrophilic, positive, and negative). CYS₇ is not varied. We prefer to leave GLY₈ as is, although a homologous protein having ALA₈ is toxic. Homologous proteins having various amino acids at position 9 are toxic; thus, we use an NNT variegation codon which allows FS²YCLPHRITNVADG. We use NNT at positions 10, 11, and 12 as well. At position 14, following the fourth CYS, we allow ALA, THR, LYS, or GLU (via an RMG codon). This variegation allows $1.053 \cdot 10^7$ amino-acid sequences, encoded by $1.68 \cdot 10^7$ DNA sequences. Libraries having $2.0 \cdot 10^7$, $3.0 \cdot 10^7$, and $5.0 \cdot 10^7$ independent transformants will, respectively, display $\approx 70\%$, $\approx 83\%$, and $\approx 95\%$ of the allowed sequences. Other variegations are also appropriate. Concerning α conotoxins, see, inter alia, ALMQ89, CRUZ85, GRAY83, GRAY84, and PARD89.

The parental mini-protein may instead be one of the proteins designated "Hybrid-I" and "Hybrid-II" by Pease et al. (PEAS90); cf. Figure 4 of PEAS90. One preferred set of residues to vary for either protein consists of:

Parental Amino acid	Variegated Codon	Allowed Amino acids	AA seqs/ DNA seqs
A5	RVT	ADGTNS	6/6
P6	VYT	PTALIV	6/6
E7	RRS	EDNKS ² RG	7/8
T8	VHG	TPALMVQKE	9/9
A9	VHG	ATPLMVQKE	9/9
A10	RMG	AEKT	4/4
K12	VHG	KQETPALMV	9/9
Q16	NNG	L ² R ² S.WPOMTKVAEG	13/15

This provides $9.55 \cdot 10^6$ amino-acid sequences encoded by $1.26 \cdot 10^7$ DNA sequences. A library comprising $5.0 \cdot 10^7$ transformants allows expression of 98.2% of all possible sequences. At each position, the parental amino acid is allowed.

5


20

25


Three Disulfide Bond Parental Mini-Proteins

The cone snails (Conus) produce venoms (conotoxins) which are 10-30 amino acids in length and exceptionally rich in disulfide bonds. They are therefore archetypal

mini-proteins. Novel mini-proteins with three disulfide bonds may be modelled after the μ -(GIIIA, GIIIB, GIIIC) or Ω -(GVIA, GVIB, GVIC, GVIIA, GVIIIB, MVIIA, MVIIIB, etc.) conotoxins. The μ -conotoxins have the following conserved structure:




No 3D structure of a μ -conotoxin has been published. Hidaka et al. (HIDA90) have established the connectivity of the disulfides. The following diagram depicts geophutoxin I (also known as μ -conotoxin GIIIA).



The connection from R19 to C20 could go over or under the strand from Q14 to C15. One preferred form of variegation is to vary the residues in one loop. Because the longest loop contains only five amino acids, it is appropriate to also vary the residues connected to the cysteines that form the loop. For example, we might vary residues 5 through 9 plus 2, 11, 19, and 22. Another useful variegation would be to vary residues 11-14 and 16-19, each

through eight amino acids. Concerning μ conotoxins, see BECK89b, BECK89c, CRUZ89, and HIDA90.

The Ω -conotoxins may be represented as follows:



- 10 The King Kong peptide has the same disulfide arrangement as the Ω -conotoxins but a different biological activity. Woodward et al. (WOOD90) report the sequences of three homologous proteins from C. textile. Within the mature toxin domain, only the cysteines are conserved. The
- 15 spacing of the cysteines is exactly conserved, but no other position has the same amino acid in all three sequences and only a few positions show even pair-wise matches. Thus we conclude that all positions (except the
- 20 cysteines) may be substituted freely with a high probability that a stable disulfide structure will form. Concerning Ω conotoxins, see HILL89 and SUNX87.

- Another mini-protein which may be used as a parental binding domain is the Cucurbita maxima trypsin inhibitor I (CMTI-I); CMTI-III is also appropriate. They are members
- 25 of the squash family of serine protease inhibitors, which also includes inhibitors from summer squash, zucchini, and cucumbers (WIEC85). McWherter et al. (MCWH89) describe synthetic sequence-variants of the squash-seed protease inhibitors that have affinity for human leukocyte elastase
- 30 and cathepsin G. Of course, any member of this family might be used.

- CMTI-I is one of the smallest proteins known, comprising only 29 amino acids held in a fixed conformation by three disulfide bonds. The structure has been
- 35 studied by Bode and colleagues using both X-ray diffraction (BODE89) and NMR (HOLA89a,b). CMTI-I is of ellip-

T06290" 56096860

soidal shape; it lacks helices or β -sheets, but consists of turns and connecting short polypeptide stretches. The disulfide pairing is Cys3-Cys20, Cys10-Cys22 and Cys16--Cys28. In the CMTI-I:trypsin complex studied by Bode et al., 13 of the 29 inhibitor residues are in direct contact with trypsin; most of them are in the primary binding segment Val2(P4)-Glu9 (P4') which contains the reactive site bond Arg5(P1)-Ile6 and is in a conformation observed also for other serine proteinase inhibitors.

10 CMTI-I has a K_i for trypsin of $\approx 1.5 \cdot 10^{-12}$ M. McWherter et al. suggested substitution of "moderately bulky hydrophobic groups" at P1 to confer HLE specificity. They found that a wider set of residues (VAL, ILE, LEU, ALA, PHE, MET, and GLY) gave detectable binding to HLE.

15 For cathepsin G, they expected bulky (especially aromatic) side groups to be strongly preferred. They found that PHE, LEU, MET, and ALA were functional by their criteria; they did not test TRP, TYR, or HIS. (Note that ALA has the second smallest side group available.)

20 A preferred initial variegation strategy would be to vary some or all of the residues ARG₁, VAL₂, PRO₄, ARG₅, ILE₆, LEU₇, MET₈, GLU₉, LYS₁₁, HIS₂₅, GLY₂₆, TYR₂₇, and GLY₂₉. If the target were HNE, for example, one could synthesize DNA embodying the following possibilities:

25		vg	Allowed	#AA seqs/
	<u>Parental</u>	<u>Codon</u>	<u>amino acids</u>	<u>#DNA seqs</u>
	ARG ₁	VNT	RSLPHITNVADG	12/12
	VAL ₂	NWT	VILFYHND	8/8
	PRO ₄	VYT	PLTIAV	6/6
30	ARG ₅	VNT	RSLPHITNVADG	12/12
	ILE ₆	NNK	all 20	20/31
	LEU ₇	VWG	LQMKVE	6/6
	<u>TYR₂₇</u>	<u>NAS</u>	<u>YHONKDE.</u>	<u>7/8</u>

This allows about $5.81 \cdot 10^6$ amino-acid sequences encoded by

35 about $1.03 \cdot 10^7$ DNA sequences. A library comprising

$5.0 \cdot 10^7$ independent transformants would give $\approx 99\%$ of the possible sequences. Other variegation schemes could also be used.

Other inhibitors of this family include:

- 5 Trypsin inhibitor I from Citrullus vulgaris (OTLE87),
Trypsin inhibitor II from Bryonia dioica (OTLE87),
Trypsin inhibitor I from Cucurbita maxima (in OTLE87),
trypsin inhibitor III from Cucurbita maxima (in OTLE87),
trypsin inhibitor IV from Cucurbita maxima (in OTLE87),
- 10 trypsin inhibitor II from Cucurbita pepo (in OTLE87),
trypsin inhibitor III from Cucurbita pepo (in OTLE87),
trypsin inhibitor IIb from Cucumis sativus (in OTLE87),
trypsin inhibitor IV from Cucumis sativus (in OTLE87),
trypsin inhibitor II from Ecballium elaterium (FAVE89),
- 15 and inhibitor CM-1 from Momordica repens (in OTLE87).

- Another mini-protein that may be used as an initial potential binding domain is the heat-stable enterotoxins derived from some enterotoxigenic E. coli, Citrobacter
- 20 freundii, and other bacteria (GUAR89). These mini-proteins are known to be secreted from E. coli and are extremely stable. Works related to synthesis, cloning, expression and properties of these proteins include: BHAT86, SEKI85, SHIM87, TAKA85, TAKE90, THOM85a,b, YOSH85,
 - 25 DALL90, DWAR89, GARI87, GUZM89, GUZM90, HOUG84, KUBO89, KUPÉ90, OKAM87, OKAM88, and OKAM90.

- Another preferred IPBD is crambin or one of its homologues, the phoratoxins and ligatoxins (LECO87). These proteins are secreted in plants. The 3D structure
- 30 of crambin has been determined. NMR data on homologues indicate that the 3D structure is conserved. Residues thought to be on the surface of crambin, phoratoxin, or ligatoxin are preferred residues to vary.


EXAMPLE XV

A MINI-PROTEIN HAVING A CROSS-LINK CONSISTING OF CU(II),
ONE CYSTEINE, TWO HISTIDINES, AND ONE METHIONINE.


Sequences such as

- 5 HIS-ASN-GLY-MET-Xaa-Xaa-Xaa-Xaa-Xaa-Xaa-HIS-ASN-GLY-CYS
and
CYS-ASN-GLY-MET-Xaa-Xaa-Xaa-Xaa-Xaa-Xaa-HIS-ASN-GLY-HIS
are likely to combine with Cu(II) to form structures as
shown in the diagram:

10



Other arrangements of HIS, MET, HIS, and CYS along the
chain are also likely to form similar structures. The
30 amino acids ASN-GLY at positions 2 and 3 and at positions
12 and 13 give the amino acids that carry the metal-
binding ligands enough flexibility for them to come
together and bind the metal. Other connecting sequences
may be used, e.g. GLY-ASN, SER-GLY, GLY-PRO, GLY-PRO-GLY,
35 or PRO-GLY-ASN could be used. It is also possible to vary
one or more residues in the loops that join the first and
second or the third and fourth metal-binding residues.
For example,



is likely to form the diagrammed structure for a wide variety of amino acids at Xaa4. It is expected that the side groups of Xaa4 and Xaa6 will be close together and on the surface of the mini-protein.

The variable amino acids are held so that they have limited flexibility. This cross-linkage has some differences from the disulfide linkage. The separation between $C_{\alpha 4}$ and $C_{\alpha 11}$ is greater than the separation of the C_{α} s of a cystine. In addition, the interaction of residues 1 through 4 and 11 through 14 with the metal ion are expected to limit the motion of residues 5 through 10 more than a disulfide between residues 4 and 11. A single disulfide bond exerts strong distance constraints on the α carbons of the joined residues, but very little directional constraint on, for example, the vector from N to C in the main-chain.

For the desired sequence, the side groups of residues 5 through 10 can form specific interactions with the target. Other numbers of variable amino acids, for example, 4, 5, 7, or 3, are appropriate. Larger spans may be used when the enclosed sequence contains segments having a high potential to form α helices or other secondary structure that limits the conformational freedom

of the polypeptide main chain. Whereas a mini-protein having four CYSSs could form three distinct pairings, a mini-protein having two HISSs, one MET, and one CYS can form only two distinct complexes with Cu. These two
 5 structures are related by mirror symmetry through the Cu. Because the two HISSs are distinguishable, the structures are different.

When such metal-containing mini-proteins are displayed on filamentous phage, the cells that produce the
 10 phage can be grown in the presence of the appropriate metal ion, or the phage can be exposed to the metal only after they are separated from the cells.

EXAMPLE XVI

A MINI-PROTEIN HAVING A CROSS-LINK CONSISTING OF ZN(II)
 15 AND FOUR CYSTEINES

A cross link similar to the one shown in Example XV is exemplified by the Zinc-finger proteins (GIBS88, GAUS87, PARR88, FRAN87, CHOW87, HARD90). One family of Zinc-fingers has two CYS and two HIS residues in conserved
 20 positions that bind Zn_{++} (PARR88, FRAN87, CHOW87, EVAN88, BERG88, CHAV88). Gibson *et al.* (GIBS88) review a number of sequences thought to form zinc-fingers and propose a three-dimensional model for these compounds. Most of these sequences have two CYS and two HIS residues in
 25 conserved positions, but some have three CYS and one HIS residue. Gauss *et al.* (GAUS87) also report a zinc-finger protein having three CYS and one HIS residues that bind zinc. Hard *et al.* (HARD90) report the 3D structure of a protein that comprises two zinc-fingers, each of which has
 30 four CYS residues. All of these zinc-binding proteins are stable in the reducing intracellular environment.

One preferred example of a CYS::zinc cross linked mini-protein comprises residues 440 to 461 of the sequence

09896095-052904

shown in Figure 1 of HARD90. The residues 444 through 456 may be variegated. One such variegation is as follows:

	Parental	Allowed	#AA / #DNA
	SER444	SER, ALA	2 / 2
5	ASP445	ASP, ASN, GLU, LYS	4 / 4
	GLU446	GLU, LYS, GLN	3 / 3
	ALA447	ALA, THR, GLY, SER	4 / 4
	SER448	SER, ALA	2 / 2
	GLY449	GLY, SER, ASN, ASP	4 / 4
10	CYS450	CYS, PHE, ARG, LEU	4 / 4
	HIS451	HIS, GLN, ASN, LYS, ASP, GLU	6 / 6
	TYR452	TYR, PHE, HIS, LEU	4 / 4
	GLY453	GLY, SER, ASN, ASP	4 / 4
	VAL454	VAL, ALA, ASP, GLY, SER, ASN, THR, ILE	8 / 8
15	LEU455	LEU, HIS, ASP, VAL	4 / 4
	THR456	THR, ILE, ASN, SER	4 / 4

This leads to $3.77 \cdot 10^7$ DNA sequences that encode the same number of amino-acid sequences. A library having $1.0 \cdot 10^8$

- 20 independent transformants will display 93% of the allowed sequences; $2.0 \cdot 10^8$ independent transformants will display 99.5% of allowed sequences.

T05250" 56036360


Unsuccessful Authorization

Processing Center Response

DECLINE

Local Response

REFER TO FIELD #1004



FORM 5605

Table 1: Single-letter codes.

5	<u>Single-letter code is used for proteins :</u>				
	a = ALA	c = CYS	d = ASP	e = GLU	f = PHE
	g = GLY	h = HIS	i = ILE	k = LYS	l = LEU
	m = MET	n = ASN	p = PRO	q = GLN	r = ARG
	s = SER	t = THR	v = VAL	w = TRP	y = TYR
10	. = STOP * = any amino acid				
	b = n or d				
	z = e or q				
	x = any amino acid				
15					
	<u>Single-letter IUB codes for DNA :</u>				
20	T, C, A, G stand for themselves				
	M for A or C				
	R for puRines A or G				
25	W for A or T				
	S for C or G				
	Y for pYrimidines T or C				
	K for G or T				
30	V for A, C, or G (not T)				
	H for A, C, or T (not G)				
	D for A, G, or T (not C)				
	B for C, G, or T (not A)				
35	N for any base.				

T06250-56056860

Table 2: Preferred Outer-Surface Proteins

	Genetic Package	Preferred Outer-Surface Protein	Reason for preference
5	M13	coat protein (gpVIII)	a) exposed amino terminus, b) predictable post-translational processing, c) numerous copies in virion. d) fusion data available
10		gp III	a) fusion data available. b) amino terminus exposed. c) working example available.
15	PhiX174	G protein	a) known to be on virion exterior, b) small enough that the <u>G-ipbd</u> gene can replace H gene.
20	<u>E. coli</u>	LamB	a) fusion data available, b) non-essential.
25		OmpC	a) topological model b) non-essential; abundant
30		OmpA	a) topological model b) non-essential; abundant c) homologues in other genera
35		OmpF	a) topological model b) non-essential; abundant
40		PhoE	a) topological model b) non-essential; abundant c) inducible
45	<u>B. subtilis</u> spores	CotC	a) no post-translational processing, b) distinctive ssequence that causes protein to localize in spore coat, c) non-essential.
		CotD	Same as for CotC.

Table 3: Ambiguous DNA for AA_seq2

5	m 1 A.T.G	k 2 A.A.r	k 3 A.A.r	s 4 T.C.n A.G.y	l 5 T.T.r C.T.n	v 6 G.T.n	l 7 T.T.r C.T.n	k 8 A.A.r
10	a 9 G.C.n	s 10 T.C.n A.G.y	v 11 G.T.n	a 12 G.C.n	v 13 G.T.n	a 14 G.C.n	t 15 A.C.n	l 16 T.T.r C.T.n
15	v 17 G.T.n	p 18 C.C.n	m 19 A.T.G	l 20 T.T.r C.T.n	s 21 T.C.n A.G.y	f 22 T.T.y	a 23 G.C.n	r 24 C.G.n A.G.r
20	p 25 C.C.n	d 26 G.A.y	f 27 T.T.y	c 28 T.G.y	l 29 T.T.r C.T.n	e 30 G.A.r	p 31 C.C.n	p 32 C.C.n
25	y 33 T.A.y	t 34 A.C.n	g 35 G.G.n	p 36 C.C.n	c 37 T.G.y	k 38 A.A.r	a 39 G.C.n	r 40 C.G.n A.G.r
30	i 41 A.T.h	i 42 A.T.h	r 43 C.G.n	y 44 T.A.y	f 45 T.T.y	y 46 T.A.y	n 47 A.A.y	a 48 G.C.n
35	k 49 A.A.r	a 50 G.C.n	g 51 G.G.n	l 52 T.T.r C.T.n	c 53 T.G.y	q 54 C.A.r	t 55 A.C.n	f 56 T.T.y
40	v 57 G.T.n	y 58 T.A.y	g 59 G.G.n	g 60 G.G.n	c 61 T.G.y	r 62 C.G.n A.G.r	a 63 G.C.n	k 64 A.A.r
45								

T05230-56056860

Table 3, continued.

5	r 65 C.G.n A.G.r	n 66 A.A.y	n 67 A.A.y	f 68 T.T.y	k 69 A.A.r	s 70 T.C.n A.G.y	a 71 G.C.n	e 72 G.A.r
10	d 73 G.A.y	c 74 T.G.y	m 75 A.T.G	r 76 C.G.n	t 77 A.C.n	c 78 T.G.y	g 79 G.G.n	g 80 G.G.n
15	a 81 G.C.n	a 82 G.C.n	e 83 G.A.r	g 84 G.G.n	d 85 G.A.y	d 86 G.A.y	p 87 C.C.n	a 88 G.C.n
20	k 89 A.A.r	a 90 G.C.n	a 91 G.C.n	f 92 T.T.y	N 93 A.A.y	s 94 T.C.n A.G.y	l 95 T.T.r C.T.n	q 96 C.A.r
25	a 97 G.C.n	s 98 T.C.n A.G.y	a 99 G.C.n	t 100 A.C.n	e 101 G.A.r	y 102 T.A.y	i 103 A.T.h	g 104 G.G.n
30	y 105 T.A.y	a 106 G.C.n	w 107 T.G.G	a 108 G.C.n	m 109 A.T.G	v 110 G.T.n	v 111 G.T.n	v 112 G.T.n
35	i 113 A.T.h	v 114 G.T.n	g 115 G.G.n	a 116 G.C.n	t 117 A.C.n	i 118 A.T.h	g 119 G.G.n	i 120 A.T.h
40	k 121 A.A.r	l 122 T.T.r C.T.n	f 123 T.T.y	k 124 A.A.r	k 125 A.A.r	f 126 T.T.y	t 127 A.C.n	s 128 T.C.n A.G.y
45	k 129 A.A.r	a 130 G.C.n	s 131 T.C.n A.G.y	.132 T.A.r T.G.A	.133 T.A.r T.G.A	.134 T.A.r T.G.A		
50								

FO62220"56096860

Table 4: Table of Restriction Enzyme Suppliers

5

Suppliers :

10

Sigma Chemical Co.
P.O.Box 14508
St. Louis, Mo. 63178

15

Bethesda Research Laboratories
P.O.Box 6009
Gaithersburg, Maryland, 20877

20

Boehringer Mannheim Biochemicals
7941 Castleway Drive
Indianapolis, Indiana, 46250

25

International Biochemicals, Inc.
P.O.Box 9558
New Haven, Connecticut, 06535

30

New England BioLabs
32 Tozer Road
Beverly, Massachusetts, 01915

Promega
2800 S. Fish Hatchery Road
Madison, Wisconsin, 53711

Stratagene Cloning Systems
11099 North Torrey Pines Road
La Jolla, California, 92037

T06230" 56035350

Table 5: Potential sites in ipbd gene.

Summary of cuts.

5	Enz = % <u>Acc</u> I has 3 elective sites : 96 169 281
	Enz = <u>Afl</u> II has 1 elective sites : 19
	Enz = <u>Apa</u> I has 2 elective sites : 102 103
	Enz = <u>Asu</u> II has 1 elective sites : 381
	Enz = <u>Ava</u> III has 1 elective sites : 314
10	Enz = <u>BspM</u> II has 1 elective sites : 72
	Enz = <u>BssH</u> II has 2 elective sites : 67 115
	Enz = % <u>BstX</u> I has 1 elective sites : 323
	Enz = + <u>Dra</u> II has 3 elective sites : 102 103 226
	Enz = + <u>EcoN</u> I has 2 elective sites : 62 94
15	Enz = + <u>Esp</u> I has 2 elective sites : 57 187
	Enz = <u>Hind</u> III has 6 elective sites : 9 23 60
	287 361 386
	Enz = <u>Kpn</u> I has 1 elective sites : 48
	Enz = <u>Mlu</u> I has 1 elective sites : 314
20	Enz = <u>Nar</u> I has 2 elective sites : 238 343
	Enz = <u>Nco</u> I has 1 elective sites : 323
	Enz = <u>Nhe</u> I has 3 elective sites : 25 289 388
	Enz = <u>Nru</u> I has 2 elective sites : 38 65
	Enz = + <u>PflM</u> I has 1 elective sites : 94
25	Enz = <u>PmaC</u> I has 1 elective sites : 228
	Enz = + <u>PpuM</u> I has 2 elective sites : 102 226
	Enz = + <u>Rsr</u> II has 1 elective sites : 102
	Enz = + <u>Sfi</u> I has 2 elective sites : 24 261
	Enz = <u>Spe</u> I has 3 elective sites : 12 45 379
30	Enz = <u>Sph</u> I has 1 elective sites : 221
	Enz = <u>Stu</u> I has 5 elective sites : 23 70 150
	287 386
	Enz = % <u>Sty</u> I has 6 elective sites : 11 44
	143 263 323 383
35	Enz = <u>Xba</u> I has 1 elective sites : 84
	Enz = <u>Xho</u> I has 1 elective sites : 85
	Enz = <u>Xma</u> III has 3 elective sites : 70 209
	242

40 Enzymes not cutting ipbd.

	<u>Avr</u> II	<u>BamH</u> I	<u>Bcl</u> I	<u>BstE</u> II
	<u>EcoR</u> I	<u>EcoR</u> V	<u>Hpa</u> I	<u>Not</u> I
	<u>Sac</u> I	<u>Sal</u> I	<u>Sau</u> I	<u>Sma</u> I
45	<u>Xma</u> I			

5

10

5

10

20

25 "most" is the value of n having the highest probability.

Table 9: best vgCodon

```

5      Program "Find Optimum vgCodon."
      INITIALIZE-MEMORY-OF-ABUNDANCES
      DO ( t1 = 0.21 to 0.31 in steps of 0.01 )
        . DO ( c1 = 0.13 to 0.23 in steps of 0.01 )
          . . DO ( a1 = 0.23 to 0.33 in steps of 0.01 )
10     Comment      calculate g1 from other concentrations
          . . . g1 = 1.0 - t1 - c1 - a1
          . . . IF( g1 .ge. 0.15 )
          . . . . DO ( a2 = 0.37 to 0.50 in steps of 0.01 )
          . . . . . DO ( c2 = 0.12 to 0.20 in steps of 0.01 )
15     Comment      Force D+E = R + K
          . . . . . g2 = (g1*a2 -.5*a1*a2)/(c1+0.5*a1)
      Comment      Calc t2 from other concentrations.
          . . . . . t2 = 1. - a2 - c2 - g2
          . . . . . IF(g2.gt. 0.1.and. t2.gt.0.1)
20     . . . . . CALCULATE-ABUNDANCES
          . . . . . COMPARE-ABUNDANCES-TO-PREVIOUS-ONES
          . . . . . ..end_IF_block
          . . . . . ..end_DO_loop ! c2
          . . . . . ..end_DO_loop ! a2
25     . . . ..end_IF_block ! if g1 big enough
          . . . ..end_DO_loop ! a1
          . . . ..end_DO_loop ! c1
          ..end_DO_loop ! t1
      WRITE the best distribution and the abundances.

```

T06230' 56095360

Table 10: Abundances obtained
from various vgCodons

A. Optimized fxS Codon, Restrained by $[D]+[E] = [K]+[R]$

5

	T	C	A	G	
1	.26	.18	.26	.30	f
2	.22	.16	.40	.22	x
3	.5	.0	.0	.5	S

10

Amino acid	Abundance	Amino acid	Abundance
A	4.80%	C	2.86%
D	6.00%	E	6.00%
F	2.86%	G	6.60%
H	3.60%	I	2.86%
K	5.20%	L	6.82%
M	2.86%	N	5.20%
P	2.88%	Q	3.60%
R	6.82%	S	7.02% mfaa
T	4.16%	V	6.60%
W	2.86% lfaa	Y	5.20%
stop	5.20%		

25

$$[D] + [E] = [K] + [R] = .12$$

$$\text{ratio} = \text{Abun}(W)/\text{Abun}(S) = 0.4074$$

30

i	$(1/\text{ratio})^j$	$(\text{ratio})^j$	stop-free
1	2.454	.4074	.9480
2	6.025	.1660	.8987
3	14.788	.0676	.8520
4	36.298	.0275	.8077
5	89.095	.0112	.7657
6	218.7	$4.57 \cdot 10^{-3}$.7258
7	536.8	$1.86 \cdot 10^{-3}$.6881

35

0095095-06601

Table 10: Abundances obtained
from various vgCodon
(continued)

5 B. Unrestrained, optimized

	T	C	A	G
1	.27	.19	.27	.27
2	.21	.15	.43	.21
10 3	.5	.0	.0	.5

Amino acid	Abundance	Amino acid	Abundance
A	4.05%	C	2.84%
D	5.81%	E	5.81%
F	2.84%	G	5.67%
H	4.08%	I	2.84%
K	5.81%	L	6.83%
M	2.84%	N	5.81%
15 P	2.85%	Q	4.08%
R	6.83%	<u>S</u>	6.89% mfaa
20 T	4.05%	V	5.67%
W	2.84% lfaa	Y	5.81%
<u>stop</u>	5.81%		

25 [D] + [E] = 0.1162 [K] + [R] = 0.1264

ratio = Abun(W)/Abun(S) = 0.41176

i	$(1/\text{ratio})^j$	$(\text{ratio})^j$	<u>stop-free</u>
1	2.4286	.41176	.9419
2	5.8981	.16955	.8872
35 3	14.3241	.06981	.8356
4	34.7875	.02875	.7871
5	84.4849	.011836	.74135
6	205.180	.004874	.69828
40 7	498.3	$2.007 \cdot 10^{-3}$.6577

T06E50"56096850

Table 10: Abundances obtained
from various vgCodon
(continued)

5 C. Optimized NNT

	T	C	A	G
1	.2071	.2929	.2071	.2929
2	.2929	.2071	.2929	.2071
10 3	1.	.0	.0	.0

	Amino acid	Abundance	Amino acid	Abundance
15	A	6.06%	C	4.29% lfaa
	D	8.58%	E	none
	F	6.06%	G	6.06%
	H	8.58%	I	6.06%
	K	none	L	8.58%
20	M	none	N	6.06%
	P	6.06%	Q	none
	R	6.06%	S	8.58% mfaa
	T	4.29% lfaa	V	8.58%
	W	none	Y	6.06%
25	stop	none		

	i	$(1/\text{ratio})^j$	$(\text{ratio})^j$	stop-free
	1	2.0	.5	1.
30	2	4.0	.25	1.
	3	8.0	.125	1.
	4	16.0	.0625	1.
	5	32.0	.03125	1.
	6	64.0	.015625	1.
35	7	128.0	.0078125	1.

005250" 52056860

Table 10: Abundances obtained
from various vgCodon
(continued)

5

D. Optimized NNG

	T	C	A	G
1	.23	.21	.23	.33
2	.215	.285	.285	.215
3	.0	.0	.0	1.0

	Amino acid	Abundance	Amino acid	Abundance
15	A	9.40%	C	none
	D	none	E	9.40%
	F	none	G	7.10%
	H	none	I	none
20	K	6.60%	L	9.50% mfaa
	M	4.90%	N	none
	P	6.00%	Q	6.00%
	R	9.50%	S	6.60%
	T	6.6 %	V	7.10%
25	W	4.90% lfaa	Y	none
	stop	6.60%		

	<u>i</u>	<u>(1/ratio)^j</u>	<u>(ratio)^j</u>	<u>stop-free</u>
30	1	1.9388	.51579	0.934
	2	3.7588	.26604	0.8723
	3	7.2876	.13722	0.8148
	4	14.1289	.07078	0.7610
	5	27.3929	$3.65 \cdot 10^{-2}$	0.7108
35	6	53.109	$1.88 \cdot 10^{-2}$	0.6639
	7	102.96	$9.72 \cdot 10^{-3}$	0.6200

FO5290"56096360

5

10

1530

<u>i</u>	<u>(1/ratio)^j</u>	<u>(ratio)^j</u>	<u>stop-free</u>
1	3.0	.33333	.96875
2	9.0	.11111	.9385
3	27.0	.03704	.90915
4	81.0	.01234567	.8807
5	243.0	.0041152	.8532
6	729.0	$1.37 \cdot 10^{-3}$.82655
7	2187.0	$4.57 \cdot 10^{-4}$.8007

Table 11: Calculate worst codon.

```

5      Program "Find worst vgCodon within Serr of given
      distribution."
      INITIALIZE-MEMORY-OF-ABUNDANCES
      Comment Serr is % error level.
      READ Serr
      Comment T1i,C1i,A1i,G1i, T2i,C2i,A2i,G2i, T3i,G3i
10     Comment are the intended nt-distribution.
      READ T1i, C1i, A1i, G1i
      READ T2i, C2i, A2i, G2i
      READ T3i, G3i
      Fdwn = 1.-Serr
15     Fup = 1.+Serr
      DO ( t1 = T1i*Fdwn to T1i*Fup in 7 steps)
      . DO ( c1 = C1i*Fdwn to C1i*Fup in 7 steps)
      . . DO ( a1 = A1i*Fdwn to A1i*Fup in 7 steps)
      . . . g1 = 1. - t1 - c1 - a1
20     . . . IF( (g1-G1i)/G1i .lt. -Serr)
      Comment g1 too far below G1i, push it back
      . . . . g1 = G1i*Fdwn
      . . . . factor = (1.-g1)/(t1 + c1 + a1)
      . . . . t1 = t1*factor
25     . . . . c1 = c1*factor
      . . . . a1 = a1*factor
      . . . . ..end_IF_block
      . . . IF( (g1-G1i)/G1i .gt. Serr)
      Comment g1 too far above G1i, push it back
30     . . . . g1 = G1i*Fup
      . . . . factor = (1.-g1)/(t1 + c1 + a1)
      . . . . t1 = t1*factor
      . . . . c1 = c1*factor
      . . . . a1 = a1*factor
35     . . . . ..end_IF_block
      . . . DO ( a2 = A2i*Fdwn to A2i*Fup in 7 steps)
      . . . . DO ( c2 = C2i*Fdwn to C2i*Fup in 7 steps)
      . . . . DO ( g2=G2i*Fdwn to G2i*Fup in 7 steps)
      Comment Calc t2 from other concentrations.
40     . . . . . t2 = 1. - a2 - c2 - g2
      . . . . . IF( (t2-T2i)/T2i .lt. -Serr)
      Comment t2 too far below T2i, push it back
      . . . . . . t2 = T2i*Fdwn
      . . . . . . factor = (1.-t2)/(a2 + c2 + g2)
45     . . . . . . a2 = a2*factor
      . . . . . . c2 = c2*factor
      . . . . . . g2 = g2*factor
      . . . . . . ..end_IF_block
      . . . . . IF( (t2-T2i)/T2i .gt. Serr)
50     Comment t2 too far above T2i, push it back
      . . . . . . t2 = T2i*Fup
      . . . . . . factor = (1.-t2)/(a2 + c2 + g2)

```

Table 11, continued.

```

. . . . . a2 = a2*factor
. . . . . c2 = c2*factor
5 . . . . . g2 = g2*factor
. . . . . ..end_IF_block
. . . . . IF(g2.gt. 0.0 .and. t2.gt.0.0)
. . . . . t3 = 0.5*(1.-Serr)
. . . . . g3 = 1. - t3
10 . . . . . CALCULATE-ABUNDANCES
. . . . . COMPARE-ABUNDANCES-TO-PREVIOUS-ONES
. . . . . t3 = 0.5
. . . . . g3 = 1. - t3
. . . . . CALCULATE-ABUNDANCES
15 . . . . . COMPARE-ABUNDANCES-TO-PREVIOUS-ONES
. . . . . t3 = 0.5*(1.+Serr)
. . . . . g3 = 1. - t3
. . . . . CALCULATE-ABUNDANCES
. . . . . COMPARE-ABUNDANCES-TO-PREVIOUS-ONES
20 . . . . . ..end_IF_block
. . . . . ..end_DO_loop ! g2
. . . . . ..end_DO_loop ! c2
. . . . . ..end_DO_loop ! a2
. . . . . ..end_DO_loop ! a1
25 . . . . . ..end_DO_loop ! c1
. . . . . ..end_DO_loop ! t1
WRITE the WORST distribution and the abundances.

```

T06230-5656860

Table 12: Abundances obtained
using optimum vgCodon assuming
5% errors

Amino acid	Abundance	Amino acid	Abundance
A	4.59%	C	2.76%
D	5.45%	E	6.02%
F	2.49% lfaa	G	6.63%
H	3.59%	I	2.71%
K	5.73%	L	6.71%
M	3.00%	N	5.19%
P	3.02%	Q	3.97%
R	7.68% mfaa	S	7.01%
T	4.37%	V	6.00%
W	3.05%	Y	4.77%
stop	5.27%		

$$\text{ratio} = \text{Abun(F)}/\text{Abun(R)} = 0.3248$$

<u>j</u>	<u>(1/ratio)^j</u>	<u>(ratio)^j</u>	<u>stop-free</u>
1	3.079	.3248	.9473
2	9.481	.1055	.8973
3	29.193	.03425	.8500
4	89.888	.01112	.8052
5	276.78	$3.61 \cdot 10^{-3}$.7627
6	852.22	$1.17 \cdot 10^{-3}$.7225
7	2624.1	$3.81 \cdot 10^{-4}$.6844

T06290-59096850

Table 13: BPTI Homologues

R #	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19
-3	-	-	-	F	-	-	-	-	-	-	-	-	-	-	-	-	Z	-	-
-2	-	-	-	Q	T	-	-	-	-	-	-	Q	-	-	-	H	G	Z	-
-1	-	-	-	T	E	-	-	-	-	-	-	P	-	-	-	D	D	G	-
1	R	R	R	P	R	R	R	R	R	R	R	L	A	R	R	R	K	R	A
2	P	P	P	P	P	P	P	P	P	P	P	R	A	P	P	P	R	P	A
3	D	D	D	D	D	D	D	D	D	D	D	K	K	D	R	T	D	S	K
4	F	F	F	L	F	F	F	F	F	F	F	L	Y	F	F	F	I	F	Y
5	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C
6	L	L	L	Q	L	L	L	L	L	L	L	I	K	E	E	N	R	N	K
7	E	E	E	L	E	E	E	E	E	E	E	L	L	L	L	L	L	L	L
8	P	P	P	P	P	P	P	P	P	P	P	H	P	P	P	P	P	P	P
9	P	P	P	Q	P	P	P	P	P	P	P	R	L	A	A	P	P	A	V
10	Y	Y	Y	A	Y	Y	Y	Y	Y	Y	Y	N	R	E	E	E	E	E	R
11	T	T	T	R	T	T	T	T	T	T	T	P	I	T	T	S	Q	T	Y
12	G	G	G	G	G	G	G	G	G	G	G	G	G	G	G	G	G	G	G
13	P	P	P	P	P	P	P	P	P	P	P	R	P	L	L	R	P	P	P
14	C	T	A	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C
15	K	K	K	K	K	V	G	A	L	I	K	Y	K	K	K	R	K	K	K
16	A	A	A	A	A	A	A	A	A	A	A	Q	R	A	A	G	G	A	K
17	R	R	R	A	A	R	R	R	R	R	R	K	K	Y	R	H	R	S	K
18	I	I	I	L	M	I	I	I	I	I	I	I	I	I	I	I	L	I	F
19	I	I	I	L	I	I	I	I	I	I	I	P	P	S	R	R	P	R	P
20	R	R	R	R	R	R	R	R	R	R	R	A	S	S	S	R	R	Q	S
21	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	F	F	F	F	I	Y	Y	F
22	F	F	F	F	F	F	F	F	F	F	F	Y	Y	H	H	Y	F	Y	Y
23	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
24	N	N	N	N	N	N	N	N	N	N	N	N	K	N	N	N	N	N	N
25	A	A	A	S	A	A	A	A	A	A	A	Q	W	L	R	L	P	S	W
26	K	K	K	T	K	K	K	K	K	K	K	K	K	A	A	E	A	K	K
27	A	A	A	S	A	A	A	A	A	A	A	K	A	A	A	S	S	S	A
28	G	G	G	N	G	G	G	G	G	G	G	K	K	Q	Q	N	R	G	K
29	L	L	L	A	F	L	L	L	L	L	L	Q	Q	Q	Q	K	M	G	Q
30	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C
31	Q	Q	Q	E	E	Q	Q	Q	Q	Q	Q	E	L	L	L	K	E	Q	L
32	T	T	T	P	T	T	T	T	T	T	T	G	P	Q	E	V	S	Q	P
33	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F
34	V	V	V	T	V	V	V	V	V	V	V	T	D	I	I	F	I	I	N
35	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	W	Y	Y	Y	Y	Y	Y	Y
36	G	G	G	G	G	G	G	G	G	G	G	S	S	G	G	G	G	G	S
37	G	G	G	G	G	G	G	G	G	G	G	G	G	G	G	G	G	G	G
38	C	T	A	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C
39	R	R	R	Q	R	R	R	R	R	R	R	G	G	G	G	G	K	R	G
40	A	A	A	G	A	A	A	A	A	A	A	G	G	G	G	G	G	G	G
41	K	K	K	N	K	K	K	K	K	K	K	N	N	N	N	N	N	N	N
42	R	R	R	N	S	R	R	R	R	R	R	S	A	A	A	A	K	Q	A
43	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N

T06230"56056860

[illegible]

- 1 BPTI
- 2 Engineered BPTI From MARK87
- 3 Engineered BPTI From MARK87
- 4 Bovine Colostrum (DUFT85)
- 5 Bovine Serum (DUFT85)
- 6 Semisynthetic BPTI, TSCH87
- 7 Semisynthetic BPTI, TSCH87
- 8 Semisynthetic BPTI, TSCH87
- 9 Semisynthetic BPTI, TSCH87
- 10 Semisynthetic BPTI, TSCH87
- 11 Engineered BPTI, AUER87
- 12 Dendroaspis polylepis polylepis (Black mamba) venom I (DUFT85)
- 13 Dendroaspis polylepis polylepis (Black Mamba) venom K (DUFT85)
- 14 Hemachatus hemachates (Ringhals Cobra) HHV II (DUFT85)
- 15 Naja nivea (Cape cobra) NNV II (DUFT85)
- 16 Vipera russelli (Russel's viper) RVV II (TAKA74)
- 17 Red sea turtle egg white (DUFT85)
- 18 Snail mucus (Helix pomania) (WAGN78)
- 19 Dendroaspis angusticeps (Eastern green mamba) C13 S1 C3 toxin (DUFT85)

Table 13, Continued

R #	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35
-5	-	-	-	-	-	-	-	-	-	-	-	-	-	D	-	-
-4	-	-	-	-	-	-	-	-	-	-	-	-	-	E	-	-
-3	-	-	-	-	-	-	-	-	-	-	-	-	T	P	-	-
-2	Z	-	L	Z	R	K	-	-	-	R	R	-	E	T	-	-
-1	P	-	Q	D	D	N	-	-	-	Q	K	-	R	T	-	-
1	R	R	H	H	R	R	I	K	T	R	R	R	G	D	K	T
2	R	P	R	P	P	P	N	E	V	H	H	P	F	L	A	V
3	K	Y	T	K	K	T	G	D	A	R	P	D	L	P	D	E
4	L	A	F	F	F	F	D	S	A	D	D	F	D	I	S	A
5	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C
6	I	E	K	Y	Y	N	E	Q	N	D	D	L	T	E	Q	N
7	L	L	L	L	L	L	L	L	L	K	K	E	S	Q	L	L
8	H	I	P	P	P	L	P	G	P	P	P	P	P	P	A	P
9	R	V	A	A	A	P	K	Y	V	P	P	P	P	P	FG	Y
10	N	A	E	D	A	E	V	S	I	D	D	Y	V	D	S	V
11	P	A	P	P	P	T	V	A	R	K	T	T	T	A	Q	Q
12	G	G	G	G	G	G	G	G	G	G	K	G	G	G	G	G
13	R	P	P	R	R	R	P	P	P	N	I	P	P	L	P	P
14	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C
15	Y	M	K	K	L	N	R	M	R	-	-	K	R	F	L	R
16	D	F	A	A	A	A	A	G	A	G	Q	A	A	G	G	A
17	K	F	S	H	Y	L	R	M	F	P	T	K	G	Y	L	F
18	I	I	I	I	M	I	F	T	I	V	V	M	F	M	F	I
19	P	S	P	P	P	P	P	S	Q	R	R	I	K	K	K	Q
20	A	A	A	R	R	A	R	R	L	A	A	R	R	L	R	L
21	F	F	F	F	F	F	Y	Y	W	F	F	Y	Y	Y	Y	W
22	Y	Y	Y	Y	Y	Y	Y	Y	F	A	Y	Y	F	N	S	F
23	Y	Y	Y	Y	Y	Y	Y	Y	F	Y	Y	Y	Y	Y	Y	F
24	N	S	N	D	N	N	N	N	D	D	K	N	N	N	N	D
25	Q	K	W	S	P	S	S	G	A	T	P	A	T	Q	G	A
26	K	G	A	A	A	H	S	T	V	R	S	K	R	E	T	V
27	K	A	A	S	S	L	S	S	K	L	A	A	T	T	S	K
28	K	N	K	N	N	H	K	M	G	K	K	G	K	K	M	G
29	Q	K	K	K	K	K	R	A	K	T	R	F	Q	C	A	K
30	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C
31	E	Y	Q	N	E	Q	E	E	V	K	V	E	E	E	E	V
32	R	P	L	K	K	K	K	T	L	A	Q	T	P	E	T	R
33	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F	F
34	D	T	H	I	I	N	I	Q	P	Q	R	V	K	I	L	S
35	W	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
36	S	S	G	G	G	G	G	G	G	R	G	G	G	G	G	G
37	G	G	G	G	G	G	G	G	G	G	G	G	G	G	G	G
38	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C
39	G	R	K	P	R	G	G	M	Q	D	D	K	K	Q	M	K
40	G	G	G	G	G	G	G	G	G	G	G	A	G	G	G	G
41	N	N	N	N	N	N	N	N	N	D	D	K	N	N	N	N
42	S	A	A	A	A	A	A	G	G	H	H	S	G	D	L	G
43	N	N	N	N	N	N	N	N	N	G	G	N	N	N	N	N

FOIA b 5 - DECLASSIFIED

Table 13, continued

R #	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35
44	R	R	R	N	N	N	N	N	K	N	N	N	R	R	N	K
45	F	F	F	F	F	F	F	F	F	F	F	F	Y	F	F	F
46	K	K	S	K	K	K	H	V	Y	K	K	R	K	S	L	Y
47	T	T	T	T	T	T	T	T	S	T	S	S	S	T	S	S
48	I	I	I	W	W	I	L	E	E	E	D	A	E	L	Q	Q
49	E	E	E	D	D	D	E	K	K	T	H	E	Q	A	K	K
50	E	E	K	E	E	E	E	E	E	L	L	D	D	E	E	E
51	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C
52	R	R	R	R	R	Q	E	L	R	R	R	M	L	E	L	K
53	R	R	H	Q	H	R	K	Q	E	C	C	R	D	Q	Q	E
54	T	T	A	T	T	T	V	T	Y	E	E	T	A	K	T	Y
55	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C
56	I	V	V	G	V	A	G	R	G	L	E	G	S	I	R	G
57	G	V	G	A	A	A	V	-	V	V	L	G	G	N	-	I
58	-	-	-	S	S	K	R	-	P	Y	Y	A	F	-	-	P
59	-	-	-	A	G	Y	S	-	G	P	R	-	-	-	-	G
60	-	-	-	-	I	G	-	-	D	-	-	-	-	-	-	E
61	-	-	-	-	-	-	-	-	E	-	-	-	-	-	-	A

- 20 Dendroaspis angusticeps (Eastern Green Mamba)
C13 S2 C3 toxin (DUFT85)
- 21 Dendroaspis polylepis polylepes (Black mamba) B toxin
(DUFT85)
- 22 Dendroaspis polylepis polylepes (Black Mamba) E toxin
(DUFT85)
- 23 Vipera ammodytes TI toxin (DUFT85)
- 24 Vipera ammodytes CTI toxin (DUFT85)
- 25 Bungarus fasciatus VIII B toxin (DUFT85)
- 26 Anemonia sulcata (sea anemone) 5 II (DUFT85)
- 27 Homo sapiens HI-14 "inactive" domain (DUFT85)
- 28 Homo sapiens HI-8 "active" domain (DUFT85)
- 29 beta bungarotoxin B1 (DUFT85)
- 30 beta bungarotoxin B2 (DUFT85)
- 31 Bovine spleen TI II (FIOR85)
- 32 Tachypleus tridentatus (Horseshoe crab) hemocyte
inhibitor (NAKA87)
- 33 Bombyx mori (silkworm) SCI-III (SASA84)
- 34 Bos taurus (inactive) BI-14
- 35 Bos taurus (active) BI-8

Table 13, continued

R #	36	37	38	39	40
-5	-	-	-	-	-
-4	-	-	-	-	-
-3	-	-	-	-	-
-2	-	-	-	-	-
-1	-	Z	-	-	-
1	R	R	R	R	R
2	P	P	P	P	P
3	D	D	D	D	D
4	F	F	F	F	F
5	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>
6	L	L	L	L	L
7	E	E	E	E	E
8	P	P	P	P	P
9	P	P	P	P	P
10	Y	Y	Y	Y	Y
11	T	T	T	T	T
12	<u>G</u>	<u>G</u>	<u>G</u>	<u>G</u>	<u>G</u>
13	P	P	P	P	P
14	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>
15	R	K	K	K	K
16	A	A	A	A	A
17	R	R	R	R	K
18	I	M	I	M	M
19	I	I	I	I	I
20	R	R	R	R	R
21	Y	Y	Y	Y	Y
22	F	F	F	F	F
23	<u>Y</u>	<u>Y</u>	<u>Y</u>	<u>Y</u>	<u>Y</u>
24	N	N	N	N	N
25	A	A	A	A	A
26	K	K	K	K	K
27	A	A	A	A	A
28	G	G	G	G	G
29	L	L	L	L	F
30	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>
31	Q	Q	Q	Q	E
32	T	P	P	P	T
33	<u>F</u>	<u>F</u>	<u>F</u>	<u>F</u>	<u>F</u>
34	V	V	V	V	V
35	<u>Y</u>	<u>Y</u>	<u>Y</u>	<u>Y</u>	<u>Y</u>
36	G	G	G	G	G
37	<u>G</u>	<u>G</u>	<u>G</u>	<u>G</u>	<u>G</u>
38	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>
39	R	R	R	R	K
40	A	A	A	A	A
41	K	K	K	K	K
42	R	S	R	R	S
43	<u>N</u>	<u>N</u>	<u>N</u>	<u>N</u>	<u>N</u>

FOIA b 5 - EXEMPT

Table 13, continued

R #	36	37	38	39	40
44	N	N	N	N	N
45	<u>F</u>	<u>F</u>	<u>F</u>	<u>F</u>	<u>F</u>
46	K	K	K	K	R
47	S	S	S	S	S
48	A	A	S	A	A
49	E	E	E	E	E
50	D	D	D	D	D
51	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>
52	E	M	M	M	M
53	R	R	R	R	R
54	T	T	T	T	T
55	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>
56	G	G	G	G	G
57	G	G	G	G	G
58	A	A	A	A	A
59	-	-	-	-	-
60	-	-	-	-	-
61	-	-	-	-	-

- 36: Engineered BPTI (KR15, ME52): Auerswald '88, Biol Chem Hoppe-Seyler, 369 Supplement, pp27-35.
- 37: Isoaprotinin G-1: Siekmann, Wenzel, Schroder, and Tschesche '88, Biol Chem Hoppe-Seyler, 369:157-163.
- 38: Isoaprotinin 2: Siekmann, Wenzel, Schroder, and Tschesche '88, Biol Chem Hoppe-Seyler, 369:157-163.
- 39: Isoaprotinin G-2: Siekmann, Wenzel, Schroder, and Tschesche '88, Biol Chem Hoppe-Seyler, 369:157-163.
- 40: Isoaprotinin 1: Siekmann, Wenzel, Schroder, and Tschesche '88, Biol Chem Hoppe-Seyler, 369:157-163.

F06230-5696860

Table 13, continued

Notes :

- a) both beta bungarotoxins have residue 15 deleted.
- b) B. mori has an extra residue between C5 and C14; we have assigned F and G to residue 9.
- c) all natural proteins have C at 5, 14, 30, 38, 50, & 55.
- d) all homologues have F33 and G37.
- e) extra C's in bungarotoxins form interchain cystine bridges

T06290"56096860

	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100	2101	2102	2103	2104	2105	2106	2107	2108	2109	2110	2111	2112	2113	2114	2115	2116	2117	2118	2119	2120	2121	2122	2123	2124	2125	2126	2127	2128	2129	2130	2131	2132	2133	2134	2135	2136	2137	2138	2139	2140	2141	2142	2143	2144	2145	2146	2147	2148	2149	2150	2151	2152	2153	2154	2155	2156	2157	2158	2159	2160	2161	2162	2163	2164	2165	2166	2167	2168	2169	2170	2171	2172	2173	2174	2175	2176	2177	2178	2179	2180	2181	2182	2183	2184	2185	2186	2187	2188	2189	2190	2191	2192	2193	2194	2195	2196	2197	2198	2199	2200	2201	2202	2203	2204	2205	2206	2207	2208	2209	2210	2211	2212	2213	2214	2215	2216	2217	2218	2219	2220	2221	2222	2223	2224	2225	2226	2227	2228	2229	2230	2231	2232	2233	2234	2235	2236	2237	2238	2239	2240	2241	2242	2243	2244	2245	2246	2247	2248	2249	2250	2251	2252	2253	2254	2255	2256	2257	2258	2259	2260	2261	2262	2263	2264	2265	2266	2267	2268	2269	2270	2271	2272	2273	2274	2275	2276	2277	2278	2279	2280	2281	2282	2283	2284	2285	2286	2287	2288	2289	2290	2291	2292	2293	2294	2295	2296	2297	2298	2299	2300	2301	2302	2303	2304	2305	2306	2307	2308	2309	2310	2311	2312	2313	2314	2315	2316	2317	2318	2319	2320	2321	2322	2323	2324	2325	2326	2327	2328	2329	2330	2331	2332	2333	2334	2335	2336	2337	2338	2339	2340	2341	2342	2343	2344	2345	2346	2347	2348	2349	2350	2351	2352	2353	2354	2355	2356	2357	2358	2359	2360	2361	2362	2363	2364	2365	2366	2367	2368	2369	2370	2371	2372	2373	2374	2375	2376	2377	2378	2379	2380	2381	2382	2383	2384	2385	2386	2387	2388	2389	2390	2391	2392	2393	2394	2395	2396	2397	2398	2399	2400	2401	2402	2403	2404	2405	2406	2407	2408	2409	2410	2411	2412	2413	2414	2415	2416	2417	2418	2419	2420	2421	2422	2
--	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	---

- 1 BPTI
- 2 synthetic BPTI, Tan & Kaiser, biochem. 16(8)1531-41
- 3 Semisynthetic BPTI, TSCH87
- 4 Semisynthetic BPTI, TSCH87
- 5 Semisynthetic BPTI, TSCH87
- 6 Semisynthetic BPTI, TSCH87
- 7 Semisynthetic BPTI, TSCH87
- 8 Engineered BPTI, AUER87
- 9 BPTI Auerswald & al GB 2 208 511A
- 10 BPTI Auerswald & al GB 2 208 511A
- 11 Engineered BPTI From MARK87
- 12 Engineered BPTI From MARK87
- 13 BPTI(KR15,ME52): Auerswald '88, Biol Chem Hoppe-Seyler, 369 Suppl, pp27-35.
- 14 BPTI CA30/CA51 Eigenbrot & al, Protein Engineering 3(7)591-598 ('90)
- 15 Isoaprotinin 2 Siekmann et al '88, Biol Chem Hoppe-Seyler, 369:157-163.
- 16 Isoaprotinin G-2: Siekmann et al '88, Biol Chem Hoppe-Seyler, 369:157-163.
- 17 BPTI Engineered, Auerswald & al GB 2 208 511A
- 18 BPTI Engineered, Auerswald & al GB 2 208 511A
- 19 BPTI Engineered, Auerswald & al GB 2 208 511A
- 20 Isoaprotinin G-1 Siekmann & al '88, Biol Chem Hoppe-Seyler, 369:157-163.
- 21 BPTI Engineered, Auerswald & al GB 2 208 511A
- 22 BPTI Engineered, Auerswald & al GB 2 208 511A
- 23 Bovine Serum (in Dufton '85)
- 24 Bovine spleen TI II (FIOR85)
- 25 Snail mucus (Helix pomatia) (WAGN78)
- 26 Hemachatus hemachates (Ringhals Cobra) HHV II (in Dufton '85)
- 27 Red sea turtle egg white (in Dufton '85)
- 28 Bovine Colostrum (in Dufton '85)
- 29 Naja nivea (Cape cobra) NNV II (in Dufton '85)
- 30 Bungarus fasciatus VIII B toxin (in Dufton '85)
- 31 Vipera ammodytes TI toxin (in Dufton '85)
- 32 Porcine ITI domain 1, (in CREI87)
- 33 Human Alzheimer's beta APP protease inhibitor, (SHIN90)
- 34 Equine ITI domain 1, in Creighton & Charles
- 35 Bos taurus (inactive) BI-8e (ITI domain 1)
- 36 Anemonia sulcata (sea anemone) 5 II (in Dufton '85)
- 37 Dendroaspis polylepis polylepes (Black Mamba) E toxin (in Dufton '85)
- 38 Vipera russelli (Russel's viper) RVV II (TAKA74)
- 39 Tachypleus tridentatus (Horseshoe crab) hemocyte inhibitor (NAKA87)
- 40 LACI 2 (Factor Xa) (WUNT88)
- 41 Vipera ammodytes CTI toxin (in Dufton '85)

Identification codes for Tables 14 and 15

- 42 Dendroaspis polylepis polylepis (Black Mamba) venom K (in Dufton '85)
- 43 Homo sapiens HI-8e "inactive" domain (in Dufton '85)
- 44 Green Mamba toxin K, (in CREI87)
- 45 Dendroaspis angusticeps (Eastern green mamba) C13 S1 C3 toxin (in Dufton '85)
- 46 LACI 3
- 47 Equine ITI domain 2, (CREI87)
- 48 LACI 1 (VIIa)
- 49 Dendroaspis polylepis polylepis (Black mamba) B toxin (in Dufton '85)
- 50 Porcine ITI domain 2, Creighton and Charles
- 51 Homo sapiens HI-8t "active" domain (in Dufton '85)
- 52 Bos taurus (active) BI-8t
- 53 Trypstatin Kito & al ('88) J Biol Chem 263(34)18104-07
- 54 Dendroaspis angusticeps (Eastern Green Mamba) C13 S2 C3 toxin (in Dufton '85)
- 55 Green Mamba I venom Creighton & Charles '87 CSHSQB 52:511-519.
- 56 beta bungarotoxin B2 (in Dufton '85)
- 57 Dendroaspis polylepis polylepis (Black mamba) venom I (in Dufton '85)
- 58 beta bungarotoxin B1 (in Dufton '85)
- 59 Bombyx mori (silkworm) SCI-III (SASA84)

T06250"56096860

Table 14: Tally of Ionizable groups

Identifier	D	E	K	R	Y	H	NH	CO2	+	ions
1	2	2	4	6	4	0	1	1	6	16
2	2	2	4	6	4	0	1	1	6	16
3	2	2	3	6	4	0	1	1	5	15
4	2	2	3	6	4	0	1	1	5	15
5	2	2	3	6	4	0	1	1	5	15
6	2	2	3	6	4	0	1	1	5	15
7	2	2	3	6	4	0	1	1	5	15
8	2	3	4	6	4	0	1	1	5	17
9	2	2	3	5	4	0	1	1	4	14
10	2	3	3	6	4	0	1	1	4	16
11	2	2	4	6	4	0	1	1	6	16
12	2	2	4	6	4	0	1	1	6	16
13	2	3	3	7	4	0	1	1	5	17
14	2	2	4	6	4	0	1	1	6	16
15	2	2	4	6	4	0	1	1	6	16
16	2	2	4	6	4	0	1	1	6	16
17	2	2	3	5	4	0	1	1	4	14
18	2	3	3	5	4	0	1	1	3	15
19	2	3	3	5	4	0	1	1	3	15
20	2	2	4	5	4	0	1	1	5	15
21	2	3	3	4	4	0	1	1	2	14
22	2	4	3	4	4	0	1	1	1	15
23	2	4	4	4	4	0	1	1	2	16
24	2	3	5	4	4	0	1	1	4	16
25	1	1	2	4	4	0	1	1	4	10
26	2	3	2	5	3	1	1	1	2	14
27	2	4	6	8	3	0	1	1	8	22
28	2	4	2	3	3	0	1	1	-1	13
29	1	4	2	7	2	2	1	1	4	16
30	1	2	5	3	4	2	1	1	5	13
31	4	1	5	3	4	2	1	1	3	15
32	1	4	3	2	4	1	1	1	0	12
33	2	6	1	5	3	0	1	1	-2	16
34	2	4	2	2	3	1	1	1	-2	12
35	2	2	3	2	4	0	1	1	1	11
36	1	5	4	5	4	1	1	1	3	17
37	0	2	6	3	3	3	1	1	7	13
38	2	5	3	7	3	2	1	1	3	19
39	3	3	5	5	4	0	1	1	4	18
40	3	7	4	3	4	0	1	1	-3	19
41	3	2	4	6	5	1	1	1	5	17
42	1	2	8	5	4	0	1	1	10	18
43	1	4	2	2	4	0	1	1	-1	11
44	1	2	9	4	5	0	1	1	10	18
45	0	2	8	4	5	0	1	1	10	16
46	1	3	5	5	3	0	1	1	6	16
47	3	4	4	3	3	0	1	1	0	16
48	3	6	5	4	1	1	1	1	0	20
49	0	3	3	5	5	0	1	1	5	13
50	2	6	4	2	3	0	1	1	-2	16

T06230" 56036360

Table 14: Tally of Ionizable groups

Identifier	D	E	K	R	Y	H	NH	CO2	+	ions
51	2	4	4	3	3	0	1	1	1	15
52	1	4	6	2	3	0	1	1	3	15
53	2	2	5	1	4	0	1	1	2	12
54	2	3	6	8	3	1	1	1	9	21
55	1	3	6	7	3	1	1	1	9	19
56	6	2	6	7	4	3	1	1	5	23
57	0	3	7	7	3	1	1	1	11	19
58	6	2	5	7	4	2	1	1	4	22
59	4	7	3	1	4	0	1	1	-7	17

T06290"56096260

[illegible]

Table 15: Frequency of Amino Acids at Each Position
in BPTI and 58 Homologues (continued)

Res. Id.	Different AAs	Contents	First
39	9	R25 G13 K6 Q4 E3 M3 L2 D2 P	R
40	2	G35 A24	A
41	3	N33 K24 D2	K
42	12	R22 A12 G8 S6 Q2 H2 N2 M D E K L	R
43	2	N57 G2	N
44	3	N40 R14 K5	N
45	2	F58 Y	F
46	11	K39 Y5 E4 S2 V2 D2 R H T A L	K
47	2	S36 T23	S
48	11	A23 I11 E6 Q6 L4 K2 T2 W2 S D R	A
49	8	E37 K8 D6 Q3 A2 P H T	E
50	7	E27 D25 K2 L2 M Q Y	D
51	2	C58 A	C
52	9	M17 R15 E8 L7 K6 Q2 T2 H V	M
53	11	R37 E6 Q5 K2 C2 H2 A N G D W	R
54	8	T41 Y5 A4 V3 I2 E2 M K	T
55	1	C59	C
56	10	G33 V9 R5 I4 E3 L A S T K	G
57	12	G34 V6 -5 A3 R2 I2 P2 D K S L N	G
58	10	A25 -15 P7 K3 S2 Y2 G2 F D R	A

106230 5503660

Coordinates taken from
Brookhaven Protein Data Bank entry 6PTI.

```

HEADER      PROTEINASE INHIBITOR (TRYPSIN) 13-MAY-87      6PTI
COMPND      BOVINE PANCREATIC TRYPSIN INHIBITOR
COMPND      2(/BPTIS, CRYSTAL FORM /III$)
AUTHOR      A.WLODAWER

```

Solvent radius = 1.40
Atomic radii given in Table 7

Areas in Å².

Residue		Total area	Not Covered by M/C	fraction	Not covered at all	fraction
ARG	1	342.45	205.09	0.5989	152.49	0.4453
PRO	2	239.12	92.65	0.3875	47.56	0.1989
ASP	3	272.39	158.77	0.5829	143.23	0.5258
PHE	4	311.33	137.82	0.4427	43.21	0.1388
CYS	5	241.06	48.36	0.2006	0.23	0.0010
LEU	6	280.98	151.45	0.5390	115.87	0.4124
GLU	7	291.39	128.91	0.4424	90.39	0.3102
PRO	8	236.12	128.71	0.5451	99.98	0.4234
PRO	9	236.09	109.82	0.4652	45.80	0.1940
TYR	10	330.97	153.63	0.4642	79.49	0.2402
THR	11	249.20	80.10	0.3214	64.99	0.2608
GLY	12	184.21	56.75	0.3081	23.05	0.1252
PRO	13	240.07	130.25	0.5426	75.27	0.3136
CYS	14	237.10	75.55	0.3186	53.52	0.2257
LYS	15	310.77	200.25	0.6444	192.00	0.6178
ALA	16	209.41	66.63	0.3182	45.59	0.2177
ARG	17	351.09	243.67	0.6940	201.48	0.5739
ILE	18	277.10	100.51	0.3627	58.95	0.2127
ILE	19	278.03	146.06	0.5254	96.05	0.3455
ARG	20	339.11	144.65	0.4266	43.81	0.1292
TYR	21	333.60	102.24	0.3065	69.67	0.2089
PHE	22	306.08	70.64	0.2308	23.01	0.0752
TYR	23	338.66	77.05	0.2275	17.34	0.0512
ASN	24	264.88	99.03	0.3739	38.69	0.1461
ALA	25	211.15	85.13	0.4032	48.20	0.2283
LYS	26	313.29	216.14	0.6899	202.84	0.6474
ALA	27	210.66	96.05	0.4560	54.78	0.2601
GLY	28	186.83	71.52	0.3828	32.09	0.1718
LEU	29	280.70	132.42	0.4718	93.61	0.3335
CYS	30	238.15	57.27	0.2405	19.33	0.0812
GLN	31	301.15	141.80	0.4709	82.64	0.2744
THR	32	251.26	138.17	0.5499	76.47	0.3043

Table 16, continued.

PHE 33	304.27	59.79	0.1965	18.91	0.0622
VAL 34	251.56	109.78	0.4364	42.36	0.1684
TYR 35	332.64	80.52	0.2421	15.05	0.0452
GLY 36	187.06	11.90	0.0636	1.97	0.0105
GLY 37	185.28	84.26	0.4548	39.17	0.2114
CYS 38	234.56	73.64	0.3139	26.40	0.1125
ARG 39	417.13	304.62	0.7303	250.73	0.6011
ALA 40	209.53	94.01	0.4487	52.95	0.2527
LYS 41	314.60	166.23	0.5284	108.77	0.3457
ARG 42	349.06	232.83	0.6670	179.59	0.5145
ASN 43	266.47	38.53	0.1446	5.32	0.0200
ASN 44	269.65	91.08	0.3378	23.39	0.0867
PHE 45	313.22	69.73	0.2226	14.79	0.0472
LYS 46	309.83	217.18	0.7010	155.73	0.5026
SER 47	224.78	69.11	0.3075	24.80	0.1103
ALA 48	211.01	82.06	0.3889	31.07	0.1473
GLU 49	286.62	161.00	0.5617	100.01	0.3489
ASP 50	299.53	156.42	0.5222	95.96	0.3204
CYS 51	238.68	24.51	0.1027	0.00	0.0000
MET 52	293.05	89.48	0.3054	66.70	0.2276
ARG 53	356.20	224.61	0.6306	189.75	0.5327
THR 54	251.53	116.43	0.4629	51.64	0.2053
CYS 55	240.40	69.95	0.2910	0.00	0.0000
GLY 56	184.66	60.79	0.3292	32.78	0.1775
GLY 57	106.58	49.71	0.4664	38.28	0.3592
ALA 58	no position given in Protein Data Bank				

"Total area" is the area measured by a rolling sphere of radius 1.4 Å, where only the atoms within the residue are considered. This takes account of conformation.

"Not covered by M/C" is the area measured by a rolling sphere of radius 1.4 Å where all main-chain atoms are considered, fraction is the exposed area divided by the total area. Surface buried by main-chain atoms is more definitely covered than is surface covered by side group atoms.

"Not covered at all" is the area measured by a rolling sphere of radius 1.4 Å where all atoms of the protein are considered.

Table 17: Plasmids used in Detailed Example I

<u>Phage</u>	<u>Contents</u>
LG1	M13mp18 with <u>Ava</u> II/ <u>Aat</u> II/ <u>Acc</u> I/ <u>Rsr</u> II/ <u>Sau</u> I adaptor
pLG2	LG1 with <u>amp</u> ^R and ColE1 of pBR322 cloned into <u>Aat</u> II/ <u>Acc</u> I sites
pLG3	pLG2 with <u>Acc</u> I site removed
pLG4	pLG3 with first part of <u>osp-pbd</u> gene cloned into <u>Rsr</u> II/ <u>Sau</u> I sites, <u>Avr</u> II/ <u>Asu</u> II sites created
pLG5	pLG4 with second part of <u>osp-pbd</u> gene cloned into <u>Avr</u> II/ <u>Asu</u> II sites, <u>BssH</u> I site created
pLG6	pLG5 with third part of <u>osp-pbd</u> gene cloned into <u>Asu</u> II/ <u>BssH</u> I sites, <u>Bbe</u> I site created
pLG7	pLG6 with last part of <u>osp-pbd</u> gene cloned into <u>Bbe</u> I/ <u>Asu</u> II sites
pLG8	pLG7 with disabled <u>osp-pbd</u> gene, same length DNA.
pLG9	pLG7 mutated to display BPTI(V15 _{BPTI})
pLG10	pLG8 + <u>tet</u> ^R gene - <u>amp</u> ^R gene
pLG11	pLG9 + <u>tet</u> ^R gene - <u>amp</u> ^R gene

T06290" 55035860

5	<u>AhaII</u>	<u>NarI</u>	<u>GdiII</u>	<u>PvuI</u>
	<u>FspI</u>	<u>BglI</u>	<u>HgiEII</u>	<u>Bsu36I</u>
10	<u>EcoRI</u>	<u>SacI</u>	<u>KpnI</u>	<u>XmaI</u>
	<u>SmaI</u>	<u>BamHI</u>	<u>XbaI</u>	<u>SalI</u>
	<u>HindIII</u>	<u>AccI</u>	<u>PstI</u>	<u>SphI</u>
15	<u>HindII</u>			

20	M13mp18			
	<u>AatII</u>	<u>AflI</u>	<u>ApaI</u>	<u>AvrII</u>
25	<u>BbvII</u>	<u>BclI</u>	<u>BspMI</u>	<u>BssHI</u>
	<u>BstBI</u>	<u>BstEII</u>	<u>BstXI</u>	<u>EaqI</u>
	<u>Eco57I</u>	<u>EcoNI</u>	<u>EcoOI09I</u>	<u>EcoRV</u>
30	<u>EspI</u>	<u>HpaI</u>	<u>MluI</u>	<u>NcoI</u>
	<u>NheI</u>	<u>NotI</u>	<u>NruI</u>	<u>NsiI</u>
35	<u>PflMI</u>	<u>PmaCI</u>	<u>PpaI</u>	<u>PpuMI</u>
	<u>RsrI</u>	<u>SacI</u>	<u>ScaI</u>	<u>SfiI</u>
	<u>SpeI</u>	<u>StuI</u>	<u>StyI</u>	<u>Tth111I</u>
40	<u>XcaI</u>	<u>XhoI</u>		

Table 20: Enzymes cutting
AmpR gene and ori

5	<u>AatII</u>	<u>BbvII</u>	<u>Eco57I</u>	<u>PpaI</u>
	<u>ScaI</u>	<u>Tth111I</u>	<u>AhaII</u>	<u>GdiII</u>
	<u>PvuI</u>	<u>FspI</u>	<u>BqlI</u>	<u>HgiEII</u>
10	<u>HindII</u>	<u>PstI</u>	<u>XbaI</u>	<u>AflIII</u>
	<u>NdeI</u>			
15				

T06250-56056B50

Table 21: Enzymes tested on Ambig DNA

	Enzyme	Recognition	Symm	cuts	Supply
5	% <u>Acc</u> I	GTMKAC	P	2 & 4	<B,M,I,N,P,T
	<u>Afl</u> II	CTTAAG	P	1 & 5	<N
	<u>Apa</u> I	GGGCCC	P	5 & 1	<M,I,N,P,T
	<u>Asu</u> II	TTCGAA	P	2 & 4	<P,N(<u>Bst</u> BI)
	<u>Ava</u> III	ATGCAT	P	5 & 1	<T; <u>Nsi</u> I:M,N,P,T; <u>Eco</u> T22I:T
10	<u>Avr</u> II	CCTAGG	P	1 & 5	<N
	<u>Bam</u> HI	GGATCC	P	1 & 5	<S,B,M,I,N,P,T
	<u>Bcl</u> I	TGATCA	P	1 & 5	<S,B,M,I,N,T
	<u>Bsp</u> MII	TCCGGA	P	1 & 5	<N
15	<u>Bss</u> HII	GCGCGC	P	1 & 5	<N,T
	+ <u>Bst</u> EII	GGTNACC	P	1 & 6	<S,B,M,N,T
	% <u>Bst</u> XI	CCANNNNN	P	8 & 4	<N,P,T
	+ <u>Dra</u> II	RGGNCCY	P	2 & 5	<M,T ; <u>Eco</u> O109I:N
	+ <u>Eco</u> NI	CCTNNNNN	P	5 & 6	<N(soon)
20	<u>Eco</u> RI	GAATTC	P	1 & 5	<S,B,M,I,N,P,T
	<u>Eco</u> RV	GATATC	P	3 & 3	<S,B,M,I,N,P,T
	+ <u>Esp</u> I	GCTNAGC	P	2 & 5	<T
	<u>Hind</u> III	AAGCTT	P	1 & 5	<S,B,M,I,N,P,T
	<u>Hpa</u> I	GTTAAC	P	3 & 3	<S,B,M,I,N,P,T
25	<u>Kpn</u> I	GGTACC	P	5 & 1	<S,B,M,I,N,P,T ; <u>Asp</u> 718:M
	<u>Mlu</u> I	ACGCGT	P	1 & 5	<M,N,P,T
	<u>Nar</u> I	GGCGCC	P	2 & 4	<B,N,T
	<u>Nco</u> I	CCATGG	P	1 & 5	<B,M,N,P,T
30	<u>Nhe</u> I	GCTAGC	P	1 & 5	<M,N,P,T
	<u>Not</u> I	GCGGCCGC	P	2 & 6	<M,N,P,T
	<u>Nru</u> I	TCGCGA	P	3 & 3	<B,M,N,T
	+ <u>Pfl</u> MI	CCANNNNN	P	7 & 4	<N
	<u>Pma</u> CI	CACGTG	P	3 & 3	<none
35	+ <u>Ppu</u> MI	RGGWCCY	P	2 & 5	<N
	+ <u>Rsr</u> II	CGGWCCG	P	2 & 5	<N,T
	<u>Sac</u> I	GAGCTC	P	5 & 1	<B(<u>Sst</u> I),M,I,N,P, T
	<u>Sal</u> I	GTCGAC	P	1 & 5	<B,M,I,N,P,T
40	+ <u>Sau</u> I	CCTNAGG	P	2 & 5	<M; <u>Cvn</u> I:B; <u>Mst</u> II :T; <u>Bsu</u> 36I:N; <u>Aoc</u> I:T
	+ <u>Sfi</u> II	GGCCNNNNNGGCC	P	8 & 5	<N,P,T
	<u>Sma</u> I	CCCGGG	P	3 & 3	<B,M,I,N,P,T
	<u>Spe</u> I	ACTAGT	P	1 & 5	<M,N,T
45	<u>Sph</u> I	GCATGC	P	5 & 1	<B,M,I,N,P,T
	<u>Stu</u> I	AGGCCT	P	3 & 3	<M,N,I(<u>Aat</u> I),P,T
	% <u>Sty</u> I	CCWWGG	P	1 & 5	<N,P,T
	<u>Xca</u> I	GTATAC	P	3 & 3	<N(soon)

T06290" 55096860

Table 21, continued.

5	<u>Xho</u> I	CTCGAG	P	1 &	5 <B,M,I,P,T; <u>Ccr</u> I: T ; <u>Paer</u> 7I:N
	<u>Xma</u> I	CCCGGG	P	1 &	5 <I,N,P,T
	<u>Xma</u> III	CGGCCG	P	1 &	5 <B; <u>Eag</u> I:N; <u>Eco</u> 52I:T
10	N_restrct =		43		

15

T06230"5505660

Table 22: ipbd gene

pbdb mod10 29III88 :

lacUV5 RsrII/AvrII/gene/TrpA attenuator/MstII; !

5'- CGGACCG TaT ! RsrII site
 CCAGGC tttaca CTTTATGCTTCCGGCTCG tataat GTG ! lacUV5
 TGG aATTGTGAGCGGATAACAATT ! lacO operator
 CCT AGGAgg CtcaCT ! Shine-Dalgarno seq.
 atg aag aaa tct ctg gtt ctt aag gct agc ! 10, M13 leader
 gtt gct gtc gcg acc ctg gta ccg atg ctg ! 20
 tct ttt gct cgt ccg gat ttc tgt ctc gag ! 30
 ccg cca tat act ggg ccc tgc aaa gcg cgc ! 40
 atc atc cgt tat ttc tac aac gct aaa gca ! 50
 ggc ctg tgc cag acc ttt gta tac ggt ggt ! 60
 tgc cgt gct aag cgt aac aac ttt aaa tcg ! 70
 gcc gaa gat tgc atg cgt acc tgc ggt ggc ! 80
 gcc gct gaa ggt gat gat ccg gcc aaa gcg ! 90
 gcc ttt aac tct ctg caa gct tct gct acc ! 100
 gaa tat atc ggt tac gcg tgg gcc atg gtg ! 110
 gtg gtt atc gtt ggt gct acc atc ggt atc ! 120
 aaa ctg ttt aag aaa ttt act tcg aaa gcg ! 130
 tct taa tag tga ggttacc ! BstEII
 agtcta agccgc ctaatga gcgggct ttttttt ! terminator
 CCTgAGG -3' ! MstII

Table 23: ipbd DNA sequence

DNA Sequence file = UV5_M13PTIM13.DNA;17

DNA Sequence title =

pbd mod10 29III88 : lac-UV5 RsrII/AvrII/gene/TrpA

attenuator/MstII; !

1 C|GGA|CCG|TAT|CCA|GGC|TTT|ACA|CTT|TAT|GCT|TCC|GGC|TCG|

41 TAT|AAT|GTG|TGG|AAT|TGT|GAG|CGG|ATA|ACA|ATT|CCT|AGG|AGG|

83 CTC|ACT|ATG|AAG|AAA|TCT|CTG|GTT|CTT|AAG|GCT|AGC|GTT|GCT|

125 GTC|GCG|ACC|CTG|GTA|CCG|ATG|CTG|TCT|TTT|GCT|CGT|CCG|GAT|

167 TTC|TGT|CTC|GAG|CCG|CCA|TAT|ACT|GGG|CCC|TGC|AAA|GCG|CGC|

209 ATC|ATC|CGT|TAT|TTC|TAC|AAC|GCT|AAA|GCA|GGC|CTG|TGC|CAG|

251 ACC|TTT|GTA|TAC|GGT|GGT|TGC|CGT|GCT|AAG|CGT|AAC|AAC|TTT|

293 AAA|TCG|GCC|GAA|GAT|TGC|ATG|CGT|ACC|TGC|GGT|GGC|GCC|GCT|

335 GAA|GGT|GAT|GAT|CCG|GCC|AAA|GCG|GCC|TTT|AAC|TCT|CTG|CAA|

377 GCT|TCT|GCT|ACC|GAA|TAT|ATC|GGT|TAC|GCG|TGG|GCC|ATG|GTG|

419 GTG|GTT|ATC|GTT|GGT|GCT|ACC|ATC|GGT|ATC|AAA|CTG|TTT|AAG|

461 AAA|TTT|ACT|TCG|AAA|GCG|TCT|TAA|TAG|TGA|GGT|TAC|CAG|TCT|

503 AAG|CCC|GCC|TAA|TGA|GCG|GGC|TTT|TTT|TTT|CCT|GAG|G

Total = 539 bases

Table 24: Summary of Restriction Cuts

```

Enz = %Acc I has 1 observed sites : 259
Enz = Acc III has 1 observed sites : 162
Enz = Acy I has 1 observed sites : 328
Enz = Afl II has 1 observed sites : 109
Enz = %Afl III has 1 observed sites : 404
Enz = Aha III has 1 observed sites : 292
Enz = Apa I has 1 observed sites : 193
Enz = Asp718 has 1 observed sites : 138
Enz = Asu II has 1 observed sites : 471
Enz = %Ava I has 1 observed sites : 175
Enz = Avr II has 1 observed sites : 76
Enz = %Ban I has 3 observed sites : 138 328 540
Enz = Bbe I has 1 observed sites : 328
Enz = +Bgl I has 1 observed sites : 352
Enz = +Bin I has 1 observed sites : 346
Enz = %BspM I has 1 observed sites : 319
Enz = BssH II has 1 observed sites : 205
Enz = +BstE II has 1 observed sites : 493
Enz = %BstX I has 1 observed sites : 413
Enz = Cfr I has 2 observed sites : 299 350
Enz = +Dra II has 1 observed sites : 193
Enz = +Esp I has 1 observed sites : 277
Enz = %Fok I has 1 observed sites : 213
Enz = Gdi II has 2 observed sites : 299 350
Enz = Hae I has 1 observed sites : 240
Enz = Hae II has 1 observed sites : 328
Enz = +Hga I has 1 observed sites : 478
Enz = %HgiC I has 3 observed sites : 138 328 540
Enz = %HgiJ II has 1 observed sites : 193
Enz = Hind III has 1 observed sites : 377
Enz = +Hph I has 1 observed sites : 340
Enz = Kpn I has 1 observed sites : 138
Enz = +Mbo II has 2 observed sites : 93 304
Enz = Mlu I has 1 observed sites : 404
Enz = Nar I has 1 observed sites : 328
Enz = Nco I has 1 observed sites : 413
Enz = Nhe I has 1 observed sites : 115
Enz = Nru I has 1 observed sites : 128
Enz = Nsp(7524) has 1 observed sites : 311
Enz = NspB II has 1 observed sites : 332
Enz = +PflM I has 1 observed sites : 184
Enz = +Pss I has 1 observed sites : 193
Enz = +Rsr II has 1 observed sites : 3
Enz = +Sau I has 1 observed sites : 535
Enz = %SfaN I has 2 observed sites : 144 209
Enz = +Sfi I has 1 observed sites : 351
Enz = Sph I has 1 observed sites : 311
Enz = Stu I has 1 observed sites : 240
Enz = %Sty I has 2 observed sites : 76 413

```

Table 24, continued.

Enz = Xca I has 1 observed sites : 259
 Enz = Xho I has 1 observed sites : 175
 Enz = Xma III has 1 observed sites : 299

Enzymes that do not cut

<u>Aat</u> II	<u>AlwN</u> I	<u>ApaI</u> I	<u>Ase</u> I	<u>Ava</u> III
<u>Bal</u> I	<u>BamH</u> I	<u>Bbv</u> I	<u>Bbv</u> II	<u>Bcl</u> I
<u>Bgl</u> II	<u>Bsm</u> I	<u>BspH</u> I	<u>Cla</u> I	<u>Dra</u> III
<u>Eco47</u> III	<u>EcoN</u> I	<u>EcoR</u> I	<u>EcoR</u> V	<u>HgiA</u> I
<u>Hinc</u> II	<u>Hpa</u> I	<u>Mst</u> I	<u>Nae</u> I	<u>Nde</u> I
<u>Not</u> I	<u>Ple</u> I	<u>PmaC</u> I	<u>PpuM</u> I	<u>Pst</u> I
<u>Pvu</u> I	<u>Pvu</u> II	<u>Sac</u> I	<u>Sac</u> II	<u>Sal</u> I
<u>Sca</u> I	<u>Sma</u> I	<u>SnaB</u> I	<u>Spe</u> I	<u>Ssp</u> I
<u>Taq</u> II	<u>Tth111</u> I	<u>Tth111</u> II	<u>Xho</u> II	<u>Xma</u> I
<u>Xmn</u> I				

T06290"56056860

Table 25: Annotated Sequence of ipbd gene

5'-	C	GGA	CCG	TAT	CCA	GGC	TTT	ACA	CTT	TAT	28
		<u>Rsr II</u>					<u>-35</u>				
	GCT	TCC	GGC	TCG	TAT	AAT	GTG	TGG			52
				<u>-10</u>							
	AAT	TGT	GAG	CGG	ATA	ACA	ATT				73
		<u>lac operator</u>									
	CCT	AGG	AGG	CTC	ACT						88
	<u>Avr II</u>										
	<u>S. D.</u>										
m	k	k	s	l	v	l	k	a	s		
1	2	3	4	5	6	7	8	9	10		
ATG	AAG	AAA	TCT	CTG	GTT	CTT	AAG	GCT	AGC		118
						<u>Afl II</u>		<u>Nhe I</u>			
v	a	v	a	t	l	v	p	m	l		
11	12	13	14	15	16	17	18	19	20		
GTT	GCT	GTC	GCG	ACC	CTG	GTA	CCG	ATG	CTG		148
		<u>Nru I</u>				<u>Kpn I</u>					
s	f	a	r	p	d	f	c	l	e		
21	22	23	24	25	26	27	28	29	30		
TCT	TTT	GCT	CGT	CCG	GAT	TTC	TGT	CTC	GAG		178
			<u>AccIII</u>					<u>Ava I</u>			
								<u>Xho I</u>			
p	p	y	t	g	p	c	k	a	r		
31	32	33	34	35	36	37	38	39	40		
CCG	CCA	TAT	ACT	GGG	CCC	TGC	AAA	GCG	CGC		208
	<u>PflM I</u>							<u>BssH II</u>			
				<u>Apa I</u>							
				<u>Dra II</u>							
				<u>Pss I</u>							
i	i	r	y	f	y	n	a	k			
41	42	43	44	45	46	47	48	49			
ATC	ATC	CGT	TAT	TTC	TAC	AAC	GCT	AAA			235

Table 25, continued.

a	g	l	c	q	t	f	v	y	g	g
50	51	52	53	54	55	56	57	58	59	60
GCA	GGC	CTG	TGC	CAG	ACC	TTT	GTA	TAC	GGT	GGT
Stu I						Acc I				
						Xca I				

268

c	r	a	k	r	n	n	f	k
61	62	63	64	65	66	67	68	69
TGC	CGT	GCT	AAG	CGT	AAC	AAC	TTT	AAA
Esp I								

295

s	a	e	d	c	m	r	t	c	g
70	71	72	73	74	75	76	77	78	79
TCG	GCC	GAA	GAT	TGC	ATG	CGT	ACC	TGC	GGT
XmaIII				Sph I					

325

g	a	a	e	g	d	d
80	81	82	83	84	85	86
GGC	GCC	GCT	GAA	GGT	GAT	GAT
Bbe I						
Nar I						

346

p	a	k	a	a
87	88	89	90	91
CCG	GCC	AAA	GCG	GCC
Sfi I				

361

f	n	s	l	q	a	s	a	t
92	93	94	95	96	97	98	99	100
TTT	AAC	TCT	CTG	CAA	GCT	TCT	GCT	ACC
Hind 3								

388

e	y	i	g	y	a	w
101	102	103	104	105	106	107
GAA	TAT	ATC	GGT	TAC	GCG	TGG
Mlu I						

409

a	m	v	v	v
108	109	110	111	112
GCC	ATG	GTG	GTG	GTT
BstX I				
Nco I				

424

Table 25, continued.

i	v	g	a	t	i	g	i
113	114	115	116	117	118	119	120
ATC	GTT	GGT	GCT	ACC	ATC	GGT	ATC

448

k	l	f	k	k	f	t	s	k	a
121	122	123	124	125	126	127	128	129	130
AAA	CTG	TTT	AAG	AAA	TTT	ACT	TCG	AAA	GCG

478

Asu II

s	.	.	.
131	132	133	134
TCT	TAA	TAG	TGA

502

BstE II

AAG	CCC	GCC	TAA	TGA	GCG	GGC	TTT	TTT	TTT
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

532

Trp terminator

CCT	GAG	G -3'
-----	-----	-------

539

Sau I

Note the following enzyme equivalences,

<u>Xma</u> III	=	<u>Eag</u> I
<u>Acc</u> III	=	<u>BspM</u> II
<u>Dra</u> II	=	<u>EcoO109</u> I
<u>Asu</u> II	=	<u>BstB</u> I
<u>Sau</u> I	=	<u>Bsu36</u> I

F06230-5505660

Table 26: DNA_seq1

5' | ccg|tcc|gtc|GGA|CCG|TAT|CCA|GGC|TTT|ACA|CTT|TAT|
 | spacer | Rsr II | | -35 |

| GCT|TCC|GGC|TCG|TAT|AAT|GTG|TGG|
 | -10 |

| AAT|TGT|GAG|CGG|ATA|ACA|ATT|
 | lac operator |

| CCT|AGG|
 | Avr II |

			s	k	a
			128	129	130
gcc gct cct	TCG	AAA	GCG		
spacer	Asu II				

s	.	.	.	
131	132	133	134	
TCT	TAA	TAG	TGA	

GGT|TAC|CAG|TCT|
 | BstE II |

| AAG|CCC|GCC|TAA|TGA|GCG|GGC|TTT|TTT|TTT|
 | Trp terminator |

| CCT|GAG|Gca|ggt|gag|cg - 3'
 | Sau I | spacer |

T06230-56036260

Table 27: DNA_synth1

5' | CCG | TCC | GTC | GGA | CCG | TAT | CCA | GGC | TTT | ACA | CTT | TAT |

| GCT | TCC | GGC | TCG | TAT | AAT | GTG | TGG |

AAT|TGT|GAG|CGG|ATA|ACA|ATT|
olig#4 = 3'-gt taa

CCT AGG
gga tcc

/ 3' = olig#3
 |GCC|GCT|CCT|TCG|AAA|GCG|
 cgg cga gga agc ttt cgc

|TCT|TAA|TAG|TGA|GGT|TAC|CAG|TCT|
aga att atc act cca atg gtc aga

|AAG|CCC|GCC|TAA|TGA|GCG|GGC|TTT|TTT|TTT|
ttc ggg cgg att act cgc ccg aaa aaa aaa

|CCT|GAG|GCA|GGT|GAG|CG
gga ctc cgt cca ctc gc - 5'

```
"Top" strand      99
"Bottom" strand   100
Overlap           23  (14 c/g and 9 a/t)
Net length        158
```

Table 28: DNA_seq2

5'- gca|cca|acg|
spacer

CCT|AGG|AGG|CTC|ACT|
Avr II
S. D.

m	k	k	s	l	v	l	k	a	s
1	2	3	4	5	6	7	8	9	10
ATG	AAG	AAA	TCT	CTG	GTT	CTT	AAG	GCT	AGC
						<u>Afl II</u>		<u>Nhe I</u>	

v	a	v	a	t	l	v	p	m	l
11	12	13	14	15	16	17	18	19	20
GTT	GCT	GTC	GCG	ACC	CTG	GTA	CCG	ATG	CTG
<u>Nru I</u>					<u>Kpn I</u>				

s	f	a	r	p	d	f	c	l	e
21	22	23	24	25	26	27	28	29	30
TCT	TTT	GCT	CGT	CCG	GAT	TTC	TGT	CTC	GAG
<u>AccIII</u>								<u>Ava I</u>	
								<u>Xho I</u>	

p	p	y	t	g	p	c	k	a	r
31	32	33	34	35	36	37	38	39	40
CCG	CCA	TAT	ACT	GGG	CCC	TGC	AAA	GCG	CGC
<u>PflM I</u>					<u>BssH II</u>				
					<u>Apa I</u>				
					<u>Dra II</u>				
					<u>Pss I</u>				

i	i	r
41	42	43
atc	atc	cgt

t	s	k
127	128	129
ACT	TCG	AAa
<u>gcg gct gcg </u>		
<u>Asu II</u> <u>spacer</u>		

- 3'

106230 5509550

Table 29: DNA_synth2

5'- GCA|CCA|ACG|CCT|AGG|AGG|CTC|ACT|ATG|AAG|AAA|TCT|CTG|GTT|CTT|AAG|GCT|AGC|GTT|GCT|GTC|GCG|ACC|CTG|GTA|CCG|ATG|CTG|
olig#6 = 3'- ggc tac gac/ 3' = olig#5
TCT|TTT|GCT|CGT|CCG|GAT|TTC|TGT|CTC|GAG|
aga aaa cga gca ggc cta aag aca gag ctcCCG|CCA|TAT|ACT|GGG|CCC|TGC|AAA|GCG|CGC|
ggc ggt ata tga ccc ggg acg ttt cgc gcgATC|ATC|CGT|
tag tag gcaACT|TCG|AAA|GCG|GCT|GCG|
tga agc ttt cgc cga cgc - 5'

"Top" strand	99
"Bottom" strand	99
Overlap	24 (14 c/g and 10 a/t)
Net length	155

Table 30: DNA_seq3

			a	r
			39	40
5'-	ccc tgc aca	GCG	CGC	
	spacer	BssH	II	

i	i	r	y	f	y	n	a	k
41	42	43	44	45	46	47	48	49
ATC	ATC	CGT	TAT	TTC	TAC	AAC	GCT	AAA

a	g	l	c	q	t	f	v	y	g	g
50	51	52	53	54	55	56	57	58	59	60
GCA	GGC	CTG	TGC	CAG	ACC	TTT	GTA	TAC	GGT	GGT
Stu I			Acc I			Xca I				

c	r	a	k	r	n	n	f	k
61	62	63	64	65	66	67	68	69
TGC	CGT	GCT	AAG	CGT	AAC	AAC	TTT	AAA
Esp I								

s	a	e	d	c	m	r	t	c	g
70	71	72	73	74	75	76	77	78	79
TCG	GCC	GAA	GAT	TGC	ATG	CGT	ACC	TGC	GGT
XmaIII			Sph I						

g	a
80	81
GGC	GCC
Bbe I	gct gaa
spacer	
Nar I	

	t	s	k
	127	128	129
ttt	acT	TCG	AAa
gcg tcg ccg - 3'			
Asu II			

T06250-55555

Table 31: DNA_synth3

5' - CCC TGC ACA GCG CGC

ATC | ATC | CGT | TAT | TTC | TAC | AAC | GCT | AAA |

olig#8 = 3'-g cca cca

/ 3' = olig#7
 |TGC|CGT|GCT|AAG|CGT|AAC|AAC|TTT|AAA|
 |acg|gca|cga|ttc|gca|ttg|ttg|aaa|ttt|

|TCG|GCC|GAA|GAT|TGC|ATG|CGT|ACC|TGC|GGT|
agc cgg ctt cta acg tac gca tgg acg cca

| GGC | GCC | GCT | GAA |
ccg cgg cgt ctt

|TTT|ACT|TCG|AAA|GCG|TCG|CCG|
aaa tga agc ttt cgc agc ggc -5'

```
"Top" strand      93
"Bottom" strand   97
Overlap           25 (15 g/c & 10 a/t)
Net length        146
```

Table 32: DNA_seq4

5'			g	a	a	e	g	d	d	
	80	81	82	83	84	85	86			
cct	cg	cct	GGC	GCC	GCT	GAA	GGT	GAT	GAT	
spacer			Bbe I							
			Nar I							

p	a	k	a	a	
87	88	89	90	91	
CCG	GCC	AAA	GCG	GCC	
Sfi I					

f	n	s	l	q	a	s	a	t	
92	93	94	95	96	97	98	99	100	
TTT	AAC	TCT	CTG	CAA	GCT	TCT	GCT	ACC	
Hind 3									

e	y	i	g	y	a	w	
101	102	103	104	105	106	107	
GAA	TAT	ATC	GGT	TAC	GCG	TGG	
Mlu I							

a	m	v	v	v	
108	109	110	111	112	
GCC	ATG	GTG	GTG	GTT	
BstX I					
Nco I					

i	v	g	a	t	i	g	i	
113	114	115	116	117	118	119	120	
ATC	GTT	GGT	GCT	ACC	ATC	GGT	ATC	

k	l	f	k	k	f	t	s	k	
121	122	123	124	125	126	127	128	129	
AAA	CTG	TTT	AAG	AAA	TTT	ACT	TCG	AAa	gcg tcg ggc
Asu II									spacer
									- 3'

T05230"56056360

Table 33: DNA_synth4

5' |GCT|CGC|CCT|GGC|GCC|GCT|GAA|GGT|GAT|GAT|

|CCG|GCC|AAA|GCG|GCC|

|TTT|AAC|TCT|CTG|CAA|GCT|TCT|GCT|ACC|

olig#10 = 3'- |GAA|TAT|ATC|GGT|TAC|GCG|TGG|
ata tag cca atg cgc acc

/ 3' = olig#9
|GCC|ATG|GTG|GTG|GTT|
cgg tac cac cac caa

|ATC|GTT|GGT|GCT|ACC|ATC|GGT|ATC|
tag caa cca cga tgg tag cca tag

|AAA|CTG|TTT|AAG|AAA|TTT|ACT|TCG|AAA|GCG|TCT|TGA|
ttt gac aaa ttc ttt aaa tga agc ttt cgc aga act - 5'

"Top" strand	100	
"Bottom" strand	93	
Overlap	25	(14 c/g and 11 a/t)
Net length	149	

T06230" 56056850

Table 34: Some interaction sets in BPTI

Res.	Number									
#	Diff.	AAs	Contents	BPTI	1	2	3	4	5	
-5	2		D -32	-						
-4	2		E -32	-						
-3	5		T P F Z -29	-						
-2	10		Z3 R3 Q2 T2 H G L K E -18	-						
-1	10		D4 T2 P2 Q2 E G N K R -18	-						
1	10		R21 A2 K2 H2 P L I T G D	R						5
2	9		P20 R4 A2 H2 N E V F L	P				s	5	
3	10		D15 K6 T3 R2 P2 S Y G A L	D				4	s	
4	7		F19 D4 L3 Y2 I2 A2 S	F				s	5	
5	1		C33	C				x	x	
6	10		L11 E5 N4 K3 Q2 I2 Y2 D2 T R	L				4		
7	5		L18 E11 K2 S Q	E			s	4		
8	7		P26 H2 A2 I L G F	P			3	4		
9	9		P17 A6 V3 R2 Q L K Y F	P		s	3	4		
10	10		Y11 E7 D4 A2 N2 R2 V2 S I D	Y	s		s	4		
11	10		T17 P5 A3 R2 I S Q Y V K	T	1	s	3	4		
12	2		G32 K	G	x		x	x		
13	5		P22 R6 L3 N I	P	1		s	4	s	
14	3		C31 T A	C	1		s	s	5	
15	12		K15 R4 Y2 M2 L2 -2 V G A I N F	K	1	s	3	4	s	
16	7		A22 G5 Q2 R K D F	A	1	s	s	s	5	
17	12		R12 K5 A2 Y3 H2 S2 F2 L M T G P	R	1	2	3		s	
18	6		I21 M4 F3 L2 V2 T	I	1	s	s		5	
19	7		I11 P10 R6 S2 K2 L Q	I	1	2	3		s	
20	5		R19 A7 S4 L2 Q	R	s	s	s		5	
21	4		Y18 F13 W I	Y		2	s	s	s	
22	6		F14 Y14 H2 A N S	F		s	3	4		
23	2		Y32 F	Y			s	s		
24	4		N26 K3 D3 S	N		s	3			
25	10		A12 S5 Q3 P3 W3 L2 T2 K G R	A			s	s		
26	9		K16 A6 T2 E2 S2 R2 G H V	K		s	3	4		
27	5		A18 S8 K3 L2 T2	A		2	3	4		
28	7		G13 K10 N5 Q2 R H M	G		2	s	s		
29	10		L9 Q7 K7 A2 F2 R2 M G T N	L		2	3			
30	1		C33	C		x	x	x		
31	7		Q12 E11 L4 K2 V2 Y N	Q		2	3	4		
32	11		T12 P5 K4 Q3 E2 L2 G V S R A	T		2	3	s		
33	1		F33	F	x	x	x	x		
34	11		V11 I8 T3 D2 N2 Q2 F H P R K	V	1	2	3	s		
35	2		Y31 W2	Y	s	s	s		5	
36	3		G27 S5 R	G	1					
37	1		G33	G	x				x	
38	3		C31 T A	C	1			s	5	
39	7		R13 G9 K4 Q3 D2 P M	R	1			4	s	

FO6250-56096350

Table 35:
Distances from C_β to
Tip of Side Group
in Å

Amino Acid type	Distance
A	0.0
C (reduced)	1.8
D	2.4
E	3.5
F	4.3
G	-
H	4.0
I	2.5
K	5.1
L	2.6
M	3.8
N	2.4
P	2.4
Q	3.5
R	6.0
S	1.5
T	1.5
V	1.5
W	5.3
Y	5.7

Notes: These distances were calculated for standard model parts with all side groups fully extended.

Table 36: Distances, BPTI residue set #2
 Distances in Å between C_β
 Hypothetical C_β was added to each Glycine.

	R17	I19	Y21	A27	G28	L29	Q31	T32	V34	A48
I19	7.7									
Y21	15.1	8.4								
A27	22.6	17.1	12.2							
G28	26.6	20.4	13.8	5.3						
L29	22.5	15.8	9.6	5.1	5.2					
Q31	16.1	10.4	6.8	6.8	10.6	6.8				
T32	11.7	5.2	6.1	12.0	15.5	10.9	5.4			
V34	5.6	6.5	11.6	17.6	21.7	18.0	11.4	8.2		
A48	18.5	11.0	5.4	12.6	13.3	8.4	8.8	8.3	15.7	
E49	22.0	14.7	8.9	16.9	16.1	12.2	13.9	13.3	19.8	5.5
M52	23.6	16.3	8.6	12.2	10.3	7.6	11.3	13.2	20.0	6.2
P9	14.0	11.3	9.0	12.2	15.4	13.3	7.9	9.2	8.7	13.9
T11	9.5	11.2	13.5	18.8	22.5	19.8	13.5	12.1	5.7	18.5
K15	7.9	14.6	20.1	27.4	31.3	27.9	21.4	18.1	10.3	24.6
A16	5.5	10.1	15.9	25.2	28.5	24.6	18.6	14.5	8.6	19.8
I18	6.1	6.0	11.2	21.3	24.4	20.2	14.7	10.4	7.0	15.0
R20	10.6	5.9	5.4	16.0	18.5	14.6	9.8	6.9	7.8	10.2
F22	15.6	10.9	5.6	10.5	12.8	10.3	6.2	8.1	10.8	10.3
N24	19.9	14.7	9.4	4.1	7.3	6.1	4.8	10.0	14.7	11.4
K26	24.4	20.1	15.2	5.4	7.7	9.8	10.1	15.3	19.0	17.0
C30	18.9	12.1	4.6	8.8	9.5	5.3	5.9	8.2	14.9	4.9
F33	10.8	7.4	7.7	12.6	16.4	13.0	6.6	5.6	5.5	12.2
Y35	8.4	7.4	9.4	18.4	21.4	17.9	12.2	9.5	5.8	14.4
S47	17.6	10.6	6.6	17.3	17.9	13.4	12.6	10.4	15.9	5.3
D50	20.0	13.6	7.2	17.2	16.8	13.5	13.5	12.9	17.6	7.6
C51	18.9	12.2	4.0	12.1	12.2	8.8	8.8	9.7	15.3	5.4
R53	25.4	18.6	11.0	17.2	15.0	13.0	15.7	16.7	22.3	9.7
R39	15.4	16.9	17.1	24.9	27.2	24.9	20.1	18.7	13.8	22.3

Distances in Å between C_β.

E49 M52 P9 T11 K15 A16 I18 R20 F22 N24

	E49	M52	P9	T11	K15	A16	I18	R20	F22	N24
<u>M52</u>	<u>6.1</u>									
P9	17.7	15.5								
T11	22.1	21.5	7.2							
K15	27.5	28.7	16.4	9.5						
A16	22.2	24.2	14.9	9.8	6.2					
I18	17.4	19.5	12.2	9.5	10.4	4.9				
R20	13.0	13.8	8.0	9.4	14.9	10.6	6.2			
F22	13.8	11.4	4.1	10.6	19.1	16.3	12.7	6.9		
N24	15.6	11.2	8.4	15.3	24.1	21.9	18.2	12.7	6.6	
K26	20.9	15.7	12.1	18.6	27.9	26.6	23.3	18.1	11.6	5.9
C30	8.7	5.6	10.6	16.6	24.1	20.2	15.7	9.8	6.8	6.9
F33	16.5	15.4	4.2	7.1	15.0	12.8	9.6	6.1	5.6	9.3
Y35	17.2	17.8	7.8	5.8	11.0	7.6	4.9	4.3	8.8	14.8
S47	4.7	9.1	15.3	18.5	23.1	17.6	12.8	9.1	12.0	15.3
D50	5.5	7.7	14.7	18.6	24.2	19.2	14.7	9.9	11.0	14.7
C51	7.1	5.4	11.0	16.4	23.5	19.2	14.6	8.7	6.9	9.6
R53	6.3	5.6	17.9	23.1	29.6	24.8	20.3	15.0	13.8	15.5
R39	23.9	24.0	13.0	9.5	12.0	11.8	12.5	12.8	14.7	20.8
	K26	C30	F33	Y35	S47	D50	C51	R53		
C30	12.4									
F33	13.9	10.1								
Y35	19.5	13.5	6.4							
S47	21.0	8.8	13.5	13.2						
D50	20.1	8.6	14.3	13.7	5.0					
C51	15.0	3.7	10.9	12.5	6.9	5.2				
R53	19.9	9.9	18.2	18.8	9.4	5.8	7.4			
R39	24.3	20.6	14.4	9.6	20.4	19.0	18.8	23.4		

Table 37: vgDNA to vary BPTI set #2.1

										208

Table 38: Result of varying set#2 of BPTI 2.1

l	e
29	30
CTC	GAG
Ava I	
Xho I	

178

p	p	y	t	g	p	c	k	a	D
31	32	33	34	35	36	37	38	39	40
CCG	CCA	TAT	ACT	GGG	CCC	TGC	AAA	GCG	GAT
PflM I									
				Apa I					
				Dra II					
				Pss I					

208

i	Q	r	y	f	y	n	a	k
41	42	43	44	45	46	47	48	49
ATC	CAG	CGT	TAT	TTC	TAC	AAC	GCT	AAA

235

E	g	L	c	q	t	f	S	y	g	g
50	51	52	53	54	55	56	57	58	59	60
GAG	GGC	CTG	TGC	CAG	ACC	TTT	TCG	TAC	GGT	GGT

268

c	r	a	k	r	n	n	f	k
61	62	63	64	65	66	67	68	69
TGC	CGT	GCT	AAG	CGT	AAC	AAC	TTT	AAA
Esp I								

295

s	W	e	d	c	m	r	t	c	g
70	71	72	73	74	75	76	77	78	79
TCG	TGG	GAA	GAT	TGC	ATG	CGT	ACC	TGC	GGT
Sph I									

325

g	a
80	81
GGC	GCC
Bbe I	
Nar I	

T06250"55035550

Table 40: Result of varying set#2 of BPTI 2.2

										<table><tr><td>l</td><td>e</td></tr><tr><td>29</td><td>30</td></tr><tr><td>CTC</td><td>GAG</td></tr><tr><td>Xho</td><td>I</td></tr></table>	l	e	29	30	CTC	GAG	Xho	I	178																									
l	e																																											
29	30																																											
CTC	GAG																																											
Xho	I																																											
<table><tr><td>p</td><td>p</td><td>y</td><td>t</td><td>g</td><td>p</td><td>c</td><td>E</td><td>a</td><td>D</td></tr><tr><td>31</td><td>32</td><td>33</td><td>34</td><td>35</td><td>36</td><td>37</td><td>38</td><td>39</td><td>40</td></tr><tr><td>CCG</td><td>CCA</td><td>TAT</td><td>ACT</td><td>GGG</td><td>CCC</td><td>TGC</td><td>GAG</td><td>GCG</td><td>GAT</td></tr></table>	p	p	y	t	g	p	c	E	a	D	31	32	33	34	35	36	37	38	39	40	CCG	CCA	TAT	ACT	GGG	CCC	TGC	GAG	GCG	GAT	<u>PflM I</u>										208			
p	p	y	t	g	p	c	E	a	D																																			
31	32	33	34	35	36	37	38	39	40																																			
CCG	CCA	TAT	ACT	GGG	CCC	TGC	GAG	GCG	GAT																																			
				<u>Apa I</u>																																								
<table><tr><td>V</td><td>Q</td><td>N</td><td>F</td><td>f</td><td>y</td><td>n</td><td>a</td><td>k</td></tr><tr><td>41</td><td>42</td><td>43</td><td>44</td><td>45</td><td>46</td><td>47</td><td>48</td><td>49</td></tr><tr><td>GTT</td><td>CAG</td><td>AAT</td><td>TTT</td><td>TTC</td><td>TAC</td><td>AAC</td><td>GCT</td><td>AAA</td></tr></table>	V	Q	N	F	f	y	n	a	k	41	42	43	44	45	46	47	48	49	GTT	CAG	AAT	TTT	TTC	TAC	AAC	GCT	AAA									235								
V	Q	N	F	f	y	n	a	k																																				
41	42	43	44	45	46	47	48	49																																				
GTT	CAG	AAT	TTT	TTC	TAC	AAC	GCT	AAA																																				
<table><tr><td>E</td><td>F</td><td>L</td><td>c</td><td>S</td><td>A</td><td>f</td><td>S</td><td>y</td><td>g</td><td>g</td></tr><tr><td>50</td><td>51</td><td>52</td><td>53</td><td>54</td><td>55</td><td>56</td><td>57</td><td>58</td><td>59</td><td>60</td></tr><tr><td>GAG</td><td>TTT</td><td>CTG</td><td>TGC</td><td>TCT</td><td>GCT</td><td>TTT</td><td>TCG</td><td>TAC</td><td>GGT</td><td>GGT</td></tr></table>	E	F	L	c	S	A	f	S	y	g	g	50	51	52	53	54	55	56	57	58	59	60	GAG	TTT	CTG	TGC	TCT	GCT	TTT	TCG	TAC	GGT	GGT											268
E	F	L	c	S	A	f	S	y	g	g																																		
50	51	52	53	54	55	56	57	58	59	60																																		
GAG	TTT	CTG	TGC	TCT	GCT	TTT	TCG	TAC	GGT	GGT																																		
<table><tr><td>c</td><td>r</td><td>a</td><td>k</td><td>r</td><td>n</td><td>n</td><td>f</td><td>k</td></tr><tr><td>61</td><td>62</td><td>63</td><td>64</td><td>65</td><td>66</td><td>67</td><td>68</td><td>69</td></tr><tr><td>TGC</td><td>CGT</td><td>GCT</td><td>AAG</td><td>CGT</td><td>AAC</td><td>AAC</td><td>TTT</td><td>AAA</td></tr></table>	c	r	a	k	r	n	n	f	k	61	62	63	64	65	66	67	68	69	TGC	CGT	GCT	AAG	CGT	AAC	AAC	TTT	AAA	<u>Esp I</u>									295							
c	r	a	k	r	n	n	f	k																																				
61	62	63	64	65	66	67	68	69																																				
TGC	CGT	GCT	AAG	CGT	AAC	AAC	TTT	AAA																																				
<table><tr><td>s</td><td>W</td><td>Q</td><td>d</td><td>c</td><td>m</td><td>r</td><td>t</td><td>c</td><td>g</td></tr><tr><td>70</td><td>71</td><td>72</td><td>73</td><td>74</td><td>75</td><td>76</td><td>77</td><td>78</td><td>79</td></tr><tr><td>TCG</td><td>TGG</td><td>CAG</td><td>GAT</td><td>TGC</td><td>ATG</td><td>CGT</td><td>ACC</td><td>TGC</td><td>GGT</td></tr></table>	s	W	Q	d	c	m	r	t	c	g	70	71	72	73	74	75	76	77	78	79	TCG	TGG	CAG	GAT	TGC	ATG	CGT	ACC	TGC	GGT					<u>Sph I</u>						325			
s	W	Q	d	c	m	r	t	c	g																																			
70	71	72	73	74	75	76	77	78	79																																			
TCG	TGG	CAG	GAT	TGC	ATG	CGT	ACC	TGC	GGT																																			
<table><tr><td>g</td><td>a</td></tr><tr><td>80</td><td>81</td></tr><tr><td>GGC</td><td>GCC</td></tr><tr><td>Bbe</td><td>I</td></tr><tr><td>Nar</td><td>I</td></tr></table>	g	a	80	81	GGC	GCC	Bbe	I	Nar	I																																		
g	a																																											
80	81																																											
GGC	GCC																																											
Bbe	I																																											
Nar	I																																											

T05290-5609550

Table 42: Result of varying set#2 of BPTI 2.3

1	e	
29	30	
CTC	GAG	178
Ava I		
Xho I		

P	E	Y	Q	g	p	c	E	a	A
31	32	33	34	35	36	37	38	39	40
CCG	GAG	TAT	CAG	GGG	CCC	TGC	GAG	GCG	GCT
				Apa I					

V	Q	N	W	f	y	n	a	k	
41	42	43	44	45	46	47	48	49	
GTT	CAG	AAT	TGG	TTC	TAC	AAC	GCT	AAA	235

Q	F	M	C	S	L	f	H	Y	g	g
50	51	52	53	54	55	56	57	58	59	60
CAG	TTT	ATG	TGC	TCT	CTT	TTT	CAT	TAC	GGT	GGT

268

c	r	a	k	r	n	n	f	k	
61	62	63	64	65	66	67	68	69	
TGC	CGT	GCT	AAG	CGT	AAC	AAC	TTT	AAA	295
Esp I									

s	w	Q	d	c	m	r	t	c	g	
70	71	72	73	74	75	76	77	78	79	
TCG	TGG	CAG	GAT	TGC	ATG	CGT	ACC	TGC	GGT	325
Sph I										

g	a
80	81
GGC	GCC
Bbe I	
Nar I	

Table 50: Preferred IPBDs

IPBD	Number Amino Acids	Structure	Cross Links	Secreted	Source Organism	Afm
Aprotinin	58	X-ray, NMR	3 SS 5-55, 14-38 30-51 (1:6, 2:4, 3:5)	yes	<u>Bos taurus</u>	trypsin
Crambin	46	X-ray, NMR	3 SS	yes	rape seed	?, Mab
CMTI-III	26	NMR	3 SS	yes	cucumbers	trypsin
ST-IA	13	NMR	3 SS	yes	<u>E. coli</u> Mabs & guanylate cyclase	
Third domain, ovomucoid	56	X-ray, NMR	3 SS	yes	<u>Coturnix</u> <u>Coturnix</u> <u>japonica</u>	trypsin
Ribonuclease A	124	X-ray, NMR		yes	<u>Bos taurus</u>	RNA, DNA
Ribonuclease	104	X-ray, NMR?		yes	<u>A. oryzae</u>	RNA, DNA
Lysozyme	129	X-ray, NMR?	4 SS	yes	<u>Gallus gallus</u>	NAG-NAM-NAG
Azurin	128	X-ray	Cu:CYS, HIS, MET		<u>P. aeruginosa</u>	Mab

Table Characteristics of Known IPBDs
(continued)

IPBD	Number Amino Acids	Structure	Cross Links	Secreted	Source Organism	AfM
α -Conotoxins	13-15	NMR	2 SS	yes	<u>Conus</u> snails	Receptor
μ -Conotoxins	20-25	NMR	3 SS	yes	<u>Conus</u> snails	Receptor
Ω -Conotoxins	25-30	-	3 SS	yes	<u>Conus</u> snails	Receptor
King-kong peptides	25-30	-	3 SS	yes	<u>Conus</u> snails	Mabs
Nuclease (staphylococcal)	141	X-ray	none	yes	<u>S aurius</u>	RNA, DNA
Charybdotoxin (scorpion toxin)	37	NMR	3 SS 7-28, 13-33 17-35 (1:4, 2:5, 3:6)	yes	<u>Leiurus</u> <u>quinquestriatus</u> <u>hebraeus</u>	Ca ⁺² -dependent K ⁺ channel
Apamin (bee venom)	12	NMR	2SS (1:3, 2:4)	yes	Bees	Mabs, Receptor(?)

Table 50
(continued)

Other suitable IPBDs
Perredoxin
Secretory trypsin inhibitor
Soybean trypsin inhibitor
SLPI (Secretory Leukocyte Protease Inhibitor) (THOM86) and SPAI (ARAK90)
Cystatin and homologues (MACH89, STUB90)
Eglin (MCPH85)
Barley inhibitor (CLOR87a, CLOR87b, SVEN82)

5

Notes:

a Designed sequence contained AGGAGG, but sequencing
30 indicates that actual DNA contains AGAGG.

SECRET

Table 102a: Annotated Sequence of gene
found in pGEM-MB1

		nucleotide number																																																																										
5	5'-(G GATCC TCTAGA GTC) GGC- from pGEM polylinker	3																																																																										
10	<u>tttaca</u> CTTTATGCTTCCGGCTCG <u>tataat</u> GTGTGG- -35 <u>lacUV5</u> -10	39																																																																										
15	<u>aATTGTGAGCGcTcACAATT-</u> lacO-symm operator	59																																																																										
20	<u>gagctc</u> <u>AG(G)AGG</u> CttaCT- <u>SacI</u> Shine-Dalgarno seq. ^a	77																																																																										
25	<table><tr><td>fM</td><td>K</td><td>K</td><td>S</td><td>L</td><td>V</td><td>L</td><td>K</td><td>A</td><td>S</td></tr><tr><td>1</td><td>2</td><td>3</td><td>4</td><td>5</td><td>6</td><td>7</td><td>8</td><td>9</td><td>10</td></tr><tr><td>ATG</td><td>AAG</td><td>AAA</td><td>TCT</td><td>CTG</td><td>GTT</td><td>CTT</td><td>AAG</td><td>GCT</td><td>AGC</td></tr><tr><td colspan="6"></td><td><u>Afl II</u></td><td><u>Nhe I</u></td><td colspan="2"></td></tr></table>	fM	K	K	S	L	V	L	K	A	S	1	2	3	4	5	6	7	8	9	10	ATG	AAG	AAA	TCT	CTG	GTT	CTT	AAG	GCT	AGC							<u>Afl II</u>	<u>Nhe I</u>			107																																		
fM	K	K	S	L	V	L	K	A	S																																																																			
1	2	3	4	5	6	7	8	9	10																																																																			
ATG	AAG	AAA	TCT	CTG	GTT	CTT	AAG	GCT	AGC																																																																			
						<u>Afl II</u>	<u>Nhe I</u>																																																																					
30	<table><tr><td>V</td><td>A</td><td>V</td><td>A</td><td>T</td><td>L</td><td>V</td><td>P</td><td>M</td><td>L</td></tr><tr><td>11</td><td>12</td><td>13</td><td>14</td><td>15</td><td>16</td><td>17</td><td>18</td><td>19</td><td>20</td></tr><tr><td>GTT</td><td>GCT</td><td>GTC</td><td>GCG</td><td>ACC</td><td>CTG</td><td>GTA</td><td>CCT</td><td>ATG</td><td>TTG</td></tr><tr><td colspan="4"></td><td><u>Nru I</u></td><td colspan="4"></td><td><u>Kpn I</u></td><td></td></tr></table>	V	A	V	A	T	L	V	P	M	L	11	12	13	14	15	16	17	18	19	20	GTT	GCT	GTC	GCG	ACC	CTG	GTA	CCT	ATG	TTG					<u>Nru I</u>					<u>Kpn I</u>		137																																	
V	A	V	A	T	L	V	P	M	L																																																																			
11	12	13	14	15	16	17	18	19	20																																																																			
GTT	GCT	GTC	GCG	ACC	CTG	GTA	CCT	ATG	TTG																																																																			
				<u>Nru I</u>					<u>Kpn I</u>																																																																			
35	<table><tr><td>S</td><td>F</td><td>A</td><td>R</td><td>P</td><td>D</td><td>F</td><td>C</td><td>L</td><td>E</td></tr><tr><td>21</td><td>22</td><td>23</td><td>24</td><td>25</td><td>26</td><td>27</td><td>28</td><td>29</td><td>30</td></tr><tr><td>TCC</td><td>TTC</td><td>GCT</td><td>CGT</td><td>CCG</td><td>GAT</td><td>TTC</td><td>TGT</td><td>CTC</td><td>GAG</td></tr><tr><td colspan="6"></td><td><u>AccIII</u></td><td colspan="2"></td><td><u>Ava I</u></td></tr></table>	S	F	A	R	P	D	F	C	L	E	21	22	23	24	25	26	27	28	29	30	TCC	TTC	GCT	CGT	CCG	GAT	TTC	TGT	CTC	GAG							<u>AccIII</u>			<u>Ava I</u>	167																																		
S	F	A	R	P	D	F	C	L	E																																																																			
21	22	23	24	25	26	27	28	29	30																																																																			
TCC	TTC	GCT	CGT	CCG	GAT	TTC	TGT	CTC	GAG																																																																			
						<u>AccIII</u>			<u>Ava I</u>																																																																			
40	M13/BPTI Jnct	<u>Xho I</u>																																																																										
45	<table><tr><td>P</td><td>P</td><td>Y</td><td>T</td><td>G</td><td>P</td><td>C</td><td>K</td><td>A</td><td>R</td></tr><tr><td>31</td><td>32</td><td>33</td><td>34</td><td>35</td><td>36</td><td>37</td><td>38</td><td>39</td><td>40</td></tr><tr><td>CCA</td><td>CCA</td><td>TAC</td><td>ACT</td><td>GGG</td><td>CCC</td><td>TGC</td><td>AAA</td><td>GCG</td><td>CGC</td></tr><tr><td colspan="5"></td><td><u>PflM I</u></td><td colspan="4"></td><td><u>BssH II</u></td></tr><tr><td colspan="5"></td><td colspan="2"></td><td><u>Apa I</u></td><td colspan="2"></td><td></td></tr><tr><td colspan="5"></td><td colspan="2"></td><td><u>Dra II</u></td><td colspan="2"></td><td></td></tr><tr><td colspan="5"></td><td colspan="2"></td><td><u>Pss I</u></td><td colspan="2"></td><td></td></tr></table>	P	P	Y	T	G	P	C	K	A	R	31	32	33	34	35	36	37	38	39	40	CCA	CCA	TAC	ACT	GGG	CCC	TGC	AAA	GCG	CGC						<u>PflM I</u>					<u>BssH II</u>								<u>Apa I</u>											<u>Dra II</u>											<u>Pss I</u>				197
P	P	Y	T	G	P	C	K	A	R																																																																			
31	32	33	34	35	36	37	38	39	40																																																																			
CCA	CCA	TAC	ACT	GGG	CCC	TGC	AAA	GCG	CGC																																																																			
					<u>PflM I</u>					<u>BssH II</u>																																																																		
							<u>Apa I</u>																																																																					
							<u>Dra II</u>																																																																					
							<u>Pss I</u>																																																																					

5

10

152025

30

35404550

5

437

K	L	F	K	K	F	T	S	K	A
121	122	123	124	125	126	127	128	129	130
AAA	CTG	TTC	AAG	AAG	TTT	ACT	TCG	AAG	GCG

Asu II

467

S	.	.	.
131	132	133	134
TCT	TAA	TGA	TAG

GGTTACC-
BstE II

486

AGTCTA AGCCCGC CTAATGA GCGGGCT TTTTTTTT-
terminator

521

aTCGA (GACctgcagggcatgc)-3'
(SalI) from pGEM polylinker

30 Notes:

^a Designed called for Shine-Dalgarno sequence, AGGAGG, but sequencing shows that actual constructed gene contains AGAGG.

35

Note the following enzyme equivalences,

Xma III = Eag I Acc III = BspM II
Dra II = EcoO109 I Asu II = BstB I

40

Table 102b : Annotated Sequence of gene
after insertion of SalI linker

5	5'-(GGATCC TCTAGA GTC) GGC- from pGEM polylinker	nucleotide number	3																																																																										
10	<u>tttaca</u> CTTTATGCTTCCGGCTCG <u>tataat</u> GTGTGG- -35 <u>lacUV5</u> -10		39																																																																										
15	<u>aATTGTGAGCGcTcACAATT-</u> lacO-symm operator		59																																																																										
20	<u>gagctc</u> <u>AGAGG</u> CttaCT- <u>SacI</u> Shine-Dalgarno seq.		77																																																																										
25	<table><tr><td>fM</td><td>K</td><td>K</td><td>S</td><td>L</td><td>V</td><td>L</td><td>K</td><td>A</td><td>S</td></tr><tr><td>1</td><td>2</td><td>3</td><td>4</td><td>5</td><td>6</td><td>7</td><td>8</td><td>9</td><td>10</td></tr><tr><td>ATG</td><td>AAG</td><td>AAA</td><td>TCT</td><td>CTG</td><td>GTT</td><td>CTT</td><td>AAG</td><td>GCT</td><td>AGC</td></tr><tr><td colspan="6"></td><td><u>Afl II</u></td><td><u>Nhe I</u></td><td colspan="2"></td></tr></table>	fM	K	K	S	L	V	L	K	A	S	1	2	3	4	5	6	7	8	9	10	ATG	AAG	AAA	TCT	CTG	GTT	CTT	AAG	GCT	AGC							<u>Afl II</u>	<u>Nhe I</u>				107																																		
fM	K	K	S	L	V	L	K	A	S																																																																				
1	2	3	4	5	6	7	8	9	10																																																																				
ATG	AAG	AAA	TCT	CTG	GTT	CTT	AAG	GCT	AGC																																																																				
						<u>Afl II</u>	<u>Nhe I</u>																																																																						
30	<table><tr><td>V</td><td>A</td><td>V</td><td>A</td><td>T</td><td>L</td><td>V</td><td>P</td><td>M</td><td>L</td></tr><tr><td>11</td><td>12</td><td>13</td><td>14</td><td>15</td><td>16</td><td>17</td><td>18</td><td>19</td><td>20</td></tr><tr><td>GTT</td><td>GCT</td><td>GTC</td><td>GCG</td><td>ACC</td><td>CTG</td><td>GTA</td><td>CCT</td><td>ATG</td><td>TTG</td></tr><tr><td colspan="4"></td><td><u>Nru I</u></td><td colspan="4"></td><td><u>Kpn I</u></td><td></td></tr></table>	V	A	V	A	T	L	V	P	M	L	11	12	13	14	15	16	17	18	19	20	GTT	GCT	GTC	GCG	ACC	CTG	GTA	CCT	ATG	TTG					<u>Nru I</u>					<u>Kpn I</u>			137																																	
V	A	V	A	T	L	V	P	M	L																																																																				
11	12	13	14	15	16	17	18	19	20																																																																				
GTT	GCT	GTC	GCG	ACC	CTG	GTA	CCT	ATG	TTG																																																																				
				<u>Nru I</u>					<u>Kpn I</u>																																																																				
35	<table><tr><td>S</td><td>F</td><td>A</td><td>R</td><td>P</td><td>D</td><td>F</td><td>C</td><td>L</td><td>E</td></tr><tr><td>21</td><td>22</td><td>23</td><td>24</td><td>25</td><td>26</td><td>27</td><td>28</td><td>29</td><td>30</td></tr><tr><td>TCC</td><td>TTC</td><td>GCT</td><td>CGT</td><td>CCG</td><td>GAT</td><td>TTC</td><td>TGT</td><td>CTC</td><td>GAG</td></tr><tr><td colspan="6"></td><td><u>AccIII</u></td><td><u>Ava I</u></td><td colspan="2"></td></tr></table>	S	F	A	R	P	D	F	C	L	E	21	22	23	24	25	26	27	28	29	30	TCC	TTC	GCT	CGT	CCG	GAT	TTC	TGT	CTC	GAG							<u>AccIII</u>	<u>Ava I</u>				167																																		
S	F	A	R	P	D	F	C	L	E																																																																				
21	22	23	24	25	26	27	28	29	30																																																																				
TCC	TTC	GCT	CGT	CCG	GAT	TTC	TGT	CTC	GAG																																																																				
						<u>AccIII</u>	<u>Ava I</u>																																																																						
40	M13/BPTI Jnct		<u>Xho I</u>																																																																										
45	<table><tr><td>P</td><td>P</td><td>Y</td><td>T</td><td>G</td><td>P</td><td>C</td><td>K</td><td>A</td><td>R</td></tr><tr><td>31</td><td>32</td><td>33</td><td>34</td><td>35</td><td>36</td><td>37</td><td>38</td><td>39</td><td>40</td></tr><tr><td>CCA</td><td>CCA</td><td>TAC</td><td>ACT</td><td>GGG</td><td>CCC</td><td>TGC</td><td>AAA</td><td>GCG</td><td>CGC</td></tr><tr><td colspan="4"></td><td><u>PflM I</u></td><td colspan="4"></td><td><u>BssH II</u></td><td></td></tr><tr><td colspan="4"></td><td colspan="2"><u>Apa I</u></td><td colspan="4"></td><td></td></tr><tr><td colspan="4"></td><td colspan="2"><u>Dra II</u></td><td colspan="4"></td><td></td></tr><tr><td colspan="4"></td><td colspan="2"><u>Pss I</u></td><td colspan="4"></td><td></td></tr></table>	P	P	Y	T	G	P	C	K	A	R	31	32	33	34	35	36	37	38	39	40	CCA	CCA	TAC	ACT	GGG	CCC	TGC	AAA	GCG	CGC					<u>PflM I</u>					<u>BssH II</u>						<u>Apa I</u>											<u>Dra II</u>											<u>Pss I</u>								197
P	P	Y	T	G	P	C	K	A	R																																																																				
31	32	33	34	35	36	37	38	39	40																																																																				
CCA	CCA	TAC	ACT	GGG	CCC	TGC	AAA	GCG	CGC																																																																				
				<u>PflM I</u>					<u>BssH II</u>																																																																				
				<u>Apa I</u>																																																																									
				<u>Dra II</u>																																																																									
				<u>Pss I</u>																																																																									

Table 102b : Annotated Sequence
of gene after insertion of SalI linker
(continued)

5	I	I	R	Y	F	Y	N	A	K	A		
	41	42	43	44	45	46	47	48	49	50		
	ATC	ATC	CGC	TAT	TTC	TAC	AAT	GCT	AAA	GC	-	226
10	G	L	C	Q	T	F	V	Y	G	G		
	51	52	53	54	55	56	57	58	59	60		
	A GGC	CTG	TGC	CAG	ACC	TTT	GTA	TAC	GGT	GGT	-	257
	Stu I						Acc I					
15							Xca I					
20	C	R	A	K	R	N	N	F	K			
	61	62	63	64	65	66	67	68	69			
	TGC	CGT	GCT	AAG	CGT	AAC	AAC	TTT	AAA	-		284
			Esp I									
25	S	A	E	D	C	M	R	T	C	G		
	70	71	72	73	74	75	76	77	78	79		
	TCG	GCC	GAA	GAT	TGC	ATG	CGT	ACC	TGC	GGT	-	314
	XmaIII				Sph I							
	BPTI/M13 boundary											
30	G	A	A	E	G	D	D	P	A	K	A	A
	80	81	82	83	84	85	86	87	88	89	90	91
	GGC	GCC	GCT	GAA	GGT	GAT	GAT	CCG	GCC	AAG	GCG	GCC
	Bbe I								Sfi I			
	Nar I											
35												
40	F	N	S	L	Q	A	S	A	T			
	92	93	94	95	96	97	98	99	100			
	TTC	AAT	TCT	CTG	CAA	GCT	TCT	GCT	ACC	-		377
					Hind 3							
45	E	Y	I	G	Y	A	W					
	101	102	103	104	105	106	107					
	GAG	TAT	ATT	GGT	TAC	GCG	TGG	-				398
50	A	M	V	V	V	I	V	G	A			
	108	109	110	111	112	113	114	115	116			
	GCC	ATG	GTG	GTG	GTT	ATC	GTT	GGT	GCT	-		425
	BstX I											
	Nco I											

Table 102 : Annotated Sequence
of osp-ipbd gene
(continued)

5 Table 102c: Calculated properties of Peptide

For the apoprotein

	Molecular weight of peptide =	16192
10	Charge on peptide =	9
	[A+G+P] =	36
	[C+F+H+I+L+M+V+W+Y] =	48
	[D+E+K+R+N+Q+S+T+.] =	48

15 For the mature protein

	Molecular weight of peptide =	13339
	Charge on peptide =	6
	[A+G+P] =	31
20	[C+F+H+I+L+M+V+W+Y] =	37
	[D+E+K+R+N+Q+S+T+.] =	41

Table 102d: Codon Usage

25		Second Base				
	First Base	t	c	a	g	Third base
30	t	3	4	2	1	t
		5	1	4	5	c
		0	0	0	0	a
		1	2	0	1	g
35	c	1	1	0	4	t
		1	1	0	2	c
		0	2	1	0	a
		5	2	1	0	g
40	a	1	2	2	0	t
		5	5	2	1	c
		0	0	5	0	a
		4	0	7	0	g
45	g	4	9	4	6	t
		1	5	0	2	c
		2	1	2	0	a
		2	5	2	2	g

Table 102e: Amino-acid frequency

Encoded polypeptide								
5	AA	#	AA	#	AA	#	AA	#
	A	20	C	6	D	4	E	4
	F	8	G	10	H	0	I	6
	K	12	L	8	M	4	N	4
10	P	6	Q	2	R	6	S	8
	T	7	V	9	W	1	Y	6
	.	1						
Mature protein								
15	AA	#	AA	#	AA	#	AA	#
	A	16	C	6	D	4	E	4
	F	7	G	10	H	0	I	6
	K	9	L	4	M	2	N	4
20	P	5	Q	2	R	6	S	5
	T	6	V	5	W	1	Y	6

T06290" 55035550

Table 102f: Enzymes used to manipulate BPTI-gp8 fusion

	<u>SacI</u>	GAGCT C
	<u>AflII</u>	C TTAAG
5	<u>NheI</u>	G CTAGC
	<u>NruI</u>	TCG CGA
	<u>KpnI</u>	GGTAC C
	<u>AccIII</u> = <u>BspMII</u>	T CCGGA
	<u>AvaI</u>	C yCGrG
10	<u>XhoI</u>	C TCGAG
	<u>PflMI</u>	CCAnnnnn nTGG
	<u>BssHII</u>	G CGCGC
	<u>ApaI</u>	GGGCC C
	<u>DraII</u> = <u>Eco109I</u>	rGGnC Cy (Same as <u>PssI</u>)
15	<u>StuI</u>	AGG CCT
	<u>AccI</u>	GT mKAC
	<u>XcaI</u>	GTA TAC
	<u>EspI</u>	GC TnAGC
	<u>XmaIII</u>	C GGCCG (Supplier ?)
20	<u>SphI</u>	GCATG C
	<u>BbeI</u>	GGCGC C (Supplier ?)
	<u>NarI</u>	GGCG CC
	<u>SfiI</u>	GGCCnnnn nGGCC
	<u>HindIII</u>	A AGCTT
25	<u>BstXI</u>	CCAnnnnnn nTGG
	<u>NcoI</u>	C CATGG
	<u>AsuII</u> = <u>BstBI</u>	TT CGAA
	<u>BstEII</u>	G GTnACC
	<u>SalI</u>	G TCGAC

T06290"55055550

Table 103 : Annotated Sequence of osp-ipbd gene

Underscored bases indicate sites of overlap between annealed synthetic duplexes.

5

5'-

10 /GGC tttaga CTTTAT, GCTTCCGGCTCG tataat GTGTGG-
lacUV5

aATTGTGAGCGcTcACAATT-
lacO-symm operator

15

gagctc AG(G)/AGG Cttact-
Sac I Shine-Dalgarno seq.

20

25

fM	K	K	S	L	V	L	K	A	S
1	2	3	4	5	6	7	8	9	10
ATG	AAG,	AAA	TCT	CTG	GTT	CTT	AAG	GCT	AGC
							Afl II	Nhe I	-

25

30

V	A	V	A	T	L	V	P	M	L	
11	12	13	14	15	16	17	18	19	20	
GTT	GCT	GTC	GCG	ACC	CTG	GTA	CCT	ATG	T/TG	
				Nru I					Kpn I	-

30

35

S	F	A	R	P	D	F	C	L	E	
21	22	23	24	25	26	27	28	29	30	
TCC	TTC	GCT	CG,T	CCG	GAT	TTC	TGT	CTC	GAG	
				↑	AccIII					
				M13/BPTI Jnct					Ava I	-
									Xho I	

35

T06230"55095B60

Table 103 : Annotated Sequence
of osp-*ipbd* gene
(continued)

5

P	P	Y	T	G	P	C	K	A	R	
31	32	33	34	35	36	37	38	39	40	
CCA	CCA	TAC	ACT	GGG	CCC	TGC	AAA	GCG	CGC	-
PflM I						BssH II				

10

Apa I					
Dra II					
Pss I					

15

I	I	R	Y	F	Y	N	A	K	A	
41	42	43	44	45	46	47	48	49	50	
ATC	ATC	CG/C	TAT	TTC	TAC	AAT	GC, T	AAA	GC	-

20

G	L	C	Q	T	F	V	Y	G	G	
51	52	53	54	55	56	57	58	59	60	
A GGC	CTG	TGC	CAG	ACC	TTT	GTA	TAC	GGT	GGT	-
Stu I						Acc I				
						Xca I				

25

C	R	A	K	R	N	N	F	K		
61	62	63	64	65	66	67	68	69		
TGC	CGT	GCT	AAG	CGT	/AAC	AAC	TTT	AAA	-	
Esp I										

30

S	A	E	D	C	M	R	T	C	G	
70	71	72	73	74	75	76	77	78	79	
TCG,	GCC	GAA	GAT	TGC	ATG	CGT	ACC	TGC	GGT	-
Xma III					Sph I					

35

BPTI/M13 boundary

G	A	A	E	G	D	D	P	A	K	A	A	
80	81	82	83	84	85	86	87	88	89	90	91	
GGC	GCC	GCT	GAA	GGT	GAT	GAT	CCG	GCC	AAG	GCG	G/CC	-
Bbe I								Sfi I				
Nar I												

40

45

F	N	S	L	Q	A	S	A	T		
92	93	94	95	96	97	98	99	100		
TTC	AAT	TCT	CTG	C, AA	GCT	TCT	GCT	ACC	-	
Hind 3										

5

E	Y	I	G	Y	A	W
101	102	103	104	105	106	107
GAG	TAT	ATT	GGT	TAC	GCG	TGG -

A	M	V	V	V	I	V	G	A	
108	109	110	111	112	113	114	115	116	
GCC	ATG	GTG	GTG	GTT	AT/C	GTT	GGT	GCT	-

15

Nco I

T	I	G	I	
117	118	119	120	
ACC,	ATC	GGG	ATC	-

K	L	F	K	K	F	T	S	K	A
121	122	123	124	125	126	127	128	129	130
AAA	CTG	TTC	AAG	AAG	TTT	ACT	TCG	AAG	GCG

Asu II

25

S	.	.	.
131	132	133	134
TCT	TAA	TGA	TAG

GGTTA/CC-
BstE II

30

AGTCTA AGCCC,GC CTAATGA GCGGGCT TTTTTTTT-
terminator

35

a/ (TCGA), -3'
(Sal I)

40

Table 104: Definition and alignment of oligonucleotides

Lines ending with "-" are continued on a following line. Blocks of ten bases are delimited by "-" within a line. When a break in one strand does not correspond to a ten-base mark in the other strand, "--" is inserted in the other strand.

	↓ Olig #801 (68 bases)
5'-GG-CTTTACACTT-TAT--GCTTCCG-	
3'-cc-gaaatgtgaa-ata	cgaaggc-
	↑
	filled in
	↓ Olig #802 (67 bases)
GCTCGTATAA-TGTGTGGAAT-TGTGAGCGCT-CACAATTGAG-CTCAGG	AGGC-TTACTATGAA-
cgagcatatt-acacacctta-acactcgcga-gtgttaactc-gagtcc--tccg-aatgatactt-	
	↓ Olig #803 (70 bases)
G--AAATCTCTG-GTTCTTAAGG-CTAGCGTTGC-TGTCGCGGACC-CTGGTACCTA-TGT	TGTCCTT-
c tttagagac-caagaattcc-gatcgcaacg-acagcgcctgg-qaccatggat-aca--acagqaa-	
↑ Olig #817 (68 bases)	
CGCTCG--TCCG-GAATTCTGTC-TCGAGCCACC-ATACACTGGG-CCCTGCAAG-CGCCGCATCAT-	
gcgagc	aggc-ctaaagacag-agctcgtggtg-tatgtgaccc-gggacgtttc-gcgcgtagta-
	↑ Olig #816 (65 bases)

Table 104: Definition and alignment of oligonucleotides
(continued)

↓ Olig #804 (67 bases)
CCG CTATTTC-TACAATGC--TA-AAGCAGGCCT-GTGCCAGACC-TTTGTATACG-GTGGTTGCCG-
ggc--gataaag-atgttacg at-ttcgtccgga-cacggtctgg-aaacatatgc-caccaacggc-
↑ Olig #815 (72 bases)

↓ Olig #805 (76 bases)
TGCTAAGCGT AACAACTTTA-AATCG--GCCGA-AGATTGCATG-CGTACCTGCG-GTGGCGCCGC-
acgatccgca--ttgttgaat-ttagc cggct-tctaacgtac-gcatggacgc-caccgcggcg-
↑ Olig #814 (67 bases)

↓ Olig #806 (67 bases)
TGAAGGTGAT-GATCCGGCCA-AGGCGG CCTT-CAATTCTCTG-C--AAGCTTCTG-CTACCGAGTA-
acttccacta-ctaggccggt-tccgac--ggaa-gttaagagac-g ttcgaaagac-gatggctcat-
↑ Olig #813 (76 bases)

↓ Olig #807 (69 bases)
TATTGGTTAC-GCGTGGGCCA-TGGTGGTGGT-TAT CGTTGGT-GCTACC--ATCG-GGATCAAACT-
ataaccaatg-cgcacccggt-accaccacca-ata--qcaacca-cgatgg tagc-cctagtttga-
↑ Olig #812 (65 bases)

**Table 104: Definition and alignment of oligonucleotides
(continued)**

↓ Olig #808 (38 bases)
 CCAGTCTTA-AGCCC--GCCTA--
 GTTCAAGAAG-TTTACTTCTGA-AGCGTCTTA-ATGATAGGGT-TA
 caagtcttc-aaatgaagct-tccgcagaat-tactatccca-at--ggtcagat-tcggg cggat-
 Olig #811 (69 bases) ↑

↓ filled in
ATGAGCGGC-TTTTTTTTA TCGA-3'
tactcgcccg-aaaaaaaaat-agct-5'
↑ Olig #810 (29 bases)

Table 104: Definition and alignment of oligonucleotides
(continued)

Overlap Sequences	Junction	T _m
AGGCTTACTATGAAG	802:817	42.
TGTCCTTCGCTCG	803:816	42.
CTATTCTACAATGC	804:815	40.
AACAACTTTAAATCG	805:814	38.
CCTTCAATTCTCTGC	806:813	44.
CGTTGGTGCTACC	807:812	42.
CCAGTCTAAGCCC	808:811	42.

All these ends, as well as the SalI end, was tested for self annealing, hair-pin loop formation, and cross hybridization. No unwanted hybridization is likely. Ideally, all fragments would be the same length, but placement of overlaps to avoid restriction sites (which are usually palindromic) and to avoid cross hybridization lead to fragments varying from 65 to 76 bases, plus two fragments of 29 to 38 bases.

Table 105: Individual sequences of Oligonucleotides 801-817.

Olig #801 (68 bases)	5'-ggcTTTAcAc TTTATgCTTc cggCTcgTAT AATgTgTgga ATTgTgAgcg CTCACAATTg AgCTcAgg-3'
Olig #802 (67 bases)	5'-AggcTTACTA TgAAGAAATc TcTggTTCTT AAggcTAGcg TTgCTgTcgc gAcccTggtA cCTATgT-3'
Olig #803 (70 bases)	5'-TgTccTTCgc TcgTccggAT TTcTgTcTcg AgccAccATA cACTggggccc TgcAAAgcgC gCATcATccg-3'
Olig #804 (67 bases)	5'-CTATTTCTAc AATgCTAAAg cAggccTgTg cCAGAccTTT gTATAcggtg gTTgcccTgc TAAgCGT-3'
Olig #805 (76 bases)	5'-AACAACTTTA AATcgccgA AgATTgCATg cGTAccTgcg gTggcgccgc TgAAGgTgAT gATccggcca Aggcgg-3'
Olig #806 (67 bases)	5'-cCTTCAATTc TcTgCAAgCT TcTgCTAccg AgTATATTgg TTAcgCGTgg gccATggTgg TggTTAT-3'
Olig #807 (69 bases)	5'-cgTTgTgCT AccATcgga TcAAAcTgTT cAAGAAgTTT ACTTcgAAgg cGTCTTAATg ATAgggTTA-3'
Olig #808 (38 bases)	5'-cCAGTCTAAG cccgcCTAAT gAgcgggCTT TTTTTTAA-3'

Table 105: Individual sequences of Oligonucleotides 801-817. (continued)

Olig #810 (29 bases)	
5'-TCGATAAAAA AAAAGCCCGC TCATTAGGC-3'	
Olig #811 (69 bases)	
5'-GGGCTTAGAC TGGTAACCCCT ATCATTAAAG CCGCTTCGAA GTAAACTTCT TGAACAGTTT GATCCCGAT-3'	
Olig #812 (65 bases)	
5'-GGTAGCAACA ACGATAACCA CCACCATGGC CCACGCGTAA CCAATATACT CGGTAGCAGA AGCTT-3'	
Olig #813 (76 bases)	
5'-GCAGAGAATT GAAGGCCGCC TTGGCCGGAT CATCACCTTC AGCGGCGCCA CCGCAGGTAC GCATGCAATC TTCGGC-3'	
Olig #814 (67 bases)	
5'-CGATTTAAAG TTGTTACGCT TAGCACGGCA AccACCGTAT ACAAAGGTCT ggcACAGGCC TgCTTTA-3'	
Olig #815 (72 bases)	
5'-GCATTGTAGA AATAGCGGAT gATgcgcgct TTgcagggcc cAGTgTATgg TggCTcgAGA CAGAAATCCG gA-3'	
Olig #816 (65 bases)	
5'-CGAGCGAAGG ACAACATAGg TaccAGgTc ggcACagCA cgcTAGcCTT AAGAAccAGA gATTT-3'	
Olig #817 (68 bases)	
5'-CTTCATAGTA AgcCTcCTga gCTCAATTgT gAgcgcTcAc AATTccAcAc ATTATAcgAg ccgGAAGC-3'	

Table 106: Signal Peptides

PhoA	M K q s t i a l a l l p l l f t p v t K A / R T...	(17)
MalE	M K I K T G A R i l l a l s a l t t m m f s a s a l a / K I...	(18)
OmpF	M M K R n i l a v i v p a l l v a g t a n a / a E...	(19)
Bla	M S I Q H F R v a l i p f f a a f c l p v f a / h p...	(>18)
LamB	M M I T L R K l p l a v a v a a g v m s a q a m a / v D...	(19)
Lpp	M K A T K l v l g a v i l g s t l l a g / c s...	(>17)
gpIII	M K K l l l f a i p l v v p p f y s h s / a E T V E...	(16)
gpIII-BPTI	M K K l l l f a i p l v v p p f y s g a / R P D...	(15)
gpVIII	M K K S L V L K a s v a v a t l v p m l s f a / a E G D D...	(16)
gpVIII-BPTI	M K K S L V L K a s v a v a t l v p m l s f a / R P D...	(15)
gpVIII'	M K K s l v l l a s v a v a t l v p m l s f a / a E G D D...	(21)

5	31 kd species ^a	14.5 kd species ^b
No DNA (control)	- ^c	-
pGEN-3Zf(-)	+	-
pGEM-MB16	+	-
pGEM-MB20	+	+
10 pGEM-MB26	+	+
pGEM-MB42	+	+
pGEM-MB46	ND	ND

15 a.) pre-beta-lactamase, encoded by the amp (bla)
gene.
 b.) pre-BPTII/VIII peptides encoded by the synthetic
gene and derived constructs.
 c.) - for absence of product; + for presence of
20 product; ND for Not Determined.

Table 108: Western analysis^a of in vivo
expressed
signal::BPTI::mature VIII protein species

5 A) expression in strain XL1-Blue				
	signal	14.5 kd species ^b	12 kd species ^c	
	pGEM-3Zf(-)	-	-d	-
	pGEM-MB16	VIII	-	-
	pGEM-MB20	VIII	++	-
10	pGEM-MB26	VIII	+++	+/-
	pGEM-MB42	phoA	++	+

B) expression in strain SEF'

	signal	14.5 kd species ^b	12 kd species ^c	
15	pGEM-MB42	phoA	+/-	+++

Notes:

a) Analysis using rabbit anti-BPTI polyclonal antibodies and horse-radish-peroxidase-conjugated goat anti-rabbit IgG antibody.

b) pro-BPTI/VIII peptides encoded by the synthetic gene and derived constructs.

c) processed BPTI/VIII peptide encoded by the synthetic gene.

25 d) not present -
weakly present +/-
present +
strong presence ++
very strong presence +++

55056850

Table 109: M13 gene III

1579	5'-GT	GAAAAAATTA	TTATTCGCAA	TTCCTTTAGT
1611	TGTTTCCTTTC	TATTCTCACT	CCGCTGAAAC	TGTTGAAAGT
1651	TGTTTAGCAA	AACCCCATAC	AGAAAATTCA	TTTACTAACG
5	1691	TCTGGAAAGA	CGACAAAAC	TTAGATCGTT
	1731	TGAGGGTTGT	CTGTGGAATG	CTACAGGCGT
	1771	ACTGGTGACG	AAACTCAGTG	TTACGGTACA
	1811	TTGGGCTTGC	TATCCCTGAA	AATGAGGGTG
	1851	GGGTGGCGGT	TCTGAGGGTG	GCGGTTCTGA
10	1891	ACTAAACCTC	CTGAGTACGG	TGATACACCT
	1931	ATACTTATAT	CAACCCTCTC	GACGGCACTT
	1971	TACTGAGCAA	AACCCGCTA	ATCCTAATCC
	2011	GAGTCTCAGC	CTCTTAATAC	TTTCATGTTT
	2051	GGTTCCGAAA	TAGGCAGGGG	GCATTAAC
15	2091	CACTGTTACT	CAAGGCACTG	ACCCCGTTAA
	2131	CAGTACACTC	CTGTATCATC	AAAAGCCATG
	2171	ACTGGAACGG	TAAATTCAGA	GACTGCGCTT
	2211	CTTTAATGAG	GATCCATTCG	TTTGTGAATA
	2251	TCGTCTGACC	TGCCTCAACC	TCCTGTCAAT
20	2291	GCTCTGGTGG	TGGTTCCTGGT	GGCGGCTCTG
	2331	CTCTGAGGGT	GGCGGTTCTG	AGGGTGGCGG
	2371	GGCGGTTCCG	GTGGTGGCTC	TGGTTCGGGT
	2411	ATGAAAAGAT	GGCAAACGCT	AATAAGGGGG
	2451	AAATGCCGAT	GAAAACGCGC	TACAGTCTGA
25	2491	AAACTTGATT	CTGTGCTAC	TGATTACGGT
	2531	ATGGTTTCAT	TGGTGACGTT	TCCGGCCTTG
	2571	TGGTGCTACT	GGTGATTTTG	CTGGCTCTAA
	2611	GCTCAAGTCG	GTGACGGTGA	TAATTCACCT
	2651	ATTTCCGTCA	ATATTTACCT	TCCCTCCCTC
30	2691	ATGTCGCCCT	TTTGTCTTTA	GCGCTGGTAA
	2731	TTTTCTATTG	ATTGTGACAA	AATAAACTTA
	2771	TCTTTGCGTT	TCTTTTATAT	GTTGCCACCT
	2811	ATTTTCTACG	TTTGCTAACA	TACTGCGTAA
	2851	TAATCATGCC	AGTTCTTTTG	GGTATTCGGT

Table 110: Introduction of NarI into gene IIIA) Wild-type III, portion encoding the signal peptide

5		M	K	K	L	L	F	A	I	P	L	
		1	2	3	4	5	6	7	8	9	10	
1579		5'-GTG	AAA	AAA	TTA	TTA	TTC	GCA	ATT	CCT	TTA	
10								/	Cleavage site			
		V	V	P	F	Y	S	H	S	A	E	T V
		11	12	13	14	15	16	17	18	19	20	21 22
1609		GTT	GTT	CCT	TTC	TAT	TCT	CAC	TCC	GCT	GAA	ACT GTT-3'
15												

B) III, portion encoding the signal peptide with NarI site

20		m	k	k	l	l	f	a	I	p	l	
		1	2	3	4	5	6	7	8	9	10	
1579		5'-gtg	aaa	aaa	tta	tta	ttc	gca	att	cct	tta	
25								/	cleavage site			
		v	v	p	f	y	s	G	A	a	e	t v
		11	12	13	14	15	16	17	18	19	20	21 22
30	1609	gtt	gtt	cct	ttc	tat	tct	GGc	Gcc	gct	gaa	act gtt-3'

T06230-55095500

Table 111: IIIsp::bpti::mautreIII fusion gene.

```

      m   k   k   l   l   f   a   I   p   l
      1   2   3   4   5   6   7   8   9  10
5' -gtg aaa aaa tta tta ttc gca att cct tta
    |<---- gene III signal peptide -----
                                     / cleavage site
                                     v
      v   v   p   f   y   s   G   A
11  12  13  14  15  16  17  18
gtt gtt cct ttc tat tct GGc Gcc
----->|

      R   P   D   F   C   L   E
      19  20  21  22  23  24  25
CGT CCG GAT TTC TGT CTC GAG -
      |AccIII|          |Ava I|
M13/BPTI Jnct          |Xho I|

      P   P   Y   T   G   P   C   K   A   R
      26  27  28  29  30  31  32  33  34  35
CCA CCA TAC ACT GGG CCC TGC AAA GCG CGC _
      |PflM I|          |BssH II|
      |Apa I|
      |Dra II|
      |Pss I|

      I   I   R   Y   F   Y   N   A   K   A
      36  37  38  39  40  41  42  43  44  45
ATC ATC CGC TAT TTC TAC AAT GCT AAA GC -

      G   L   C   Q   T   F   V   Y   G   G
      46  47  48  49  50  51  52  53  54  55
A GGC CTG TGC CAG ACC TTT GTA TAC GGT GGT -
|Stu I|          |Acc I|
                  |Xca I|

      C   R   A   K   R   N   N   F   K
      56  57  58  59  60  61  62  63  64
TGC CGT GCT AAG CGT AAC AAC TTT AAA -
      |Esp I|

```

45

S	A	E	D	C	M	R	T	C	G
65	66	67	68	69	70	71	72	73	74
TCG	GCC	GAA	GAT	TGC	ATG	CGT	ACC	TGC	GGT
<u>XmaIII</u>				<u>Sph I</u>					

10

G	A
75	76
GGC	GCC
Bbe I	

15

Nar I

BPTI/M13 boundary

	G	A	a	e	t	v	e	s
	77	78	79	80	81	82	83	84
20	GGc	Gcc	gct	gaa	act	gtt	GAA	AGT

	1651	TGTTTAGCAA	AACCCCATAC	AGAAAATTCA	TTTACTAACG
	1691	TCTGGAAAGA	CGACAAAAC	TTAGATCGTT	ACGCTAACTA
	1731	TGAGGGTTGT	CTGTGGAATG	CTACAGGCGT	TGTAGTTTGT
25	1771	ACTGGTGACG	AAACTCAGTG	TTACGGTACA	TGGGTTCCCTA
	1811	TTGGGCTTGC	TATCCCTGAA	AATGAGGGTG	GTGGCTCTGA
	1851	GGGTGGCGGT	TCTGAGGGTG	GCGGTTCTGA	GGGTGGCGGT
	1891	ACTAAACCTC	CTGAGTACGG	TGATACACCT	ATTCGGGGCT
	1931	ATACTTATAT	CAACCCTCTC	GACGGCACTT	ATCCGCCTGG
30	1971	TACTGAGCAA	AACCCCGCTA	ATCCTAATCC	TTCTCTTGAG
	2011	GAGTCTCAGC	CTCTTAATAC	TTTCATGTTT	CAGAATAATA
	2051	GGTTCCGAAA	TAGGCAGGGG	GCATTAAC	TTTATACGGG
	2091	CACTGTTACT	CAAGGCACTG	ACCCCGTTAA	AACTTATTAC
	2131	CAGTACACTC	CTGTATCATC	AAAAGCCATG	TATGACGCTT
35	2171	ACTGGAACGG	TAAATTGAGA	GACTGCGCTT	TCCATTCTGG
	2211	CTTTAATGAG	GATCCATTCG	TTTGTGAATA	TCAAGGCCAA
	2251	TCGTCTGACC	TGCCTCAACC	TCCTGTCAAT	GCTGGCGGCG
	2291	GCTCTGGTGG	TGGTTCCTGG	GGCGGCTCTG	AGGGTGGTGG
	2331	CTCTGAGGGT	GGCGGTTCTG	AGGGTGGCGG	CTCTGAGGGA
40	2371	GGCGGTTCCG	GTGGTGGCTC	TGGTTCGGGT	GATTTTGATT
	2411	ATGAAAAGAT	GGCAAACGCT	AATAAGGGGG	CTATGACCGA
	2451	AAATGCCGAT	GAAAACGCGC	TACAGTCTGA	CGCTAAAGGC

Table 111, continued

	2491	AAACTTGATT	CTGTCGCTAC	TGATTACGGT	GCTGCTATCG
	2531	ATGGTTTCAT	TGGTGACGTT	TCCGGCCTTG	CTAATGGTAA
	2571	TGGTGCTACT	GGTGATTTTG	CTGGCTCTAA	TTCCCAAATG
5	2611	GCTCAAGTCG	GTGACGGTGA	TAATTCACCT	TTAATGAATA
	2651	ATTTCCGTCA	ATATTTACCT	TCCCTCCCTC	AATCGGTTGA
	2691	ATGTCGCCCT	TTTGTCTTTA	GCGCTGGTAA	ACCATATGAA
	2731	TTTTCTATTG	ATTGTGACAA	AATAAACTTA	TTCCGTGGTG
	2771	TCTTTGCGTT	TCTTTTATAT	GTTGCCACCT	TTATGTATGT
10	2811	ATTTTCTACG	TTTGCTAACA	TACTGCGTAA	TAAGGAGTCT
	2851	<u>TAATCATGCC</u>	AGTTCTTTTG	GGTATTCCGT	--

F06290"5055850

5

BamHI

ctg TTGACA attaatcatcgGCTCG tataat GTGTGG-
 -35 tac -10

aATTGTGAGCGcTcACAATT-
lacO-symm operator

GAGCTC T ggagga AATAAA-
SacI Shine-Dalgarno seq.

fM	K	K	S	L	V	L	K	A	S	
1	2	3	4	5	6	7	8	9	10	
ATG	AAG	AAA	TCT	CTG	GTT	CTT	AAG	GCT	AGC	-
						Afl II		Nhe I		

V	A	V	A	T	L	V	P	M	L
11	12	13	14	15	16	17	18	19	20
GTT	GCT	GTC	GCG	ACC	CTG	GTA	CCT	ATG	TTG
Nru I					Kpn I				

S	F	A	R	P	D	F	C	L	E
21	22	23	24	25	26	27	28	29	30
TCC	TTC	GCT	CGT	CCG	GAT	TTC	TGT	CTC	GAG
AccIII						Ava I			

M13/BPTI Jct	Xho I
--------------	-------

P	P	Y	T	G	P	C	K	A	R
31	32	33	34	35	36	37	38	39	40
CCA	CCA	TAC	ACT	GGG	CCC	TGC	AAA	GCG	CGC
PflM I					BssH II				
Apa I									
Dra II									
Pss I									

5

10

152025

30

35404550

BstX I
Nco I

Table 112 : Annotated Sequence of
 Ptac::RBS(GGAGGAAATAAA)::
VIII-signal::mature-bpti::mature-VIII-coat-protein gene
 (continued)

5

T	I	G	I
117	118	119	120
ACC	ATC	GGG	ATC

 -

10

K	L	F	K	K	F	T	S	K	A
121	122	123	124	125	126	127	128	129	130
AAA	CTG	TTC	AAG	AAG	TTT	ACT	TCG	AAG	GCG

 -
Asu II

15

S	.	.	.
131	132	133	134
TCT	TAA	TGA	TAG

GGTTACC-
BstE II

20

AGTCTA AGCCCGC CTAATGA GCGGGCT TTTTTTTT-
terminator

25

aTCGA GACctgca GGTGACC ggcattgc-3'
SalI

T06230" 55035350

Table 113 : Annotated Sequence of
 pGEM-MB42 comprising Ptac::RBS(GGAGGAAATAAA)::
phoA-signal::mature-bpti::mature-VIII-coat-protein

5 5'-GGATCC actcccccattcccc
 |
 BamHI

10 ctg TTGACA attaatcatcgGCTCG tataat GTGTGG-
 -35 tac -10

15 aATTGTGAGCGcTcACAATT-
 lacO-symm operator

20 GAGCTCCATGGGAGAAAATAAA | M | K | Q | S | T |
 |SacI| | 1 | 2 | 3 | 4 | 5 |
 ATG AAA CAA AGC ACG | -
 <----- phoA signal peptide

25 | I | A | L | L | P | L | L | F | T | P | V | T |
 6 7 8 9 10 11 12 13 14 15 16 17 |
 ATC GCA CTC TTA CCG TTA CTG TTT ACC CCT GTG ACA | -
 ----- phoA signal continues -----

30 (There are no residues 20-23.)

35 | K | A | R | P | D | F | C | L | E |
 18 19 24 25 26 27 28 29 30 |
 AAA GCC CGT CCG GAT TTC TGT CTC GAG -
 phoA signal-> | AccIII | Ava I |
 phoA/BPTI Jnct | Xho I |
 |<----- BPTI insert ----->

40 | P | P | Y | T | G | P | C | K | A | R |
 31 32 33 34 35 36 37 38 39 40 |
 CCA CCA TAC ACT GGG CCC TGC AAA GCG CGC | -
 | PflM I | | BssH II |

45 | Apa I |
 | Dra II |
 | Pss I |

0936095 062901
 106290 5595860

Table 113 : Annotated Sequence of
 Ptac::RBS (GGAGGAAATAAA)::
phoA-signal::mature-bpti::mature-VIII-coat-protein gene
 (continued)

5

I	I	R	Y	F	Y	N	A	K	A	
41	42	43	44	45	46	47	48	49	50	
ATC	ATC	CGC	TAT	TTC	TAC	AAT	GCT	AAA	GC	-

10

G	L	C	Q	T	F	V	Y	G	G	
51	52	53	54	55	56	57	58	59	60	
GGC	CTG	TGC	CAG	ACC	TTT	GTA	TAC	GGT	GGT	-
<u>Stu I</u>						<u>Acc I</u>				
						<u>Xca I</u>				

15

C	R	A	K	R	N	N	F	K		
61	62	63	64	65	66	67	68	69		
TGC	CGT	GCT	AAG	CGT	AAC	AAC	TTT	AAA		-
<u>Esp I</u>										

20

S	A	E	D	C	M	R	T	C	G	
70	71	72	73	74	75	76	77	78	79	
TCG	GCC	GAA	GAT	TGC	ATG	CGT	ACC	TGC	GGT	-
<u>XmaIII</u>				<u>Sph I</u>						

----- BPTI insert -----

25

BPTI/M13 boundary

G	A	A	E	G	D	D	P	A	K	A	A	
80	81	82	83	84	85	86	87	88	89	90	91	
GGC	GCC	GCT	GAA	GGT	GAT	GAT	CCG	GCC	AAG	GCG	GCC	-
<u>Bbe I</u>								<u>Sfi I</u>				
<u>Nar I</u>												

-- BPTI--> <----- mature gene VIII coat protein -----

30

F	N	S	L	Q	A	S	A	T		
92	93	94	95	96	97	98	99	100		
TTC	AAT	TCT	CTG	CAA	GCT	TCT	GCT	ACC		-
<u>Hind 3</u>										

35

E	Y	I	G	Y	A	W				
101	102	103	104	105	106	107				
GAG	TAT	ATT	GGT	TAC	GCG	TGG				-

40

45

Table 113 : Annotated Sequence of
 Ptac::RBS(GGAGGAAATAAA)::
phoA-signal::mature-bpti::mature-VIII-coat-protein gene
 (continued)

5

A	M	V	V	V	I	V	G	A
108	109	110	111	112	113	114	115	116
GCC	ATG	GTG	GTG	GTT	ATC	GTT	GGT	GCT

10 BstX I
 Nco I

15

T	I	G	I
117	118	119	120
ACC	ATC	GGG	ATC

-

20

K	L	F	K	K	F	T	S	K	A
121	122	123	124	125	126	127	128	129	130
AAA	CTG	TTC	AAG	AAG	TTT	ACT	TCG	AAG	GCG

-
 Asu II

25

S	.	.	.
131	132	133	134
TCT	TAA	TGA	TAG

GGTTACC-
BstE II

30

AGTCTA AGCCCGC CTAATGA GCGGGCT TTTTTTTT-
terminator

35

aTCGA GACctgca GGTCGAC-3'
 SalI

T002200-560365B50

Table 114: Neutralization of Phage Titer Using
Agarose-immobilized Anhydro-Trypsin

5	Phage Type	Addition	Percent Residual Titer As a Function of Time (hours)		
			1	2	4
10	MK-BPTI	5 μ l IS	99	104	105
		2 μ l IAT	82	71	51
		5 μ l IAT	57	40	27
		10 μ l IAT	40	30	24
15	MK	5 μ l IS	106	96	98
		2 μ l IAT	97	103	95
		5 μ l IAT	110	111	96
		10 μ l IAT	99	93	106

Legend:

IS = Immobilized streptavidin

IAT = Immobilized anhydro-trypsin

1062200 5605660

Table 115: Affinity Selection of MK-BPTI Phage
on Immobilized Anhydro-Trypsin

	Phage Type	Addition	Percent of Total Phage Recovered in Elution Buffer
5	MK-BPTI	5 μ l IS	<<1 ^a
		2 μ l IAT	5
		5 μ l IAT	20
		10 μ l IAT	50
10	MK	5 μ l IS	<<1 ^a
		2 μ l IAT	<<1
		5 μ l IAT	<<1
		10 μ l IAT	<<1

15

Legend:

IS = Immobilized streptavidin

IAT = Immobilized anhydro-trypsin

^a not detectable.

20

T05230" 5696860

Table 116: translation of Signal-III::bpti::mature-III

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15
 fM K K L L L F A I P L V V P F Y
 GTG AAA AAA TTA TTA TTC GCA ATT CCT TTA GTT CCT TTC TAT
 |<----- gene III signal peptide ----->|

5

16 17 18 19 20 21 22 23 24 25 26 27 28 29 30
 S G A R P P D F C L E P P Y T G
 TCT GGC GCC cgt ccg gat ttc tgt ctc gag cca cca tac act ggg
 ----->|<----- BPTI insertion ----->|

10

31 32 33 34 35 36 37 38 39 40 41 42 43 44 45
 P C K A R I I R Y F Y N A K A
ccc tgc aaa gcg cgc atc atc cgc tat ttc tac aat gct aaa gca

15

46 47 48 49 50 51 52 53 54 55 56 57 58 59 60
 G L C Q T F V Y G C R A K R
ggc ctg tgc cag acc ttt gta tac ggt ggt tgc cgt gct aag cgt

20

61 62 63 64 65 66 67 68 69 70 71 72 73 74 75
 N N F K S A E D C M R T C G G
aac aac ttt aaa tcg gcc gaa gat tgc atg cgt acc tgc ggt ggc

25

76 77 78 79 80 81 82 83 84 85 86 87 88 89 90
 A G A A E T V E S C L A K P H
gcc gcc gcc GCT GAA ACT GTT GAA AGT TGT TTA GCA AAA CCC CAT
 |<----- mature gene III protein ----->|

30

91 92 93 94 95 96 97 98 99 100 101 102 103 104 105
 T E N S F T N V W K D K T L
 ACA GAA AAT TCA TTT ACT AAC GTC TGG AAA GAC GAC AAA ACT TTA

Table 116: translation of Signal-III::bpti::mature-III (continued)

5	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120
	D	R	Y	A	N	Y	E	G	C	L	W	N	A	T	G
	GAT	CGT	TAC	GCT	AAC	TAT	GAG	GGT	TGT	CTG	TGG	AAT	GCT	ACA	GGC
	121	122	123	124	125	126	127	128	129	130	131	132	133	134	135
	V	V	V	C	T	G	D	E	T	Q	C	Y	G	T	W
	GTT	GTA	GTT	TGT	ACT	GGT	GAC	GAA	ACT	CAG	TGT	TAC	GGT	ACA	TGG
10	136	137	138	139	140	141	142	143	144	145	146	147	148	149	150
	V	P	I	G	L	A	I	P	E	N	E	G	G	G	S
	GTT	CCT	ATT	GGG	CTT	GCT	ATC	CCT	GAA	AAT	GAG	GGT	GGT	GGC	TCT
15	151	152	153	154	155	156	157	158	159	160	161	162	163	164	165
	E	G	G	G	S	E	G	G	G	S	E	G	G	G	T
	GAG	GGT	GGC	GGT	TCT	GAG	GGT	GGC	GGT	TCT	GAG	GGT	GGC	GGT	ACT
20	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180
	K	P	P	E	Y	G	D	T	P	I	P	G	Y	T	Y
	AAA	CCT	CCT	GAG	TAC	GGT	GAT	ACA	CCT	ATT	CCG	GGC	TAT	ACT	TAT
25	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195
	I	N	P	L	D	G	T	Y	P	P	G	T	E	Q	N
	ATC	AAC	CCT	CTC	GAC	GGC	ACT	TAT	CCG	CCT	GGT	ACT	GAG	CAA	AAC
	196	197	198	199	200	201	202	203	204	205	206	207	208	209	210
	P	A	N	P	N	P	S	L	E	E	S	Q	P	L	N
	CCC	GCT	AAT	CCT	AAT	CCT	TCT	CTT	GAG	GAG	TCT	CAG	CCT	CTT	AAT
30	211	212	213	214	215	216	217	218	219	220	221	222	223	224	225
	T	F	M	F	Q	N	N	R	F	R	N	R	Q	G	A
	ACT	TTC	ATG	TTT	CAG	AAT	AAT	AGG	TTC	CGA	AAT	AGG	CAG	GGG	GCA

Table 116: translation of Signal-III::bpti::mature-III (continued)

5	226	L	TTA	ACT	GTT	TAT	ACG	GGC	GGC	ACT	GTT	ACT	CAA	GGC	ACT	GAC	CCC	GTT
	227	T																
	228	V																
	229	Y																
	230	T																
	231	G																
	232	T																
	233	V																
	234	T																
	235	Q																
	236	G																
	237	T																
	238	D																
	239	P																
	240	V																
10	241	K	AAA	ACT	TAT	TAC	CAG	TAC	TAC	ACT	CCT	GTA	TCA	TCA	AAA	GCC	ATG	TAT
	242	T																
	243	Y																
	244	Y																
	245	Q																
	246	Y																
	247	T																
	248	P																
	249	V																
	250	S																
	251	S																
	252	K																
	253	A																
	254	M																
	255	Y																
15	256	D	GAC	GCT	TAC	TGG	AAC	GGT	AAA	TTC	AGA	GAC	TGC	TGC	GCT	TTC	CAT	TCT
	257	A																
	258	Y																
	259	W																
	260	N																
	261	G																
	262	K																
	263	F																
	264	R																
	265	D																
	266	C																
	267	A																
	268	F																
	269	H																
	270	S																
20	271	G	GGC	TTT	AAT	GAG	GAT	CCA	TTC	GTT	TGT	GAA	TAT	CAA	GGC	CAA	TCG	
	272	F																
	273	N																
	274	E																
	275	D																
	276	P																
	277	F																
	278	V																
	279	C																
	280	E																
	281	Y																
	282	Q																
	283	G																
	284	Q																
	285	S																
25	286	S	TCT	GAC	CTG	CCT	CAA	CCT	CCT	CCT	AAT	GCT	GGC	GGC	TCT	TCT	GGT	
	287	L																
	288	P																
	289	Q																
	290	P																
	291	P																
	292	P																
	293	V																
	294	N																
	295	A																
	296	G																
	297	G																
	298	G																
	299	S																
	300	G																
30	301	G	GGT	GGT	TCT	TCT	GGC	GGC	GGC	TCT	GAG	GGT	GGT	GGC	TCT	GAG	GGT	GGC
	302	S																
	303	G																
	304	G																
	305	G																
	306	G																
	307	S																
	308	E																
	309	G																
	310	G																
	311	G																
	312	S																
	313	E																
	314	G																
	315	G																
25	316	G	GGT	TCT	GAG	GGT	GGC	GGC	GGC	TCT	GAG	GGT	GGC	GGC	TCC	GGT	GGT	GGC
	317	S																
	318	E																
	319	G																
	320	G																
	321	G																
	322	S																
	323	E																
	324	G																
	325	G																
	326	G																
	327	S																
	328	G																
	329	G																
	330	G																
30	331	S	TCT	GGT	TCC	GGT	GAT	TTT	GAT	TAT	GAA	AAG	ATG	GCA	AAC	AAC	GCT	AAT
	332	G																
	333	S																
	334	G																
	335	D																
	336	F																
	337	D																
	338	Y																
	339	E																
	340	K																
	341	M																
	342	A																
	343	N																
	344	A																
	345	N																

Table 116: translation of Signal-III::bpti::mature-III (continued)

5	346	347	348	349	350	351	352	353	354	355	356	357	358	359	360
	K	G	A	M	T	E	N	A	D	E	N	A	L	Q	S
	AAG	GGG	GCT	ATG	ACC	GAA	AAT	GCC	GAT	GAA	AAC	GCG	CTA	CAG	TCT
10	361	362	363	364	365	366	367	368	369	370	371	372	373	374	375
	D	A	K	G	K	L	D	S	V	A	T	D	Y	G	A
	GAC	GCT	AAA	GGC	AAA	CTT	GAT	TCT	GTC	GCT	ACT	GAT	TAC	GGT	GCT
	376	377	378	379	380	381	382	383	384	385	386	387	388	389	390
	A	I	D	G	F	I	G	D	V	S	G	L	A	N	G
	GCT	ATC	GAT	GGT	TTC	ATT	GGT	GAC	GTT	TCC	GGC	CTT	GCT	AAT	GGT
15	391	392	393	394	395	396	397	398	399	400	401	402	403	404	405
	N	G	A	T	G	D	F	A	G	S	N	S	Q	M	A
	AAT	GGT	GCT	ACT	GGT	GAT	TTT	GCT	GGC	TCT	AAT	TCC	CAA	ATG	GCT
20	406	407	408	409	410	411	412	413	414	415	416	417	418	419	420
	Q	V	G	D	G	D	N	S	P	L	M	N	N	F	R
	CAA	GTC	GGT	GAC	GGT	GAT	AAT	TCA	CCT	TTA	ATG	AAT	AAT	TTC	CGT
25	421	422	423	424	425	426	427	428	429	430	431	432	433	434	435
	Q	Y	L	P	S	L	P	Q	S	V	E	C	R	P	F
	CAA	TAT	TTA	CCT	TCC	CTC	CCT	CAA	TCG	GTT	GAA	TGT	CGC	CCT	TTT
30	436	437	438	439	440	441	442	443	444	445	446	447	448	449	450
	V	F	S	A	G	K	P	Y	E	F	S	I	D	C	D
	GTC	TTT	AGC	GCT	GGT	AAA	CCA	TAT	GAA	TTT	TCT	ATT	GAT	TGT	GAC

Table 116: translation of Signal-III::bpti::mature-III (continued)

5	451	452	453	454	455	456	457	458	459	460	461	462	463	464	465
	K	I	N	L	F	R	G	V	F	A	F	L	L	Y	V
	AAA	ATA	AAC	TAA	TTC	CGT	GGT	GTC	TTT	GCG	TTT	CTT	TTA	TAT	GTT
	<----- uncharged anchor region ----->														
10	466	467	468	469	470	471	472	473	474	475	476	477	478	479	480
	A	T	F	M	Y	V	F	S	T	F	A	N	I	L	R
	GCC	ACC	TTT	ATG	TAT	GTA	TTT	TCT	ACG	TTT	GCT	AAC	ATA	CTG	CGT
	----- uncharged anchor region continues ----->														

481 482 483 484 485
 N K E S
 AAT AAG GAG TCT TAA

20	Molecular weight of peptide	=	58884
	Charge on peptide	=	-20
	[A+G+P]	=	143
	[C+F+H+I+L+M+V+W+Y]	=	140
	[D+E+K+R+N+Q+S+T+.]	=	202

Table 116: translation of Signal-III::bpti::mature-III (continued)

Second Base		a		g	
	t	c	a	t	c
5	15	21	15	8	t
	12	5	10	6	c
	10	4	0	0	a
	0	3	0	4	g
10	6	20	2	8	t
	3	4	0	3	c
	1	4	9	1	a
	4	3	7	0	g
15	5	19	21	1	t
	5	4	11	1	c
	2	4	16	1	a
	8	2	4	2	g
20	13	22	14	41	t
	6	7	12	29	c
	4	5	12	1	a
	1	3	16	4	g
25					
	AA	#	AA	#	AA
	A	37	C	14	D
	F	27	G	75	H
	K	20	L	24	M
	P	31	Q	16	R
	T	29	V	23	W
	.	1			
30					
	AA	#	AA	#	AA
	E	26	D	26	E
	I	2	H	2	I
	N	9	M	9	N
	S	15	R	15	S
	Y	4	W	4	Y
25					
	AA	#	AA	#	AA
	E	28	D	28	E
	I	12	H	12	I
	N	32	M	32	N
	S	35	R	35	S
	Y	25	W	25	Y

Table 130: Sampling of a Library encoded by (NNK)⁶

A. Numbers of hexapeptides in each class

5 total = 64,000,000 stop-free sequences.

α can be one of [WMFYCIKDENVH]Q]

Φ can be one of [PTAVG]

Ω can be one of [SLR]

10	$\alpha\alpha\alpha\alpha\alpha\alpha$	=	2985984.	$\Phi\alpha\alpha\alpha\alpha\alpha$	=	7464960.
	$\Omega\alpha\alpha\alpha\alpha\alpha$	=	4478976.	$\Phi\Phi\alpha\alpha\alpha\alpha$	=	7776000.
	$\Phi\Omega\alpha\alpha\alpha\alpha$	=	9331200.	$\Omega\Omega\alpha\alpha\alpha\alpha$	=	2799360.
	$\Phi\Phi\Phi\alpha\alpha\alpha$	=	4320000.	$\Phi\Phi\Omega\alpha\alpha\alpha$	=	7776000.
15	$\Phi\Omega\Omega\alpha\alpha\alpha$	=	4665600.	$\Omega\Omega\Omega\alpha\alpha\alpha$	=	933120.
	$\Phi\Phi\Phi\Phi\alpha\alpha$	=	1350000.	$\Phi\Phi\Phi\Omega\alpha\alpha$	=	3240000.
	$\Phi\Phi\Omega\Omega\alpha\alpha$	=	2916000.	$\Phi\Omega\Omega\Omega\alpha\alpha$	=	1166400.
	$\Omega\Omega\Omega\Omega\alpha\alpha$	=	174960.	$\Phi\Phi\Phi\Phi\Phi\alpha$	=	225000.
	$\Phi\Phi\Phi\Phi\Omega\alpha$	=	675000.	$\Phi\Phi\Phi\Omega\Omega\alpha$	=	810000.
20	$\Phi\Phi\Omega\Omega\Omega\alpha$	=	486000.	$\Phi\Omega\Omega\Omega\Omega\alpha$	=	145800.
	$\Omega\Omega\Omega\Omega\Omega\alpha$	=	17496.	$\Phi\Phi\Phi\Phi\Phi\Phi$	=	15625.
	$\Phi\Phi\Phi\Phi\Phi\Omega$	=	56250.	$\Phi\Phi\Phi\Phi\Omega\Omega$	=	84375.
	$\Phi\Phi\Phi\Omega\Omega\Omega$	=	67500.	$\Phi\Phi\Omega\Omega\Omega\Omega$	=	30375.
	$\Phi\Omega\Omega\Omega\Omega\Omega$	=	7290.	$\Omega\Omega\Omega\Omega\Omega\Omega$	=	729.

25 $\Phi\Phi\Omega\Omega\alpha\alpha$, for example, stands for the set of peptides having two amino acids from the α class, two from Φ , and two from Ω arranged in any order. There are, for example, $729 = 3^6$ sequences composed entirely of S, L, and R.

30

0956095.0956095

Table 130: Sampling of a Library encoded by (NNK)⁶
(continued)

5 B. Probability that any given stop-free DNA sequence
will encode a hexapeptide from a stated class.

		P	% of class
	αααααα...	3.364E-03	(1.13E-07)
	ϕααααα...	1.682E-02	(2.25E-07)
10	ηααααα...	1.514E-02	(3.38E-07)
	ϕϕαααα...	3.505E-02	(4.51E-07)
	ϕηαααα...	6.308E-02	(6.76E-07)
	ηηαααα...	2.839E-02	(1.01E-06)
	ϕϕϕααα...	3.894E-02	(9.01E-07)
15	ϕϕηααα...	1.051E-01	(1.35E-06)
	ϕηηααα...	9.463E-02	(2.03E-06)
	ηηηααα...	2.839E-02	(3.04E-06)
	ϕϕϕϕαα...	2.434E-02	(1.80E-06)
	ϕϕϕηαα...	8.762E-02	(2.70E-06)
20	ϕϕηηαα...	1.183E-01	(4.06E-06)
	ϕηηηαα...	7.097E-02	(6.08E-06)
	ηηηηαα...	1.597E-02	(9.13E-06)
	ϕϕϕϕϕα...	8.113E-03	(3.61E-06)
	ϕϕϕϕηα...	3.651E-02	(5.41E-06)
25	ϕϕϕηηα...	6.571E-02	(8.11E-06)
	ϕϕηηηα...	5.914E-02	(1.22E-05)
	ϕηηηηα...	2.661E-02	(1.83E-05)
	ηηηηηα...	4.790E-03	(2.74E-05)
	ϕϕϕϕϕϕ...	1.127E-03	(7.21E-06)
30	ϕϕϕϕϕη...	6.084E-03	(1.08E-05)
	ϕϕϕϕηη...	1.369E-02	(1.62E-05)
	ϕϕϕηηη...	1.643E-02	(2.43E-05)
	ϕϕηηηη...	1.109E-02	(3.65E-05)
	ϕηηηηη...	3.992E-03	(5.48E-05)
35	ηηηηηη...	5.988E-04	(8.21E-05)

FOIb290-56096850

Table 130: Sampling of a Library encoded by (NNK)⁶
(continued)

C. Number of different stop-free amino-acid sequences in
each class expected for various library sizes

Library size = 1.0000E+06

total = 9.7446E+05 % sampled = 1.52

Class	Number	%	Class	Number	%
ααααα...	3362.6(.1)	ϕαααα...	16803.4(.2)
ηαααα...	15114.6(.3)	ϕϕααα...	34967.8(.4)
ϕηααα...	62871.1(.7)	ηηααα...	28244.3(1.0)
ϕϕααα...	38765.7(.9)	ϕϕηαα...	104432.2(1.3)
ϕηηαα...	93672.7(2.0)	ηηηαα...	27960.3(3.0)
ϕϕϕαα...	24119.9(1.8)	ϕϕϕηα...	86442.5(2.7)
ϕϕηηα...	115915.5(4.0)	ϕηηηα...	68853.5(5.9)
ηηηηα...	15261.1(8.7)	ϕϕϕϕα...	7968.1(3.5)
ϕϕϕηα...	35537.2(5.3)	ϕϕϕηη...	63117.5(7.8)
ϕϕηηη...	55684.4(11.5)	ϕηηηη...	24325.9(16.7)
ηηηηη...	4190.6(24.0)	ϕϕϕϕϕ...	1087.1(7.0)
ϕϕϕϕα...	5767.0(10.3)	ϕϕϕϕη...	12637.2(15.0)
ϕϕϕηη...	14581.7(21.6)	ϕϕηηη...	9290.2(30.6)
ϕηηηη...	3073.9(42.2)	ηηηηη...	408.4(56.0)

Library size = 3.0000E+06

total = 2.7885E+06 % sampled = 4.36

ααααα...	10076.4(.3)	ϕαααα...	50296.9(.7)
ηαααα...	45190.9(1.0)	ϕϕααα...	104432.2(1.3)
ϕηααα...	187345.5(2.0)	ηηααα...	83880.9(3.0)
ϕϕααα...	115256.6(2.7)	ϕϕηαα...	309107.9(4.0)
ϕηηαα...	275413.9(5.9)	ηηηαα...	81392.5(8.7)
ϕϕϕαα...	71074.5(5.3)	ϕϕϕηα...	252470.2(7.8)
ϕϕηηα...	334106.2(11.5)	ϕηηηα...	194606.9(16.7)
ηηηηα...	41905.9(24.0)	ϕϕϕϕα...	23067.8(10.3)
ϕϕϕηα...	101097.3(15.0)	ϕϕϕηη...	174981.0(21.6)
ϕϕηηη...	148643.7(30.6)	ϕηηηη...	61478.9(42.2)
ηηηηη...	9801.0(56.0)	ϕϕϕϕϕ...	3039.6(19.5)
ϕϕϕϕα...	15587.7(27.7)	ϕϕϕϕη...	32516.8(38.5)
ϕϕϕηη...	34975.6(51.8)	ϕϕηηη...	20215.5(66.6)
ϕηηηη...	5879.9(80.7)	ηηηηη...	667.0(91.5)

Table 130: Sampling of a Library encoded by (NNK)⁶
(continued)

Library size = 1.0000E+07

5

total = 8.1204E+06 % sampled = 12.69

	ααααα...	33455.9(1.1)	φαααα...	166342.4(2.2)
	Παααα...	148871.1(3.3)	φφααα...	342685.7(4.4)
10	ΦΠααα...	609987.6(6.5)	ΠΠααα...	269958.3(9.6)
	φφφαα...	372371.8(8.6)	φφΠαα...	983416.4(12.6)
	φΠΠαα...	856471.6(18.4)	ΠΠΠαα...	244761.5(26.2)
	φφφφα...	222702.0(16.5)	φφφΠα...	767692.5(23.7)
	φφΠΠα...	972324.6(33.3)	φΠΠΠα...	531651.3(45.6)
15	ΠΠΠΠα...	104722.3(59.9)	φφφφφ...	68111.0(30.3)
	φφφφΠ...	281976.3(41.8)	φφφΠΠ...	450120.2(55.6)
	φφΠΠΠ...	342072.1(70.4)	φΠΠΠΠ...	122302.6(83.9)
	ΠΠΠΠΠ...	16364.0(93.5)	φφφφφ...	8028.0(51.4)
	φφφφφ...	37179.9(66.1)	φφφφΠ...	67719.5(80.3)
20	φφφΠΠ...	61580.0(91.2)	φφΠΠΠ...	29586.1(97.4)
	φΠΠΠΠ...	7259.5(99.6)	ΠΠΠΠΠ...	728.8(100.0)

Library size = 3.0000E+07

25

total = 1.8633E+07 % sampled = 29.11

	ααααα...	99247.4(3.3)	φαααα...	487990.0(6.5)
	Παααα...	431933.3(9.6)	φφααα...	983416.5(12.6)
	ΦΠααα...	1712943.0(18.4)	ΠΠααα...	734284.6(26.2)
30	φφφαα...	1023590.0(23.7)	φφΠαα...	2592866.0(33.3)
	φΠΠαα...	2126605.0(45.6)	ΠΠΠαα...	558519.0(59.9)
	φφφφα...	563952.6(41.8)	φφφΠα...	1800481.0(55.6)
	φφΠΠα...	2052433.0(70.4)	φΠΠΠα...	978420.5(83.9)
	ΠΠΠΠα...	163640.3(93.5)	φφφφφ...	148719.7(66.1)
35	φφφφΠ...	541755.7(80.3)	φφφΠΠ...	738960.1(91.2)
	φφΠΠΠ...	473377.0(97.4)	φΠΠΠΠ...	145189.7(99.6)
	ΠΠΠΠΠ...	17491.3(100.0)	φφφφφ...	13829.1(88.5)
	φφφφφ...	54058.1(96.1)	φφφφΠ...	83726.0(99.2)
	φφφΠΠ...	67454.5(99.9)	φφΠΠΠ...	30374.5(100.0)
40	φΠΠΠΠ...	7290.0(100.0)	ΠΠΠΠΠ...	729.0(100.0)

Table 130: Sampling of a Library encoded by (NNK)⁶
(continued)

Library size =		7.6000E+07	
5	total = 3.2125E+07		% sampled = 50.19
	αααααα...	245057.8(8.2)	ϕααααα... 1175010.0(15.7)
	ηααααα...	1014733.0(22.7)	ϕηαααα... 2255280.0(29.0)
10	ϕηαααα...	3749112.0(40.2)	ηηαααα... 1504128.0(53.7)
	ϕϕαααα...	2142478.0(49.6)	ϕϕηααα... 4993247.0(64.2)
	ϕηηααα...	3666785.0(78.6)	ηηηααα... 840691.9(90.1)
	ϕϕϕααα...	1007002.0(74.6)	ϕϕϕηαα... 2825063.0(87.2)
	ϕϕηηαα...	2782358.0(95.4)	ϕηηηαα... 1154956.0(99.0)
15	ηηηηαα...	174790.0(99.9)	ϕϕϕϕαα... 210475.6(93.5)
	ϕϕϕϕηα...	663929.3(98.4)	ϕϕϕηηα... 808298.6(99.8)
	ϕϕηηηα...	485953.2(100.0)	ϕηηηηα... 145799.9(100.0)
	ηηηηηα...	17496.0(100.0)	ϕϕϕϕϕα... 15559.9(99.6)
	ϕϕϕϕϕη...	56234.9(100.0)	ϕϕϕϕηη... 84374.6(100.0)
20	ϕϕϕηηη...	67500.0(100.0)	ϕϕηηηη... 30375.0(100.0)
	ϕηηηηη...	7290.0(100.0)	ηηηηηη... 729.0(100.0)
Library size =		1.0000E+08	
25	total = 3.6537E+07		% sampled = 57.09
	αααααα...	318185.1(10.7)	ϕααααα... 1506161.0(20.2)
	ηααααα...	1284677.0(28.7)	ϕηαααα... 2821285.0(36.3)
	ϕηαααα...	4585163.0(49.1)	ηηαααα... 1783932.0(63.7)
30	ϕϕαααα...	2566085.0(59.4)	ϕϕηααα... 5764391.0(74.1)
	ϕηηααα...	4051713.0(86.8)	ηηηααα... 888584.3(95.2)
	ϕϕϕααα...	1127473.0(83.5)	ϕϕϕηαα... 3023170.0(93.3)
	ϕϕηηαα...	2865517.0(98.3)	ϕηηηαα... 1163743.0(99.8)
	ηηηηαα...	174941.0(100.0)	ϕϕϕϕαα... 218886.6(97.3)
35	ϕϕϕϕηα...	671976.9(99.6)	ϕϕϕηηα... 809757.3(100.0)
	ϕϕηηηα...	485997.5(100.0)	ϕηηηηα... 145800.0(100.0)
	ηηηηηα...	17496.0(100.0)	ϕϕϕϕϕα... 15613.5(99.9)
	ϕϕϕϕϕη...	56248.9(100.0)	ϕϕϕϕηη... 84375.0(100.0)
	ϕϕϕηηη...	67500.0(100.0)	ϕϕηηηη... 30375.0(100.0)
40	ϕηηηηη...	7290.0(100.0)	ηηηηηη... 729.0(100.0)

Table 130: Sampling of a Library encoded by (NNK)⁶
(continued)

Library size = 3.0000E+08

5

total = 5.2634E+07 % sampled = 82.24

	ααααα... 856451.3(28.7)	ϕαααα... 3668130.0(49.1)
	Ωαααα... 2854291.0(63.7)	ϕϕααα... 5764391.0(74.1)
10	ϕΩααα... 8103426.0(86.8)	ΩΩααα... 2665753.0(95.2)
	ϕϕϕαα... 4030893.0(93.3)	ϕϕΩαα... 7641378.0(98.3)
	ϕΩΩαα... 4654972.0(99.8)	ΩΩΩαα... 933018.6(100.0)
	ϕϕϕϕα... 1343954.0(99.6)	ϕϕϕΩα... 3239029.0(100.0)
	ϕϕΩΩα... 2915985.0(100.0)	ϕΩΩΩα... 1166400.0(100.0)
15	ΩΩΩΩα... 174960.0(100.0)	ϕϕϕϕϕ... 224995.5(100.0)
	ϕϕϕϕΩ... 674999.9(100.0)	ϕϕϕΩΩ... 810000.0(100.0)
	ϕϕΩΩΩ... 486000.0(100.0)	ϕΩΩΩΩ... 145800.0(100.0)
	ΩΩΩΩΩ... 17496.0(100.0)	ϕϕϕϕϕϕ... 15625.0(100.0)
	ϕϕϕϕϕΩ... 56250.0(100.0)	ϕϕϕϕΩΩ... 84375.0(100.0)
20	ϕϕϕΩΩΩ... 67500.0(100.0)	ϕϕΩΩΩΩ... 30375.0(100.0)
	ϕΩΩΩΩΩ... 7290.0(100.0)	ΩΩΩΩΩΩ... 729.0(100.0)

Library size = 1.0000E+09

25

total = 6.1999E+07 % sampled = 96.87

	ααααα... 2018278.0(67.6)	ϕαααα... 6680917.0(89.5)
	Ωαααα... 4326519.0(96.6)	ϕϕααα... 7690221.0(98.9)
30	ϕΩααα... 9320389.0(99.9)	ΩΩααα... 2799250.0(100.0)
	ϕϕϕαα... 4319475.0(100.0)	ϕϕΩαα... 7775990.0(100.0)
	ϕΩΩαα... 4665600.0(100.0)	ΩΩΩαα... 933120.0(100.0)
	ϕϕϕϕα... 1350000.0(100.0)	ϕϕϕΩα... 3240000.0(100.0)
	ϕϕΩΩα... 2916000.0(100.0)	ϕΩΩΩα... 1166400.0(100.0)
35	ΩΩΩΩα... 174960.0(100.0)	ϕϕϕϕϕ... 225000.0(100.0)
	ϕϕϕϕΩ... 675000.0(100.0)	ϕϕϕΩΩ... 810000.0(100.0)
	ϕϕΩΩΩ... 486000.0(100.0)	ϕΩΩΩΩ... 145800.0(100.0)
	ΩΩΩΩΩ... 17496.0(100.0)	ϕϕϕϕϕϕ... 15625.0(100.0)
	ϕϕϕϕϕΩ... 56250.0(100.0)	ϕϕϕϕΩΩ... 84375.0(100.0)
40	ϕϕϕΩΩΩ... 67500.0(100.0)	ϕϕΩΩΩΩ... 30375.0(100.0)
	ϕΩΩΩΩΩ... 7290.0(100.0)	ΩΩΩΩΩΩ... 729.0(100.0)

Table 130: Sampling of a Library encoded by (NNK)⁶
(continued)

Library size = 3.0000E+09

5

total = 6.3890E+07 % sampled = 99.83

	ααααα... 2884346.0(96.6)	ϕαααα... 7456311.0(99.9)
	ηαααα... 4478800.0(100.0)	ϕϕααα... 7775990.0(100.0)
10	ϕηααα... 9331200.0(100.0)	ηηααα... 2799360.0(100.0)
	ϕϕααα... 4320000.0(100.0)	ϕϕηαα... 7776000.0(100.0)
	ϕηηαα... 4665600.0(100.0)	ηηηαα... 933120.0(100.0)
	ϕϕϕαα... 1350000.0(100.0)	ϕϕϕηα... 3240000.0(100.0)
	ϕϕηηα... 2916000.0(100.0)	ϕηηηα... 1166400.0(100.0)
15	ηηηηα... 174960.0(100.0)	ϕϕϕϕα... 225000.0(100.0)
	ϕϕϕηη... 675000.0(100.0)	ϕϕϕηη... 810000.0(100.0)
	ϕϕηηη... 486000.0(100.0)	ϕηηηη... 145800.0(100.0)
	ηηηηη... 17496.0(100.0)	ϕϕϕϕϕ... 15625.0(100.0)
	ϕϕϕϕη... 56250.0(100.0)	ϕϕϕϕη... 84375.0(100.0)
20	ϕϕϕηηη... 67500.0(100.0)	ϕϕηηηη... 30375.0(100.0)
	ϕηηηηη... 7290.0(100.0)	ηηηηηη... 729.0(100.0)

FOI b7E b7C b7D b7F b7G b7H b7I b7J b7K b7L b7M b7N b7O b7P b7Q b7R b7S b7T b7U b7V b7W b7X b7Y b7Z

Table 130, continued

D. Formulae for tabulated quantities.

- 5 Lsize is the number of independent transformants.
 31**6 is 31 to sixth power; 6*3 means 6 times 3.
 $A = \text{Lsize}/(31^{**6})$
 α can be one of [WMFYCIKDENVHQ.]
 Φ can be one of [PTAVG]
- 10 Ω can be one of [SLR]
 $F0 = (12)^{**6}$ $F1 = (12)^{**5}$ $F2 = (12)^{**4}$
 $F3 = (12)^{**3}$ $F4 = (12)^{**2}$ $F5 = (12)$
 $F6 = 1$
- 15 $\alpha\alpha\alpha\alpha\alpha\alpha = F0 * (1-\exp(-A))$
 $\Phi\alpha\alpha\alpha\alpha\alpha = 6 * 5 * F1 * (1-\exp(-2*A))$
 $\Omega\alpha\alpha\alpha\alpha\alpha = 6 * 3 * F1 * (1-\exp(-3*A))$
 $\Phi\Phi\alpha\alpha\alpha\alpha = (15) * 5^{**2} * F2 * (1-\exp(-4*A))$
 $\Phi\Omega\alpha\alpha\alpha\alpha = (6*5)*5*3 * F2 * (1-\exp(-6*A))$
- 20 $\Omega\Omega\alpha\alpha\alpha\alpha = (15) * 3^{**2} * F2 * (1-\exp(-9*A))$
 $\Phi\Phi\Phi\alpha\alpha\alpha = (20)*(5^{**3}) * F3 * (1-\exp(-8*A))$
 $\Phi\Phi\Omega\alpha\alpha\alpha = (60)*(5*5*3)*F3 * (1-\exp(-12*A))$
 $\Phi\Omega\Omega\alpha\alpha\alpha = (60)*(5*3*3)*F3*(1-\exp(-18*A))$
 $\Omega\Omega\Omega\alpha\alpha\alpha = (20)*(3)^{**3}*F3*(1-\exp(-27*A))$
- 25 $\Phi\Phi\Phi\Phi\alpha\alpha = (15)*(5)^{**4}*F4*(1-\exp(-16*A))$
 $\Phi\Phi\Phi\Omega\alpha\alpha = (60)*(5)^{**3}*3*F4*(1-\exp(-24*A))$
 $\Phi\Phi\Omega\Omega\alpha\alpha = (90)*(5*5*3*3)*F4*(1-\exp(-36*A))$
 $\Phi\Omega\Omega\Omega\alpha\alpha = (60)*(5*3*3*3)*F4*(1-\exp(-54*A))$
 $\Omega\Omega\Omega\Omega\alpha\alpha = (15)*(3)^{**4} * F4 * (1-\exp(-81*A))$
- 30 $\Phi\Phi\Phi\Phi\Phi\alpha = (6)*(5)^{**5} * F5 * (1-\exp(-32*A))$
 $\Phi\Phi\Phi\Phi\Omega\alpha = 30*5*5*5*5*3*F5*(1-\exp(-48*A))$
 $\Phi\Phi\Phi\Omega\Omega\alpha = 60*5*5*5*3*3*F5*(1-\exp(-72*A))$
 $\Phi\Phi\Omega\Omega\Omega\alpha = 60*5*5*3*3*3*F5*(1-\exp(-108*A))$
 $\Phi\Omega\Omega\Omega\Omega\alpha = 30*5*3*3*3*3*F5*(1-\exp(-162*A))$
- 35 $\Omega\Omega\Omega\Omega\Omega\alpha = 6*3*3*3*3*3*F5*(1-\exp(-243*A))$
 $\Phi\Phi\Phi\Phi\Phi\Phi = 5^{**6} * (1-\exp(-64*A))$
 $\Phi\Phi\Phi\Phi\Phi\Omega = 6*3*5^{**5}*(1-\exp(-96*A))$
 $\Phi\Phi\Phi\Phi\Omega\Omega = 15*3*3*5^{**4}*(1-\exp(-144*A))$
 $\Phi\Phi\Phi\Omega\Omega\Omega = 20*3^{**3}*5^{**3}*(1-\exp(-216*A))$
- 40 $\Phi\Phi\Omega\Omega\Omega\Omega = 15*3^{**4}*5^{**2}*(1-\exp(-324*A))$
 $\Phi\Omega\Omega\Omega\Omega\Omega = 6*3^{**5}*5*(1-\exp(-486*A))$
 $\Omega\Omega\Omega\Omega\Omega\Omega = 3^{**6}*(1-\exp(-729*A))$
 total = $\alpha\alpha\alpha\alpha\alpha\alpha + \Phi\alpha\alpha\alpha\alpha\alpha + \Omega\alpha\alpha\alpha\alpha\alpha + \Phi\Phi\alpha\alpha\alpha\alpha + \Phi\Omega\alpha\alpha\alpha\alpha +$
 $\Omega\Omega\alpha\alpha\alpha\alpha + \Phi\Phi\Phi\alpha\alpha\alpha + \Phi\Phi\Omega\alpha\alpha\alpha + \Phi\Omega\Omega\alpha\alpha\alpha + \Omega\Omega\Omega\alpha\alpha\alpha +$
 $\Phi\Phi\Phi\Phi\alpha\alpha + \Phi\Phi\Phi\Omega\alpha\alpha + \Phi\Phi\Omega\Omega\alpha\alpha + \Phi\Omega\Omega\Omega\alpha\alpha + \Omega\Omega\Omega\Omega\alpha\alpha +$
 $\Phi\Phi\Phi\Phi\Phi\alpha + \Phi\Phi\Phi\Phi\Omega\alpha + \Phi\Phi\Phi\Omega\Omega\alpha + \Phi\Phi\Omega\Omega\Omega\alpha + \Phi\Omega\Omega\Omega\Omega\alpha +$
 $\Omega\Omega\Omega\Omega\Omega\alpha + \Phi\Phi\Phi\Phi\Phi\Phi + \Phi\Phi\Phi\Phi\Phi\Omega + \Phi\Phi\Phi\Phi\Omega\Omega + \Phi\Phi\Phi\Omega\Omega\Omega +$
 $\Phi\Phi\Omega\Omega\Omega\Omega + \Phi\Omega\Omega\Omega\Omega\Omega + \Omega\Omega\Omega\Omega\Omega\Omega$
- 45

Table 131: Sampling of a Library
Encoded by (NNT)⁴(NNG)²

X can be F,S,Y,C,L,P,H,R,I,T,N,V,A,D,G

5

Γ can be L²,R²,S,W,P,Q,M,T,K,V,A,E,G

Library comprises $8.55 \cdot 10^6$ amino-acid sequences; $1.47 \cdot 10^7$ DNA sequences.

10

Total number of possible aa sequences= 8,555,625

15 x LVPTARGFYCHIND
 S S
 θ VPTAGWQMKES
 Ω LR

20 The first, second, fifth, and sixth positions can
hold x or S; the third and fourth position can hold θ or
Ω. I have lumped sequences by the number of xs, Ss, θs,
and Ωs.

For example xxθNSS stands for:

25 [xxθNSS, xSθNxs, xSθNSx, SSθNxx, SxθNxs, SxθNSx,
 xxNθSS, xSθNxs, xSθNSx, SSθNxx, SxθNxs, SxθNSx]

30 The following table shows the likelihood that any
particular DNA sequence will fall into one of the defined
classes.

	Library size =	1.0	Sampling =	.00001%
	total.....	1.0000E+00	%sampled.....	1.1688E-07
35	xxθθxx.....	3.1524E-01	xxθNxx.....	2.2926E-01
	xxΩNxx.....	4.1684E-02	xxθθxs.....	1.8013E-01
	xxθNxs.....	1.3101E-01	xxΩNxs.....	2.3819E-02
	xxθθSS.....	3.8600E-02	xxθNSS.....	2.8073E-02
	xxΩNSS.....	5.1042E-03	xSθθSS.....	3.6762E-03
40	xSθNSS.....	2.6736E-03	xSΩNSS.....	4.8611E-04
	SSθθSS.....	1.3129E-04	SSθNSS.....	9.5486E-05
	SSΩNSS.....	1.7361E-05		

Table 131: Sampling of a Library
 Encoded by (NNT)⁴(NNG)²
 (continued)

5 The following sections show how many sequences of
 each class are expected for libraries of different sizes.

Library size = 1.0000E+05

10	total.....		9.9137E+04	fraction sampled = 1.1587E-02		
	Type	Number	%	Type	Number	%
	xx00xx.....	31416.9(.7)	xx00xx.....	22771.4(1.3)
	xx00xx.....	4112.4(2.7)	xx00xs.....	17891.8(1.3)
15	xx00xs.....	12924.6(2.7)	xx00xs.....	2318.5(5.3)
	xx00ss.....	3808.1(2.7)	xx00ss.....	2732.5(5.3)
	xx00ss.....	483.7(10.3)	xs00ss.....	357.8(5.3)
	xs00ss.....	253.4(10.3)	xs00ss.....	43.7(19.5)
	ss00ss.....	12.4(10.3)	ss00ss.....	8.6(19.5)
20	ss00ss.....	1.4(35.2)			

Library size = 1.0000E+06

	total.....	9.2064E+05	fraction sampled = 1.0761E-01			
25	xx00xx.....	304783.9(6.6)	xx00xx.....	214394.0(12.7)		
	xx00xx.....	36508.6(23.8)	xx00xs.....	168452.5(12.7)		
	xx00xs.....	114741.4(23.8)	xx00xs.....	18383.8(41.9)		
	xx00ss.....	33807.7(23.8)	xx00ss.....	21666.6(41.9)		
	xx00ss.....	3114.6(66.2)	xs00ss.....	2837.3(41.9)		
30	xs00ss.....	1631.5(66.2)	xs00ss.....	198.4(88.6)		
	ss00ss.....	80.1(66.2)	ss00ss.....	39.0(88.6)		
	ss00ss.....	3.9(98.7)				

Library size = 3.0000E+06

35	total.....	2.3880E+06	fraction sampled = 2.7912E-01
	xx00xx.....	855709.5(18.4)	xx00xx..... 565051.6(33.4)
	xx00xx.....	85564.7(55.7)	xx00xs..... 443969.1(33.4)
	xx00xs.....	268917.8(55.7)	xx00xs..... 35281.3(80.4)
40	xx00ss.....	79234.7(55.7)	xx00ss..... 41581.5(80.4)
	xx00ss.....	4522.6(96.1)	xs00ss..... 5445.2(80.4)
	xs00ss.....	2369.0(96.1)	xs00ss..... 223.7(99.9)
	ss00ss.....	116.3(96.1)	ss00ss..... 43.9(99.9)
	ss00ss.....	4.0(100.0)	

TABLE 131: SAMPLING OF A LIBRARY

Table 131: Sampling of a Library
 Encoded by (NNT)⁴(NNG)²
 (continued)

5	Library size = 8.5556E+06	
	total..... 4.9303E+06 fraction sampled = 5.7626E-01	
	xx00xx..... 2046301.0(44.0)	xx00xx..... 1160645.0(68.7)
	xx00xx..... 138575.9(90.2)	xx00xs..... 911935.6(68.7)
10	xx00xs..... 435524.3(90.2)	xx00xs..... 43480.7(99.0)
	xx00ss..... 128324.1(90.2)	xx00ss..... 51245.1(99.0)
	xx00ss..... 4703.6(100.0)	xs00ss..... 6710.7(99.0)
	xs00ss..... 2463.8(100.0)	xs00ss..... 224.0(100.0)
	ss00ss..... 121.0(100.0)	ss00ss..... 44.0(100.0)
15	ss00ss..... 4.0(100.0)	
	Library size = 1.0000E+07	
	total..... 5.3667E+06 fraction sampled = 6.2727E-01	
20	xx00xx..... 2289093.0(49.2)	xx00xx..... 1254877.0(74.2)
	xx00xx..... 143467.0(93.4)	xx00xs..... 985974.9(74.2)
	xx00xs..... 450896.3(93.4)	xx00xs..... 43710.7(99.6)
	xx00ss..... 132853.4(93.4)	xx00ss..... 51516.1(99.6)
	xx00ss..... 4703.9(100.0)	xs00ss..... 6746.2(99.6)
25	xs00ss..... 2464.0(100.0)	xs00ss..... 224.0(100.0)
	ss00ss..... 121.0(100.0)	ss00ss..... 44.0(100.0)
	ss00ss..... 4.0(100.0)	
	Library size = 3.0000E+07	
30	total..... 7.8961E+06 fraction sampled = 9.2291E-01	
	xx00xx..... 4040589.0(86.9)	xx00xx..... 1661409.0(98.3)
	xx00xx..... 153619.1(100.0)	xx00xs..... 1305393.0(98.3)
	xx00xs..... 482802.9(100.0)	xx00xs..... 43904.0(100.0)
35	xx00ss..... 142254.4(100.0)	xx00ss..... 51744.0(100.0)
	xx00ss..... 4704.0(100.0)	xs00ss..... 6776.0(100.0)
	xs00ss..... 2464.0(100.0)	xs00ss..... 224.0(100.0)
	ss00ss..... 121.0(100.0)	ss00ss..... 44.0(100.0)
	ss00ss..... 4.0(100.0)	

009965 55096860

Table 131: Sampling of a Library
 Encoded by (NNT)⁴(NNG)²
 (continued)

5	Library size = 5.0000E+07	
	total.....	8.3956E+06 fraction sampled = 9.8130E-01
	xx00xx.....	4491779.0(96.6) xx00xx..... 1688387.0(99.9)
	xx00xx.....	153663.8(100.0) xx00xs..... 1326590.0(99.9)
10	xx00xs.....	482943.4(100.0) xx00ss..... 43904.0(100.0)
	xx00ss.....	142295.8(100.0) xx00ss..... 51744.0(100.0)
	xx00ss.....	4704.0(100.0) xs00ss..... 6776.0(100.0)
	xs00ss.....	2464.0(100.0) xs00ss..... 224.0(100.0)
	ss00ss.....	121.0(100.0) ss00ss..... 44.0(100.0)
15	ss00ss.....	4.0(100.0)
	Library size = 1.0000E+08	
	total.....	8.5503E+06 fraction sampled = 9.9938E-01
20	xx00xx.....	4643063.0(99.9) xx00xx..... 1690302.0(100.0)
	xx00xx.....	153664.0(100.0) xx00xs..... 1328094.0(100.0)
	xx00xs.....	482944.0(100.0) xx00xs..... 43904.0(100.0)
	xx00ss.....	142296.0(100.0) xx00ss..... 51744.0(100.0)
	xx00ss.....	4704.0(100.0) xs00ss..... 6776.0(100.0)
25	xs00ss.....	2464.0(100.0) xs00ss..... 224.0(100.0)
	ss00ss.....	121.0(100.0) ss00ss..... 44.0(100.0)
	ss00ss.....	4.0(100.0)

106250 5695550

Table 132: Relative efficiencies of various simple variegation codons

5	Number of codons		
	5	6	7
	#DNA/#AA	#DNA/#AA	#DNA/#AA
	[#DNA]	[#DNA]	[#DNA]
vgCodon	(#AA)	(#AA)	(#AA)
10 NNK	8.95	13.86	21.49
assuming	$[2.86 \cdot 10^7]$	$[8.87 \cdot 10^8]$	$[2.75 \cdot 10^{10}]$
stops vanish	$(3.2 \cdot 10^6)$	$(6.4 \cdot 10^7)$	$(1.28 \cdot 10^9)$
NNT	1.38	1.47	1.57
15	$[1.05 \cdot 10^6]$	$[1.68 \cdot 10^7]$	$[2.68 \cdot 10^8]$
	$(7.59 \cdot 10^5)$	$(1.14 \cdot 10^7)$	$(1.71 \cdot 10^8)$
NNG	2.04	2.36	2.72
assuming	$[7.59 \cdot 10^5]$	$[1.14 \cdot 10^6]$	$[1.71 \cdot 10^8]$
20 stops vanish	$(3.7 \cdot 10^5)$	$(4.83 \cdot 10^6)$	$(6.27 \cdot 10^7)$

T06230" 56096850

Table 140. Affect of anti BPTI IgG on phage titer.

Phage Strain	Input	+Anti-BPTI	+Anti-BPTI +Protein A(a)	Eluted Phage
M13MP18	100 (b)	98	92	$7 \cdot 10^{-4}$
5 BPTI.3	100	26	21	6
M13MB48 (c)	100	90	36	0.8
M13MB48 (d)	100	60	40	2.6
(a)	Protein A-agarose beads.			
(b)	Percentage of input phage measured as plaque forming units			
(c)	Batch number 3			
(d)	Batch number 4			

Table 141. Affect of anti-BPTI or protein A on phage titer.

Strain	Input	No Addition	+Anti-BPTI	+Protein A (a)	+Anti-BPTI +Protein A
M13MP18	100(b)	107	105	72	65
M13MB48(b)	100	92	$7 \cdot 10^{-3}$	58	$<10^{-4}$
(a)	Protein A-agarose beads				
(b)	Percentage of input phage measured as plaque forming units				
(c)	Batch number 5				

Table 142 Affect of anti-BPTI and non-immune serum on phage titer

5	Strain	Input	+Anti-BPTI	+NRS (a)	+Anti-BPTI +Protein A (b)	+NRS +Protein A
	M13MP18	100(c)	65	104	71	88
	M13MB48(d)	100	30	125	13	121
10	M13MB48(e)	100	2	105	0.7	110
	(a)	Purified IgG from normal rabbit serum.				
	(b)	Protein A-agarose beads.				
	(c)	Percentage of input phage measured as plaque forming units				
15	(d)	Batch number 4				
	(e)	Batch number 5				

T06290 55036260

Table 143. Loss in titer of display phage with anhydrotrypsin.

5	Strain	Anhydrotrypsin Beads		Streptavidin Beads	
		Start	Post Incubation	Start	Post Incubation
	M13MP18	100 (a)	121	ND	ND
	M13MB48	100	58	100	98
	5AA Pool	100	44	100	93

10

(a) Plaque forming units expressed as a percentage of input.

Table 144. Binding of Display Phage to Anhydrotrypsin.

15

Experiment 1.

20	Strain	Eluted Phage (a)	Relative to
			M13MP18
	M13MP18	0.2 (a)	1.0
	BPTI-IIIIMK	7.9	39.5
	M13MB48	11.2	56.0

Experiment 2.

25

30	Strain	Eluted Phage (a)	Relative to
			M13mp18
	M13mp18	0.3	1.0
	BPTI-IIIIMK	12.0	40.0
	M13MB56	17.0	56.7

(a) Plaque forming units acid eluted from beads, expressed as a percentage of the input.

0336093 062904
106290 55036360

Table 145. Binding of Display Phage to Anhydrotrypsin or Trypsin.

5	Strain	Anhydrotrypsin Beads		Trypsin Beads	
		Eluted Phage (a)	Relative Binding (b)	Eluted Phage	Relative Binding
	M13MP18	0.1	1	2.3×10^{-4}	1.0
	BPTI-IIIIMK	9.1	91	1.17	5×10^3
10	M13.3X7	25.0	250	1.4	6×10^3
	M13.3X11	9.2	92	0.27	1.2×10^3

(a) Plaque forming units eluted from beads, expressed as a percentage of the input.

15 (b) Relative to the non-display phage, M13MP18.

Table 146. Binding of Display Phage to Trypsin or Human Neutrophil Elastase.

20	Strain	Trypsin Beads		HNE Beads	
		Eluted Phage (a)	Relative Binding(b)	Eluted Phage	Relative Binding
25	M13MP18	5×10^{-4}	1	3×10^{-4}	1.0
	BPTI-IIIIMK	1.0	2000	5×10^{-3}	16.7
	M13MB48	0.13	260	9×10^{-3}	30.0
	M13.3X7	1.15	2300	1×10^{-3}	3.3
	M13.3X11	0.8	1600	2×10^{-3}	6.7
30	BPTI3.CL	1×10^{-3}	2	4.1	1.4×10^4

(c) .

(a) Plaque forming units acid eluted from the beads, expressed as a percentage of input.

35 (b) Relative to the non-display phage, M13MP18.

(C) BPTI-IIIMK (K15L MGNG)

Table 155
Distance in Å between alpha carbons in octapeptides:

5 Extended Strand: angle of $C_{\alpha}1-C_{\alpha}2-C_{\alpha}3 = 138^\circ$

	1	2	3	4	5	6	7	8
1	-							
2	3.8	-						
10 3	7.1	3.8	-					
4	10.7	7.1	3.8	-				
5	14.2	10.7	7.1	3.8	-			
6	17.7	14.1	10.7	7.1	3.8	-		
7	21.2	17.7	14.1	10.6	7.0	3.8	-	
15 8	24.6	20.9	17.5	13.9	10.6	7.0	3.8	-

Reverse turn between residues 4 and 5.

	1	2	3	4	5	6	7	8
1	-							
2	3.8	-						
3	7.1	3.8	-					
4	10.6	7.0	3.8	-				
25 5	11.6	8.0	6.1	3.8	-			
6	9.0	5.8	5.5	5.6	3.8	-		
7	6.2	4.1	6.3	8.0	7.0	3.8	-	
8	5.8	6.0	9.1	11.6	10.7	7.2	3.8	-

30 Alpha helix: angle of $C_{\alpha}1-C_{\alpha}2-C_{\alpha}3 = 93^\circ$

	1	2	3	4	5	6	7	8
1	-							
2	3.8	-						
35 3	5.5	3.8	-					
4	5.1	5.4	3.8	-				
5	6.6	5.3	5.5	3.8	-			
6	9.3	7.0	5.6	5.5	3.8	-		
40 7	10.4	9.3	6.9	5.4	5.5	3.8	-	
8	11.3	10.7	9.5	6.8	5.6	5.6	3.8	-

105250" 55035850

Table 156

Distances between alpha carbons in closed mini-proteins of
the form disulfide cyclo(CXXXXC)

Minimum distance

		1	2	3	4	5	6
10	1	-					
	2	3.8	-				
	3	5.9	3.8	-			
	4	5.6	6.0	3.8	-		
15	5	4.7	5.9	6.0	3.8	-	
	6	4.8	5.3	5.1	5.2	3.8	-

20 Average distance

		1	2	3	4	5	6
25	1	-					
	2	3.8	-				
	3	6.3	3.8	-			
	4	7.5	6.4	3.8	-		
	5	7.1	7.5	6.3	3.8	-	
	6	5.6	7.5	7.7	6.4	3.8	-

30

Maximum distance

		1	2	3	4	5	6
35	1	-					
	2	3.8	-				
	3	6.7	3.8	-			
	4	9.0	6.9	3.8	-		
	5	8.7	8.8	6.8	3.8	-	
	6	6.6	9.2	9.1	6.8	3.8	-

40

T06230-5605650

Table 160: pH Profile of BPTI-III MK phage and Shad 1 phage binding to Cat G beads.

BPTI-IIIIMK			
5	pH	Total pfu in Fraction	Percentage of Input
	7	3.7×10^5	3.7×10^{-2}
	6	3.1×10^5	3.1×10^{-2}
	5	1.4×10^5	1.4×10^{-2}
	4.5	3.1×10^4	3.1×10^{-3}
10	4	7.1×10^3	7.1×10^{-4}
	3.5	2.6×10^3	2.6×10^{-4}
	3	2.5×10^3	2.5×10^{-4}
	2.5	8.8×10^2	8.8×10^{-5}
	2	7.6×10^2	7.6×10^{-5}
15	(total input = 1×10^9 phage)		
	Shad 1		
	7	2.5×10^5	1.1×10^{-2}
	6	6.3×10^4	2.7×10^{-3}
20	5	7.4×10^4	3.1×10^{-3}
	4.5	7.1×10^4	3.0×10^{-3}
	4	4.1×10^4	1.7×10^{-3}
	3.5	3.3×10^4	1.4×10^{-3}
	3	2.5×10^3	1.1×10^{-4}
25	2.5	1.4×10^4	5.7×10^{-4}
	2	5.2×10^3	2.2×10^{-4}
	(total input = 2.35×10^8 phage).		

106250" 5605650

5

10

The total input for BPTI-III MK phage was $1.85 \cdot 10^{10}$ plaque-forming units while the input for MK phage was $4.65 \cdot 10^{11}$ plaque-forming units.

TABLE 202
Elution of BPTI-III MK and BPTI(K15L)-III MA Phage from
Immobilized Trypsin and HNE

5				
	Type of Phage	Immobil- ized Protease	Total Plaque- Forming Units in Elution Fraction	Percentage of Input Phage Recovered
10				
	BPTI-III MK	Trypsin	$2.1 \cdot 10^7$	$4.1 \cdot 10^{-1}$
15	BPTI-III MK	HNE	$2.6 \cdot 10^5$	$5 \cdot 10^{-3}$
	BPTI(K15L)- III MA	Trypsin	$5.2 \cdot 10^4$	$5 \cdot 10^{-3}$
20	BPTI(K15L)- III MA	HNE	$1.0 \cdot 10^6$	$1.0 \cdot 10^{-1}$
25	The total input of BPTI-III MK phage was $5.1 \cdot 10^9$ pfu and the input of BPTI(K15L)-III MA phage was $9.6 \cdot 10^8$ pfu.			

135230" 56095850

TABLE 203

Effect of pH on the Dissociation of
Bound BPTI-III MK and
BPTI(K15L)-III MA Phage from Immobilized HNE

pH	BPTI-III MK		BPTI(K15L)-III MA	
	Total Plaque- Forming Units in Fraction	% of Input Phage	Total Plaque- Forming Units in Fraction	% of Input Phage
7.0	$5.0 \cdot 10^4$	$2 \cdot 10^{-3}$	$1.7 \cdot 10^5$	$3.2 \cdot 10^{-2}$
6.0	$3.8 \cdot 10^4$	$2 \cdot 10^{-3}$	$4.5 \cdot 10^5$	$8.6 \cdot 10^{-2}$
5.0	$3.5 \cdot 10^4$	$1 \cdot 10^{-3}$	$2.1 \cdot 10^6$	$4.0 \cdot 10^{-1}$
4.0	$3.0 \cdot 10^4$	$1 \cdot 10^{-3}$	$4.3 \cdot 10^6$	$8.2 \cdot 10^{-1}$
3.0	$1.4 \cdot 10^4$	$1 \cdot 10^{-3}$	$1.1 \cdot 10^6$	$2.1 \cdot 10^{-1}$
2.2	$2.9 \cdot 10^4$	$1 \cdot 10^{-3}$	$5.9 \cdot 10^4$	$1.1 \cdot 10^{-2}$
Percentage of Input Phage = $8.0 \cdot 10^{-3}$ Recovered Percentage of Input Phage = 1.56 Recovered				
The total input of BPTI-III MK phage was $0.030 \text{ ml} \times (8.6 \cdot 10^{10} \text{ pfu/ml}) = 2.6 \cdot 10^9$.				
The total input of BPTI(K15L)-III MA phage was $0.030 \text{ ml} \times (1.7 \cdot 10^{10} \text{ pfu/ml}) = 5.2 \cdot 10^8$.				
Given that the infectivity of BPTI(K15L)-III MA phage is 5 fold lower than that of BPTI-III MK phage, the phage inputs utilized above ensure that an equivalent number of phage particles are added to the immobilized HNE.				

TABLE 204

Effect of Mutation of Residues 39 to 42 of BPTI
on the ability of BPTI(K15L)-III MA to Bind to
Immobilized HNE

10	pH	BPTI(K15L)-III MA		BPTI(K15L, MGNG)-III MA	
		Total Plaque- Forming Units	% Input	Total Plaque- Forming Units	% Input
15	7.0	$3.0 \cdot 10^5$	$8.2 \cdot 10^{-2}$	$4.5 \cdot 10^5$	$1.63 \cdot 10^{-1}$
	6.0	$3.6 \cdot 10^5$	$1.00 \cdot 10^{-1}$	$6.3 \cdot 10^5$	$2.27 \cdot 10^{-1}$
	5.5	$5.3 \cdot 10^5$	$1.46 \cdot 10^{-1}$	$7.3 \cdot 10^5$	$2.64 \cdot 10^{-1}$
20	5.0	$5.6 \cdot 10^5$	$1.52 \cdot 10^{-1}$	$8.7 \cdot 10^5$	$3.16 \cdot 10^{-1}$
	4.75	$9.9 \cdot 10^5$	$2.76 \cdot 10^{-1}$	$1.3 \cdot 10^6$	$4.60 \cdot 10^{-1}$
25	4.5	$3.1 \cdot 10^5$	$8.5 \cdot 10^{-2}$	$3.6 \cdot 10^5$	$1.30 \cdot 10^{-1}$
	4.25	$5.2 \cdot 10^5$	$1.42 \cdot 10^{-1}$	$5.0 \cdot 10^5$	$1.80 \cdot 10^{-1}$
30	4.0	$5.1 \cdot 10^4$	$1.4 \cdot 10^{-2}$	$1.3 \cdot 10^5$	$4.8 \cdot 10^{-2}$
	3.5	$1.3 \cdot 10^4$	$4 \cdot 10^{-3}$	$3.8 \cdot 10^4$	$1.4 \cdot 10^{-2}$
35	Total Percentage = 1.00 Recovered		Total Percentage = 1.80 Recovered		

The total input of BPTI(K15L)-III MA phage was
 $0.030 \text{ ml} \times (1.2 \cdot 10^{10} \text{ pfu/ml}) = 3.6 \cdot 10^8 \text{ pfu}$.

40 The total input of BPTI(K15L, MGNG)-III MA phage was
 $0.030 \text{ ml} \times (9.2 \cdot 10^9 \text{ pfu/ml}) = 2.8 \cdot 10^8 \text{ pfu}$.

105230"56056350

TABLE 205

Fractionation of a Mixture of
BPTI-III MK and
BPTI(K15L,MGNG)-III MA Phage
on Immobilized HNE

pH	BPTI-III MK		BPTI(K15L,MGNG)-III MA	
	Total Kanamycin Transducing Units	% of Input	Total Ampicillin Transducing Units	% of Input
7.0	$4.01 \cdot 10^3$	$4.5 \cdot 10^{-3}$	$1.39 \cdot 10^5$	$3.13 \cdot 10^{-1}$
6.0	$7.06 \cdot 10^2$	$8 \cdot 10^{-4}$	$7.18 \cdot 10^4$	$1.62 \cdot 10^{-1}$
5.0	$1.81 \cdot 10^3$	$2.0 \cdot 10^{-3}$	$1.35 \cdot 10^5$	$3.04 \cdot 10^{-1}$
4.0	$1.49 \cdot 10^3$	$1.7 \cdot 10^{-3}$	$7.43 \cdot 10^5$	1.673

The total input of BPTI-III MK phage was
 $0.015 \text{ ml} \times (5.94 \cdot 10^9 \text{ kanamycin transducing units/ml}) =$
 $8.91 \cdot 10^7 \text{ kanamycin transducing units.}$

The total input of BPTI(K15L,MGNG)-III MA phage was
 $0.015 \text{ ml} \times (2.96 \cdot 10^9 \text{ ampicillin transducing units/ml}) =$
 $4.44 \cdot 10^7 \text{ ampicillin transducing units.}$

TABLE 206

5 Characterization of the Affinity of
BPTI(K15V,R17L)-III MA Phage for Immobilized HNE

10	pH	BPTI(K15V,R17L)-III MA		BPTI(K15L,MGNG)-III MA	
		Total Plaque- Forming Units Recovered	Percentage of Input Phage	Total Plaque- Forming Units Recovered	Percentage of Input Phage
15	7.0	$3.19 \cdot 10^6$	$8.1 \cdot 10^{-2}$	$9.42 \cdot 10^4$	$4.6 \cdot 10^{-2}$
	6.0	$5.42 \cdot 10^6$	$1.38 \cdot 10^{-1}$	$1.61 \cdot 10^5$	$7.9 \cdot 10^{-2}$
	5.0	$9.45 \cdot 10^6$	$2.41 \cdot 10^{-1}$	$2.85 \cdot 10^5$	$1.39 \cdot 10^{-1}$
20	4.5	$1.39 \cdot 10^7$	$3.55 \cdot 10^{-1}$	$4.32 \cdot 10^5$	$2.11 \cdot 10^{-1}$
	4.0	$2.02 \cdot 10^7$	$5.15 \cdot 10^{-1}$	$1.42 \cdot 10^5$	$6.9 \cdot 10^{-2}$
25	3.75	$9.20 \cdot 10^6$	$2.35 \cdot 10^{-1}$	-	-
	3.5	$4.16 \cdot 10^6$	$1.06 \cdot 10^{-1}$	$5.29 \cdot 10^4$	$2.6 \cdot 10^{-2}$
	3.0	$2.65 \cdot 10^6$	$6.8 \cdot 10^{-2}$	-	-
30	Total Input = 1.73 Recovered		Total Input = 0.57 Recovered		

35 Total input of BPTI(K15V,R17L)-III MA phage was
 $0.040 \text{ ml} \times (9.80 \cdot 10^{10} \text{ pfu/ml}) = 3.92 \cdot 10^9 \text{ pfu}.$

Total input of BPTI(K15L,MGNG)-III MA phage was
 $0.040 \text{ ml} \times (5.13 \cdot 10^9 \text{ pfu/ml}) = 2.05 \cdot 10^8 \text{ pfu}.$

0366095.02904
106230"5603650

TABLE 207

Sequence of the EpiNE α Clone Selected
From the Mini-Library

5

10

1	1	1	1	1	1	1	2	2
3	4	5	6	7	8	9	0	1
P	C	V	A	M	F	Q	R	Y
CCT	TGC	GTG	GCT	ATG	TTC	CAA	CGC	TAT

105230" 5603650

SEQUENCES OF THE EpINE CLONES IN THE P1 REGION

CLONE IDENTIFIERS		SEQUENCE									
10	3, 9, 16, 17, 18, 19	1 3 P	1 4 C	1 5 V	1 6 G	1 7 F	1 8 F	1 9 S	2 0 R	2 1 Y	EpiNE3
		CCT.TGC.GTC.GGT.TTC.TTC.TCA.CGC.TAT									
15	6	1 3 P	1 4 C	1 5 V	1 6 G	1 7 F	1 8 F	1 9 Q	2 0 R	2 1 Y	EpiNE6
		CCT.TGC.GTC.GGT.TTC.TTC.CAA.CGC.TAT									
20	7, 13, 14 15, 20	1 3 P	1 4 C	1 5 V	1 6 A	1 7 M	1 8 F	1 9 P	2 0 R	2 1 Y	EpiNE7
		CCT.TGC.GTC.GCT.ATG.TTC.CCA.CGC.TAT									
25	4	1 3 P	1 4 C	1 5 V	1 6 A	1 7 I	1 8 F	1 9 P	2 0 R	2 1 Y	EpiNE4
		CCT.TGC.GTC.GCT.ATC.TTC.CCA.CGC.TAT									
30	8	1 3 P	1 4 C	1 5 V	1 6 A	1 7 I	1 8 F	1 9 K	2 0 R	2 1 S	EpiNE8
		CCT.TGC.GTC.GCT.ATC.TTC.AAA.CGC.TCT									
35	1, 10 11, 12	1 3 P	1 4 C	1 5 I	1 6 A	1 7 F	1 8 F	1 9 P	2 0 R	2 1 Y	EpiNE1
		CCT.TGC.ATC.GCT.TTC.TTC.CCA.CGC.TAT									
40	5	1 3 P	1 4 C	1 5 I	1 6 A	1 7 F	1 8 F	1 9 Q	2 0 R	2 1 Y	EpiNE5
		CCT.TGC.ATC.GCT.TTC.TTC.CAA.CGC.TAT									
45	2	1 3 P	1 4 C	1 5 I	1 6 A	1 7 L	1 8 F	1 9 K	2 0 R	2 1 Y	EpiNE2
		CCT.TGC.ATC.GCT.TTG.TTC.AAA.CGC.TAT									

Table 209: DNA sequences and predicted amino acid sequences around the P1 region of BPTI analogues selected for binding to Cathepsin G.

5	Clone	P1				
		15	16	17	18	19
	BPTI	AAA . GCG . CGC . ATC . ATC				
		LYS ALA ARG ILE ILE				
10						
	Epic 1	ATG . GGT . TTC . TCC . AAA				
	(a)	MET GLY PHE SER LYS				
15	Epic 7	ATG . GCT . TTG . TTC . AAA				
		MET ALA LEU PHE LYS				
	Epic 8	TTC . GCT . ATC . ACC . CCA				
	(b)	PHE ALA ILE THR PRO				
20						
	Epic 10	ATG . GCT . TTG . TTC . CAA				
		MET ALA LEU PHE GLN				
	Epic 20	ATG . GCT . ATC . TCC . CCA				
25		MET ALA ILE SER PRO				

(a) Clones 11 and 31 also had the identical sequence.

(b) Clone 8 also contained the mutation Tyr 10 to ASN.

105250" 55095350

Table 210
 Derivatives of EpiNE7 Obtained
 by Variegation at positions 34, 36, 39, 40 and 41

5	EpiNE7	♦♦♦♦♦	♦♦♦♦♦
	RPDFCLEPPYTGPCvAmfpRYFYNAKAGLCQTFVYGGCmgngNNFKSAEDCMRTCGGA		
	1 2 3 4 5		
10	123456789012345678901234567890123456789012345678		
	EpiNE7.6	↓ ↓ ↓ ↓ ↓	♦ ♦ ♦ ♦ ↓
	RPDFCLEPPYTGPCvAmfpRYFYNAKAGLCQTFlyGcKgkGNNFKSAEDCMRTCGGA		
15	EpiNE7.8, EpiNE7.9, and EpiNE7.31		
	RPDFCLEPPYTGPCvAmfpRYFYNAKAGLCQTFeyGcWakGNNFKSAEDCMRTCGGA		
	EpiNE7.11		
20	RPDFCLEPPYTGPCvAmfpRYFYNAKAGLCQTFgYagCrakGNNFKSAEDCMRTCGGA		
	EpiNE7.7		
	RPDFCLEPPYTGPCvAmfpRYFYNAKAGLCQTFeyGcChaeGNNFKSAEDCMRTCGGA		
25	EpiNE7.4 and EpiNE7.14		
	RPDFCLEPPYTGPCvAmfpRYFYNAKAGLCQTFlyGcWaqGNNFKSAEDCMRTCGGA		
	EpiNE7.5		
30	RPDFCLEPPYTGPCvAmfpRYFYNAKAGLCQTFrYgGclaeGNNFKSAEDCMRTCGGA		
	EpiNE7.10 and EpiNE7.20		
	RPDFCLEPPYTGPCvAmfpRYFYNAKAGLCQTFdYgGchadGNNFKSAEDCMRTCGGA		
	EpiNE7.1		
35	RPDFCLEPPYTGPCvAmfpRYFYNAKAGLCQTFkyGclahGNNFKSAEDCMRTCGGA		
	EpiNE7.16		
	RPDFCLEPPYTGPCvAmfpRYFYNAKAGLCQTFtYgGcwanGNNFKSAEDCMRTCGGA		
40	EpiNE7.19		
	RPDFCLEPPYTGPCvAmfpRYFYNAKAGLCQTFnYgGcegkGNNFKSAEDCMRTCGGA		
	EpiNE7.12		
	RPDFCLEPPYTGPCvAmfpRYFYNAKAGLCQTFqYgGcegyGNNFKSAEDCMRTCGGA		
45	EpiNE7.17		
	RPDFCLEPPYTGPCvAmfpRYFYNAKAGLCQTFqYgGclgeGNNFKSAEDCMRTCGGA		
	EpiNE7.21		
50	RPDFCLEPPYTGPCvAmfpRYFYNAKAGLCQTFhYgGcwggGNNFKSAEDCMRTCGGA		

T05290"5696860

Table 210: Derivatives of EpiNE7 Obtained
by Variegation at positions 34, 36, 39, 40 and 41
(continued)

5	EpiNE7	◆◆◆◆	◆◆◆◆
	RPDFCLEPPYTGPCvAmfpRYFYNAKAGLCQTFVYGGCmgngNNFKSAEDCMRTCGGA		
	1 2 3 4 5		
	123456789012345678901234567890123456789012345678		
10	EpiNE7.22	↓↓↓↓↓	◆ ◆ ◆◆↓
	RPDFCLEPPYTGPCvAmfpRYFYNAKAGLCQTFhYgGCwgeGNNFKSAEDCMRTCGGA		
15	EpiNE7.23		
	RPDFCLEPPYTGPCvAmfpRYFYNAKAGLCQTFkYgGCwgkGNNFKSAEDCMRTCGGA		
	EpiNE7.24		
	RPDFCLEPPYTGPCvAmfpRYFYNAKAGLCQTFkYgGChgnGNNFKSAEDCMRTCGGA		
20	EpiNE7.25		
	RPDFCLEPPYTGPCvAmfpRYFYNAKAGLCQTFpYgGCwakGNNFKLAEDCMRTCGGA		
	EpiNE7.26		
25	RPDFCLEPPYTGPCvAmfpRYFYNAKAGLCQTFkYgGCwghGNNFKSAEDCMRTCGGA		
	EpiNE7.27		
	RPDFCLEPPYTGPCvAmfpRYFYNAKAGLCQTFnYgGCwgkGNNFKSAEDCMRTCGGA		
30	EpiNE7.28		
	RPDFCLEPPYTGPCvAmfpRYFYNAKAGLCQTFtYgGClghGNNFKSAEDCMRTCGGA		
	EpiNE7.29		
	RPDFCLEPPYTGPCvAmfpRYFYNAKAGLCQTFtYgGClgyGNNFKSAEDCMRTCGGA		
35	EpiNE7.30, EpiNE7.34, and EpiNE7.35		
	RPDFCLEPPYTGPCvAmfpRYFYNAKAGLCQTFkYgGCwaeGNNFKSAEDCMRTCGGA		
	EpiNE7.32		
40	RPDFCLEPPYTGPCvAmfpRYFYNAKAGLCQTFgYgGCwgeGNNFKSAEDCMRTCGGA		
	EpiNE7.33		
	RPDFCLEPPYTGPCvAmfpRYFYNAKAGLCQTFeYgGCwanGNNFKSAEDCMRTCGGA		
45	EpiNE7.36		
	RPDFCLEPPYTGPCvAmfpRYFYNAKAGLCQTFvYgGChgdGNNFKSAEDCMRTCGGA		
	EpiNE7.37		
	RPDFCLEPPYTGPCvAmfpRYFYNAKAGLCQTFmYgGCqgkGNNFKSAEDCMRTCGGA		
50	EpiNE7.38		
	RPDFCLEPPYTGPCvAmfpRYFYNAKAGLCQTFyYgGCwakGNNFKSAEDCMRTCGGA		

Table 210 (continued)
 Derivatives of EpiNE7 Obtained
 by Variegation at positions 34, 36, 39, 40 and 41

5

EpiNE7

♦♦♦♦♦

RPDFCLEPPYTGPCvAmfpRYFYNAKAGLCQTFVYGGCmgngNNFKSAEDCMRTC GGA

1

2

3

4

5

10123456789012345678901234567890123456789012345678

EpiNE7.39

↓↓↓↓↓

♦ ♦ ♦♦↓

RPDFCLEPPYTGPCvAmfpRYFYNAKAGLCQTFmYgGCwgdGNNFKSAEDCMRTC GGA

15

EpiNE7.40

RPDFCLEPPYTGPCvAmfpRYFYNAKAGLCQTFtYgGChgnGNNFKSAEDCMRTC GGA

T06250"55095860

Table 210: Derivatives of EpiNE7 Obtained
by Variegation at positions 34, 36, 39, 40 and 41
(continued)

5

Notes:

- a) ♦ indicates variegated residue. * indicates imposed change. ↓ indicates carry over from EpiNE7.
- 10 b) The sequence M₃₉-GNG in EpiNE7 (indicated by *) was imposed to increase similarity to ITI-D1.
- 15 b) Lower case letters in EpiNE7.6 to 7.38 indicate changes from BPTI that were selected in the first round (residues 15-19) or positions where the PBD was variegated in the second round (residues 34, 36, 39, 40, and 41).
- 20 c) All EpiNE7 derivatives have G₄₂.

106290" 55095350

TABLE 211

Effects of antisera on phage infectivity

5	Phage (dilution of stock)	Incubation Conditions	pfu/ml	Relative Titer
10	MA-ITI (10 ⁻¹)	PBS	1.2·10 ¹¹	1.00
		NRS	6.8·10 ¹⁰	0.57
		anti-ITI	1.1·10 ¹⁰	0.09
15	MA-ITI (10 ⁻³)	PBS	7.7·10 ⁸	1.00
		NRS	6.7·10 ⁸	0.87
		anti-ITI	8.0·10 ⁶	0.01
20	MA (10 ⁻¹)	PBS	1.3·10 ¹²	1.00
		NRS	1.4·10 ¹²	1.10
		anti-ITI	1.6·10 ¹²	1.20
25	MA (10 ⁻³)	PBS	1.3·10 ¹⁰	1.00
		NRS	1.2·10 ¹⁰	0.92
		anti-ITI	1.5·10 ¹⁰	1.20

T05250"56036360

TABLE 212

Fractionation of EpiNE-7 and MA-ITI phage on HNE beads

Sample	EpiNE-7		MA-ITI	
	Total pfu in sample	Fraction of input	Total pfu in sample	Fraction of input
INPUT	$3.3 \cdot 10^9$	1.00	$3.4 \cdot 10^{11}$	1.00
15Final TBS-TWEEN Wash	$3.8 \cdot 10^5$	$1.2 \cdot 10^{-4}$	$1.8 \cdot 10^6$	$5.3 \cdot 10^{-6}$
20pH 7.0	$6.2 \cdot 10^5$	$1.8 \cdot 10^{-4}$	$1.6 \cdot 10^6$	$4.7 \cdot 10^{-6}$
pH 6.0	$1.4 \cdot 10^6$	$4.1 \cdot 10^{-4}$	$1.0 \cdot 10^6$	$2.9 \cdot 10^{-6}$
pH 5.5	$9.4 \cdot 10^5$	$2.8 \cdot 10^{-4}$	$1.6 \cdot 10^6$	$4.7 \cdot 10^{-6}$
25pH 5.0	$9.5 \cdot 10^5$	$2.9 \cdot 10^{-4}$	$3.1 \cdot 10^5$	$9.1 \cdot 10^{-7}$
pH 4.5	$1.2 \cdot 10^6$	$3.5 \cdot 10^{-4}$	$1.2 \cdot 10^5$	$3.5 \cdot 10^{-7}$
30pH 4.0	$1.6 \cdot 10^6$	$4.8 \cdot 10^{-4}$	$7.2 \cdot 10^4$	$2.1 \cdot 10^{-7}$
pH 3.5	$9.5 \cdot 10^5$	$2.9 \cdot 10^{-4}$	$4.9 \cdot 10^4$	$1.4 \cdot 10^{-7}$
pH 3.0	$6.6 \cdot 10^5$	$2.0 \cdot 10^{-4}$	$2.9 \cdot 10^4$	$8.5 \cdot 10^{-8}$
35pH 2.5	$1.6 \cdot 10^5$	$4.8 \cdot 10^{-5}$	$1.4 \cdot 10^4$	$4.1 \cdot 10^{-8}$
pH 2.0	$3.0 \cdot 10^5$	$9.1 \cdot 10^{-5}$	$1.7 \cdot 10^4$	$5.0 \cdot 10^{-8}$
40SUM*	$6.4 \cdot 10^6$	$3 \cdot 10^{-3}$	$5.7 \cdot 10^6$	$2 \cdot 10^{-5}$

* SUM is the total pfu (or fraction of input) obtained from
all pH elution fractions

0959095.06301
T05250 5695860

TABLE 213

Fractionation of EpiC-10 and MA-ITI phage on Cat-G beads

10	Sample	EpiC-10		MA-ITI	
		Total pfu in sample	Fraction of input	Total pfu in sample	Fraction of input
	INPUT	$5.0 \cdot 10^{11}$	1.00	$4.6 \cdot 10^{11}$	1.00
15	Final TBS-TWEEN Wash	$1.8 \cdot 10^7$	$3.6 \cdot 10^{-5}$	$7.1 \cdot 10^6$	$1.5 \cdot 10^{-5}$
20	pH 7.0	$1.5 \cdot 10^7$	$3.0 \cdot 10^{-5}$	$6.1 \cdot 10^6$	$1.3 \cdot 10^{-5}$
	pH 6.0	$2.3 \cdot 10^7$	$4.6 \cdot 10^{-5}$	$2.3 \cdot 10^6$	$5.0 \cdot 10^{-6}$
	pH 5.5	$2.5 \cdot 10^7$	$5.0 \cdot 10^{-5}$	$1.2 \cdot 10^6$	$2.6 \cdot 10^{-6}$
25	pH 5.0	$2.1 \cdot 10^7$	$4.2 \cdot 10^{-5}$	$1.1 \cdot 10^6$	$2.4 \cdot 10^{-6}$
	pH 4.5	$1.1 \cdot 10^7$	$2.2 \cdot 10^{-5}$	$6.7 \cdot 10^5$	$1.5 \cdot 10^{-6}$
30	pH 4.0	$1.9 \cdot 10^6$	$3.8 \cdot 10^{-6}$	$4.4 \cdot 10^5$	$9.6 \cdot 10^{-7}$
	pH 3.5	$1.1 \cdot 10^6$	$2.2 \cdot 10^{-6}$	$4.4 \cdot 10^5$	$9.6 \cdot 10^{-7}$
	pH 3.0	$4.8 \cdot 10^5$	$9.6 \cdot 10^{-7}$	$3.6 \cdot 10^5$	$7.8 \cdot 10^{-7}$
35	pH 2.5	$2.0 \cdot 10^5$	$4.0 \cdot 10^{-7}$	$2.7 \cdot 10^5$	$5.9 \cdot 10^{-7}$
	pH 2.0	$2.4 \cdot 10^5$	$4.8 \cdot 10^{-7}$	$3.2 \cdot 10^5$	$7.0 \cdot 10^{-7}$
40	SUM*	$9.9 \cdot 10^7$	$2 \cdot 10^{-4}$	$1.4 \cdot 10^7$	$3 \cdot 10^{-5}$

* SUM is the total pfu (or fraction of input) obtained from all pH elution fractions

FOI 5095352

TABLE 214

Abbreviated fractionation of display phage on HNE beads

5	DISPLAY PHAGE				
	EpINE-7	MA-ITI 2	MA-ITI-E7 1	MA-ITI-E7 2	
10	INPUT (pfu)	1.00 (1.8·10 ⁹)	1.00 (1.2·10 ¹⁰)	1.00 (3.3·10 ⁹)	1.00 (1.1·10 ⁹)
15	WASH	6·10 ⁻⁵	1·10 ⁻⁵	2·10 ⁻⁵	2·10 ⁻⁵
	pH 7.0	3·10 ⁻⁴	1·10 ⁻⁵	2·10 ⁻⁵	4·10 ⁻⁵
	pH 3.5	3·10 ⁻³	3·10 ⁻⁶	8·10 ⁻⁵	8·10 ⁻⁵
20	pH 2.0	1·10 ⁻³	1·10 ⁻⁶	6·10 ⁻⁶	2·10 ⁻⁵
25	SUM*	4.3·10 ⁻³	1.4·10 ⁻⁵	1.1·10 ⁻⁴	1.4·10 ⁻⁴

* SUM is the total fraction of input pfu obtained from all pH elution fractions

F05250-3609550

TABLE 215

Fractionation of EpINE-7 and MA-ITI-E7 phage on HNE beads

10	Sample	EpINE-7		MA-ITI-E7	
		Total pfu in sample	Fraction of input	Total pfu in sample	Fraction of input
	INPUT	$1.8 \cdot 10^9$	1.00	$3.0 \cdot 10^9$	1.00
15	PH 7.0	$5.2 \cdot 10^5$	$2.9 \cdot 10^{-4}$	$6.4 \cdot 10^4$	$2.1 \cdot 10^{-5}$
	pH 6.0	$6.4 \cdot 10^5$	$3.6 \cdot 10^{-4}$	$4.5 \cdot 10^4$	$1.5 \cdot 10^{-5}$
20	pH 5.5	$7.8 \cdot 10^5$	$4.3 \cdot 10^{-4}$	$5.0 \cdot 10^4$	$1.7 \cdot 10^{-5}$
	pH 5.0	$8.4 \cdot 10^5$	$4.7 \cdot 10^{-4}$	$5.2 \cdot 10^4$	$1.7 \cdot 10^{-5}$
	pH 4.5	$1.1 \cdot 10^6$	$6.1 \cdot 10^{-4}$	$4.4 \cdot 10^4$	$1.5 \cdot 10^{-5}$
25	pH 4.0	$1.7 \cdot 10^6$	$9.4 \cdot 10^{-4}$	$2.6 \cdot 10^4$	$8.7 \cdot 10^{-6}$
	pH 3.5	$1.1 \cdot 10^6$	$6.1 \cdot 10^{-4}$	$1.3 \cdot 10^4$	$4.3 \cdot 10^{-6}$
30	pH 3.0	$3.8 \cdot 10^5$	$2.1 \cdot 10^{-4}$	$5.6 \cdot 10^3$	$1.9 \cdot 10^{-6}$
	pH 2.5	$2.8 \cdot 10^5$	$1.6 \cdot 10^{-4}$	$4.9 \cdot 10^3$	$1.6 \cdot 10^{-6}$
	pH 2.0	$2.9 \cdot 10^5$	$1.6 \cdot 10^{-4}$	$2.2 \cdot 10^3$	$7.3 \cdot 10^{-7}$
35	SUM*	$7.6 \cdot 10^6$	$4.1 \cdot 10^{-3}$	$3.1 \cdot 10^5$	$1.1 \cdot 10^{-4}$

40* SUM is the total pfu (or fraction of input) obtained from all pH elution fractions

T0520-5603660

CITATIONS

AKOH72:

Ako, H, RJ Foster, and CA Ryan,

- 5 "The preparation of anhydro-trypsin and its reactivity
with naturally occurring proteinase inhibitors",
Biochem Biophys Res Commun (USA) (1972), 47(6)1402-7.

ALBR83a:

- 10 Albrecht, G, K Hochstrasser, and OL Schonberger,
"Kunitz-type proteinase inhibitors derived by limited
proteolysis of the inter- α -trypsin inhibitor, IX:
isolation and characterization of the inhibitory parts of
inter- α -trypsin inhibitors from several mammalian sera",
15 Hoppe-Seyler's Z Physiol Chem (1983), 364:1697-1702.

ALBR83b:

- Albrecht, GJ, K Hochstrasser, and J-P Salier,
"Elastase inhibition by the inter- α -trypsin inhibitor and
20 derived inhibitors of man and cattle",
Hoppe-Seyler's Z Physiol chem (1983), 364:1703-1708.

ALMA83a:

- Almassy, RC, JC Fontecilla-Camps, FL Suddath, and CE Bugg,
25 "Structure of scorpion neurotoxin at 1.8 Å resolution",
Entry 1SN3 in Brookhaven Protein Data Bank, (1983).

ALMA83b:

- Almassy, RC, JC Fontecilla-Camps, FL Suddath, and CE Bugg,
30 "Structure of variant-3 scorpion neurotoxin from
Centruroides Sculpturatus ewing refined at 1.8 Å
resolution",
J Mol Biol (1983), 170:497ff.

35 ALMQ89:

- Almqvist, RG, SR Kadambi, DM Yasuda, FL Weitl, WE Polgar,
and LR Toll,
"Paralytic activity of (des-Glu1)conotoxin GI analogs in
the mouse diaphragm",
40 Int J Pept Protein Res, (Dec 1989), 34(6)455-62.

ANFI73:

- Anfinsen, CB,
"Principles that govern the folding of protein chains",
45 Science (1973), 181(96)223-30.

ARGO87:

- Argos, P,
"Analysis of Sequence-similar Pentapeptides in Unrelated
50 Protein Tertiary Structures",
J Mol Biol (1987), 197:331-348.

ARAK90:

Araki, K, M Kuwada, O Ito, J Kuroki, and S Tachibana,
 "Four disulfide bonds allocation of Na⁺, K⁺-ATPase
 inhibitor (SPAI)",

- 5 Biochem Biophys Res Comm (1990), 172(1)42-46.

ARMS81:

Armstrong, J, RN Perham, and JE Walker,
 "Domain structure of Bacteriophage fd Adsorption Protein",

- 10 FEBS Lett (1981), 135(1)167-172.

ARMS83:

Armstrong, J, JA Hewitt, and RN Perham,
 "Chemical modification of the coat protein in bacterio-
 15 phage fd and orientation of the virion during assembly and
 disassembly",
 EMBO J (1983), 2(10)1641-6.

ARNA90:

- 20 Arnaout, MA,
 "Leukocyte Adhesion Molecules Deficiency: Its STructural
 Basis, Pathophysiology and Implications for Modulating the
 Inflammatory Response",
 Immunological Reviews (1990), 114:_____.

25

AUER87:

Auerswald, E-A, W Schroeder, and M Kotick,
 "Synthesis, Cloning and Expression of Recombinant Aprotin-
 in",

- 30 Biol Chem Hoppe-Seyler (1987), 368:1413-1425.

AUER88:

Auerswald, E-A, D Hoerlein, G Reinhardt, W Schroder, and
 E Schnabel,
 35 "Expression, Isolation, and Characterization of Recom-
 binant [Arg¹⁵,Glu⁵²]Aprotinin",
 Bio Chem Hoppe-Seyler (1988), 369(Supplement):27-35.

AUER89:

- 40 Auerswald, E-A, W Bruns, D Hoerlein, G Reinhardt, E Schna-
 bel, and W Schroder,
 "Variants of bovine pancreatic trypsin inhibitor produced
 by recombinant DNA technology",
 UK Patent Application GB 2,208,511 A.

45

AUER90:

Auerswald, E-A, W Schroeder, E Schnabel, W Bruns, G Rein-
 hard, and M Kotick,
 "Homologs of Aprotinin produced from a recombinant host,
 50 process ecpression vector and recombinant host therefor
 and pharmaceutical use thereof",
 US Patent 4,894,436 (16 Jan 1990).

AUSU87:

Ausubel, FM, R Brent, RE Kingston, DD Moore, JG Seidman, JA Smith, and K Struhl, Editors

- 5 Current Protocols in Molecular Biology,
Greene Publishing Associates and Wiley-Interscience,
Publishers: John Wiley & Sons, New York, 1987.

BAKE87:

- 10 Baker, K, N Mackman, and IB Holland,
"Genetics and Biochemistry of the Assembly of Proteins
into the Outer Membrane of E. coli",
Prog Biophys molec Biol (1987), 49:89-115.

BALD85:

- 15 Balduyck, M, M Davril, C Mizon, M Smyrlaki, A Hayem, and
J Mizon,
"Human urinary proteinase inhibitor: inhibitory properties
and interaction with bovine trypsin",
20 Biol Chem Hoppe-Seyler (1985), 366:9-14.

BANN81:

- Banner, DW, C Nave, and DA Marvin,
"Structure of the protein and DNA in fd filamentous
25 bacterial virus",
Nature (1981), 289:814-816.

BARB85:

- Barbe, J, JA Vericat, M Llagostera, and R Guerrero,
30 "Expression of the SOS genes of Escherichia coli in
Salmonella typhimurium",
Microbiologia (1985), 1(1-2)77-87.

BECK80:

- 35 Beck, E,
"Nucleotide sequence of the gene ompA coding the outer
membrane protein II* of Escherichia coli K-12",
Nucl Acid Res (1980), 8(13)3011-3024.

BECK83:

- 40 Beckwith, J, and TJ Silhavy,
"Genetic Analysis of Protein Export in Escherichia coli",
Methods in Enzymology (1983), 97:3-11.

BECK88b:

- 45 Beckmann, J, A Mehlich, W Schroeder, HR Wenzel, and
H Tschesche,
"Preparation of chemically 'mutated' aprotinin homologues
by semisynthesis: P1 substitutions change inhibitory
50 specificity",
Eur J Biochem (1988), 176:675-82.

005290-56096860

BECK89a:

- Beckmann, J, A Mehlich, W Schroeder, HR Wenzel, and
H Tschesche,
"Semisynthesis of Arg¹⁵, Glu¹⁵, Met¹⁵, and Nle¹⁵-Aprotinin
5 Involving Enzymatic Peptide Bond Resynthesis",
J Protein Chem (1989), 8(1)101-113.

BECK89b:

- Becker, S, E Atherton, H Michel, and RD Gordon,
10 "Synthesis and characterization of conotoxin IIIa",
J Protein Chem, (Jun 1989), 8(3)393-4.

BECK89c:

- Becker, S, E Atherton, and RD Gordon,
15 "Synthesis and characterization of mu-conotoxin IIIa",
Eur J Biochem, (Oct 20 1989), 185(1)79-84.

BENS84:

- Benson, SA, E Bremer, and TJ Silhavy,
20 "Intragenic regions required for LamB export",
Proc Natl Acad Sci USA (1984), 81:3830-34.

BENS87b:

- Benson, SA, and E Bremer,
25 "In vivo selection and characterization of internal
deletions in the lamB::lacZ gene fusion",
Gene (1987), 52(2-3)165-73.

BENS87c:

- Benson, SA, MN Hall, and BA Rasmussen,
30 "Signal Sequence Mutations That Alter Coupling of
Secretion and Translation of an Escherichia coli Outer
Membrane Protein",
J Bacteriol (1987), 169(10)4686-91.

35

BENS88:

- Benson, SA, JL Occi, BA Sampson,
"Mutations that alter the pore function of the OmpF porin
of Escherichia coli K12",
40 J Mol Biol (1988) 203(4)961-70.

BENZ88a:

- Benz, R, and K Bauer,
"Permeation of hydrophilic molecules through the outer
45 membrane of gram-negative bacteria",
Eur J Biochem (1988), 176:1-19.

BENZ88b:

- Benz, R,
50 "Structure and Function of Porins from Gram-Negative
Bacteria",
Ann Rev Microbiol (1988), 42:359-93.

T05230-56936360

BERG88:

Berg, JM,

- 5 "Proposed structure for the zinc-binding domains from
transcription factor IIIA and related proteins",
Proc Natl Acad Sci USA (1988), 85:99-102.

BETT88:

Better, M, CP Chang, RR Robinson, and AH Horwitz,

- 10 "Escherichai coli Secretion of an Active Chimeric
Antibody Fragment",
Science (1988), 240:1041-1043.

BHAT86:

Bhatnagar, PK, and JC Frantz,

- 15 "Synthesis and Antigenic activity of E. coli ST and its
analogues",
Develop biol Standard (1986), 63:79-87.

BIRD67:

Birdsell, DC, and EH Cota-Robles,

- 20 "Production and Ultrastructure of lysozyme and
ethylenediaminetetraacetate-lysozyme spheroplasts of E.
coli",
25 J Bacteriol (1967), 93:427-437.

BIET86:

Bieth, JG,

- 30 "Elastase: Catalytic and Biological Properties",
pp. 217-320 in Regulation of Matrix Accumulation, Editor:
RP Mecham, Academic Press, Orlando, 1986.

BLOW72:

Blow et al.,

- 35 J Mol Biol (1972), 69:137ff.

BODE89:

Bode, W, HJ Greyling, R Huber, J Otlewski, and T Wilusz,
"The refined 2.0 A X-ray crystal structure of the complex
40 formed between bovine beta-trypsin and CMTI-I, a trypsin
inhibitor from squash seeds (Cucurbita maxima). Topolog-
ical similarity of the squash seed inhibitors with the
carboxypeptidase A inhibitor from potatoes",
FEBS Lett (Jan 2 1989), 242(2)285-92.

45

BOEK80:

Boeke, JD, M Russel, and P Model,

- "Processing of Filamentous Phage Pre-coat Protein: Effect
of Sequence Variations near the Signal Peptidase Cleavage
50 Site",
J Mol Biol (1980), 144:103-116.

Boeke, JD, P Model, and ND Zinder,
"Effects fo Bacteriophage fl Gene III Protein on the Host
Cell Membrane",

- BOQU87:
Boquet, PL, C Manoil, and J Beckwith,
"Use of TnphoA to Detect Genes for Exported Proteins in
10 Escherichia coli: Identification of the Plasmid-Encoded
Gene for a Periplasmic Acid Phosphatase",
J Bacteriol (1987), 169:1663-1669.

15 Botstein, D, and D Shortle,
"Strategies and applications of in vitro mutagenesis",
Science, (1985), 229(4719)1193-201.

20 Bouges-Bocquet, B, H Villarroja, and M Hofnung,
"Linker Mutagenesis in the Gene of an Outer Membrane
Protein of Escherichia coli, LamB",
J Cellular Biochem (1984), 24:217-28.

Boulain, JC, A Charbita and M Hofnung,
"Mutagenesis by random linker insertion into the lamB gene
of Escherichia coli K12",
Mol Gen Genet, (1986), 205(2)339-48.

Brawerman, G,
"Determinants of messenger RNA stability",
Cell (1987), 48(1)5-6.

Calamia, J, and C Manoil,
"lac permease of Escherichia coli: topology and sequence
elements promoting membrane insertion",
40 Proc Natl Acad Sci USA, (Jul 1990), 87(13)4937-41.

Campanelli, D, M Melchior, Yiping Fu, M Nakata, H Shuman,
C Nathan, and JE Gabay,
45 "Cloning of cDNA for Proteinase 3: A Serine Protease,
Antibiotic, and Autoantigen from Human Neutrophils",
J Exp Med (Dec 1990), 172:1709-15.

CARM90:

- Carmel, G, D Hellstern, D Henning, and JW Coulton,
 "Insertion mutagenesis of the gene encoding the ferri-
 chrome-iron receptor of Escherichia coli K-12",
 5 J Bacteriol, (Apr 1990), 172(4)1861-9.

CARU85:

- Caruthers, MH,
 "Gene Synthesis Machines: DNA Chemistry and Its Uses",
 10 Science (1985), 230:281-285.

CARU87:

- Caruthers, MH, P Gottlieb, LP Bracco, and L Cummings,
 "The Thymine 5-Methyl Group: A Protein-DNA Contact Site
 15 Useful for Redesigning Cro Repressor to Recognize a New
 Operator",
 in Protein Structure, Folding, and Design 2, 1987,
 Ed. D Oxender (New York, AR Liss Inc) p.9ff.

CAST79:

- Castillo, MJ, K Nakajima, M Zimmerman, and JC Powers,
 "Sensitive substrates for human leukocyte and porcine
 pancreatic elastase: a study of the merits of various
 chromophoric and fluorogenic leaving groups in assays for
 25 serine proteases",
 Anal Biochem (1979), 99(1)53-64.

CATR87:

- Catron, KM, and CA Schnaitman,
 30 "Export of Protein in Escherichia coli: a Novel Mutation
 in ompC Affects Expression of Other Major Outer Membrane
 Proteins",
 J Bacteriol (1987), 169:4327-34.

CHAM82:

- Chambers, RW, I Kucan, and Z Kucan,
 "Isolation and characterization of phi-X174 mutants
 carrying lethal missense mutations in gene G",
 Nucleic Acids Res (1982), 10(20)6465-73.

40

CHAN79:

- Chang, CN, P Model, and G Blobel,
 "Membrane biogenesis: Cotranslational integration of the
 bacteriophage f1 coat protein into an Escherichia coli
 45 membrane fraction",
 Proc Natl Acad Sci USA (1979), 76:1251-1255.

CHAP90:

Chapot, MP, Y Eshdat, S Marullo, JG Guillet, A Charbit, AD Strosberg, and C Delavier-Klutchko,
 "Localization and characterization of three different
 5 beta-adrenergic receptors expressed in *Escherichia coli*",
 Eur J Biochem (1990), 187(1)137-44.

CHAR84:

Charbit, A, J-M Clement, and M Hofnung,
 10 "Further Sequence Analysis of the Phage Lambda Receptor
 Site",
 J Mol Biol (1984), 175:395-401.

CHAR86a:

15 Charbit, A, JC Boulain, A Ryter, and M Hofnung,
 "Probing the topology of a bacterial membrane protein by
 genetic insertion of a foreign epitope; expression at the
 cell surface",
 EMBO J, (1986), 5(11)3029-37.

20

CHAR86b:

Charbit, A, J-C Boulain, and M Hofnung,
 "Une methode genetique pour exposer un epitope choisi a la
 surface de la bacteria *Escherichia coli*. Perspectives [A
 25 genetic method to expose a chosen epitope on the surface
 of the bacteria *E. coli*]",
 Comptes Rendu Acad Sci, Paris, (1986), 302:617-24.

CHAR87:

30 Charbit, A, E Sobczak, ML Michel, A Molla, P Tiollais, and
 M Hofnung,
 "Presentation of two epitopes of the preS2 region of
 hepatitis B virus on live recombinant bacteria",
 J Immunol (1987), 139:1658-64.

35

CHAR88a:

Charbit, A, K Gehring, H Nikaido, T Ferenci, and M
 Hofnung,
 "Maltose transport and starch binding in phage-resistant
 40 point mutants of maltoporin. Functional and topological
 implications",
 J Mol Biol (1988), 201(3)487-96.

CHAR88b:

45 Charbit, A, A Molla, W Saurin, and M Hofnung,
 "Versatility of a vector for expressing foreign
 polypeptides at the surface of gram-negative bacteria",
 Gene (1988), 70(1)181-9.

FOI 290-560960

CHAR88c:

Charbit, A, S Van der Werf, V Mimic, JC Boulain, M Girard,
and M Hofnung,

- 5 "Expression of a poliovirus neutralization epitope at the
surface of recombinant bacteria: first immunization
results",
Ann Inst Pasteur Microbiol (1988), 139(1)45-58.

CHAR90:

- 10 Charbit, A, A Molla, J Ronco, JM Clement, V Favier,
EM Bahraoui, L Montagnier, A Leguern, and M Hofnung,
"Immunogenicity and antigenicity of conserved peptides
from the envelope of HIV-1 expressed at the surface of
recombinant bacteria",
15 AIDS (1990), 4(6)545-51.

CHAV88:

Chavrier, P, P Lemaire, O Revelant, R Bravo, and
P Charnay,

- 20 "Characterization of a Mouse Multigene Family That Encodes
Zinc Finger Structures",
Molec Cell Biol (1988), 8(3)1319-26.

CHAZ85:

- 25 Chazin, WJ, DP Goldenberg, TE Creighton, and K Wuthrich,
"Comparative studies of conformation and internal mobility
in native and circular basic pancreatic trypsin inhibitor
by ¹H nuclear magnetic resonance in solution",
Eur J Biochem (1985), 152: (2)429-37.

CHOT75:

Chothia, C, and J Janin,
"Principles of protein-protein recognition",
Nature (1975), 256:705-708.

35

CHOT76:

Chothia, C, S Wodak, and J Janin,
"Role of subunit interfaces in the allosteric mechanism of
hemoglobin",

- 40 Proc Natl Acad Sci USA (1976), 73:3793-7.

CHOU74:

Chou, PY, and GD Fasman,
"Prediction of protein conformation"

- 45 Biochemistry (1974), 13: (2)222-45.

CHOU78a:

Chou, PY, and GD Fasman,
"Prediction of the secondary structure of proteins from
their amino acid sequence",
Adv Enzymol (1978), 47:45-148.

50

FOI 290 56095850

CHOU78b:

Chou, PY, and GD Fasman,
 "Empirical predictions of protein conformation"
 Annu Rev Biochem (1978), 47:251-76.

5

CHOW87:

Chowdhury, K, U Deutsch, and P Gruss,
 "A Multigene Family Encoding Several 'Finger' Structures
 Is Present and Differentially Active in Mammalian
 Genomes",
 Cell (1987), 48:771-778.

10

CLEM81:

Clement, JM, and M Hofnung,
 "The sequence of the lambda receptor, an outer membrane
 protein of E. coli K12",
 Cell (1981), 27:507-514.

15

CLEM83:

Clement JM, E Lepouce, C Marchal, and M Hofnung,
 "Genetic Study of a membrane protein: DNA sequence
 alterations due to 17 LamB point mutations affecting
 adsorption of phage lambda",
 EMBO J (1983), 2:77-80.

20

CLIC88:

Click, EM, GA McDonald, and CA Schnaitman,
 "Translational Control of Exported Proteins
 That Results from OmpC Porin Overexpression",
 J Bacteriol (1988), 170:2005-2011.

25

CLOR86:

Clore, GM, AT Brunger, M Karplus, AM Gronenborn,
 "Application of Molecular Dynamics with Interproton
 Distance Restraints to Three-dimensional Protein Structure
 Determination: A model study of Crambin",
 J Mol Biol (1986), 191:523-551.

35

CLOR87a:

Clore, GM, AM Gronenborn, M Kjaer, and FM Poulsen,
 "The determination of the three-dimensional structure of
 barley serine proteinase inhibitor 2 by nuclear magnetic
 resonance distance geometry and restrained molecular
 dynamics",

40

Protein Engineering (1987), 1(4)305-311.

CLOR87b:

Clore, GM, AM Gronenborn, MNG James, M Kjaer, CA McPhalen,
 and FM Poulsen,
 "Comparison of the solution and X-ray structures of barley
 serine proteinase inhibitor 2",
 Protein Engineering (1987), 1(4)313-318.

50

T05290"56096850

CLUN84:

- Clune, A, K-S Lee, and T Ferenci,
 "Affinity Engineering of Maltoporin: Variants with
 5 Enhanced Affinity for Particular Ligands",
 Biochem and Biophys Res Comm (1984), 121:34-40.

CREI74:

- Creighton, TE,
 10 "Intermediates in the Refolding of Reduced Pancreatic
 Trypsin Inhibitor",
 J Mol Biol (1974), 87:579-602.

CREI77a:

- 15 Creighton, TE,
 "Conformational Restrictions on the Pathway of Folding and
 Unfolding of the Pancreatic Trypsin Inhibitor",
 J Mol Biol (1977), 113:275-293.

20 CREI77b:

Creighton, TE,
 Energetics of Folding and Unfolding of Pancreatic Trypsin
 Inhibitor",
 J Mol Biol (1977), 113:295-312.

25

CREI80:

- Creighton, TE,
 "Role of the Environment in the Refolding of Reduced
 Pancreatic Trypsin Inhibitor",
 30 J Mol Biol (1980), 144:521-550.

CREI84:

- Creighton, TE,
Proteins: Structures and Molecular Principles,
 35 W H Freeman & Co, New York, 1984.

CREI87:

- Creighton, TE, and IG Charles,
 "Biosynthesis, Processing, and Evolution of Bovine
 40 Pancreatic Trypsin Inhibitor",
 Cold Spring Harb Symp Quant Biol (1987), 52:511-519.

CREI88:

- Creighton, TE,
 45 "Disulphide Bonds and Protein Stability",
 BioEssays (1988), 8(2)57-63.

CRIS84:

- Crissman, JW, and GP Smith,
 50 "Gene-III Protein of Filamentous Phages: Evidence for a
 Carboxyl-Terminal Domain with a Role in Morphogenesis",
 Virology (1984), 132:445-55.

T05330-56096860

Cruz, LJ, WR Gray, BM Olivera, RD Zeikus, L Kerr,
D Yoshikami, and E Moczydlowski,

- CRUZ89:
10 Cruz, LJ, G Kupryszewski, GW LeCheminant, WR Grey,
BM Oliveria, and J Rivier,
"mu-Conotoxin GIIIA, a Peptide Ligand for Muscle Sodium
Channels: Chemical Synthesis, Radiolabeling, and Receptor
Characterization",
15 Biochem (1989), 28:3437-3442.

Cwirala, SE, EA Peters, RW Barrett, and WJ Dower,
"Peptides on Phage: A vast library of peptides for
identifying ligands",
Proc Natl Acad Sci USA, (August 1990), 87:6378-6382.

25 Dailey, D, GL Schieven, MY Lim, H Marquardt, T Gilmore,
J Thorner, and GS Martin,
"Novel yeast protein kinase (YPK1 gene product) is a 40-
kilodalton phosphotyrosyl protein associated with protein-
tyrosine kinase activity",
Mol Cell Biol (Dec 1990), 10(12)6244-56.

Dallas, WS,
"The Heat-Stable Toxin I Gene from Escherichia coli 18D",
J Bacteriol (1990); 172(9)5490-93.

Dargent, B, A Charbit, M Hofnung, and F Pattus,
"Effect of point mutations on the in-vitro pore properties
of maltoporin, a protein of Escherichia coli outer
40 membrane",
J Mol Biol (1988), 201(3)497-506.

45 Dawkins, R,
The Blind Watchmaker,
W W Norton & Co, New York, 1986.

50 Day, LA, CJ Marzec, SA Reisberg, and A Casadevall,
"DNA Packing in Filamentous Bacteriophage",
Ann Rev Biophys Biophys Chem (1988), 17:509-39.

Biochem Biophys Res Commun (1978), 80(4)993-9.

- DEVR84:
de Vries, G, CK raymond, and RA Ludwig,
"Extension of bacteriophage λ host range: Selection,
cloning, and characterization of a constitutive λ receptor
5 gene",
Proc Natl Acad Sci USA (1984), 81:6080-4.
- DIAR90:
Diarra-Mehrpour, M, J Bourguignon, R Sesboue, J-P Salier,
10 T Leveillard and J-P Martin,
"Structural analysis of the human inter- α -trypsin
inhibitor light-chain gene",
Eur J Biochem (1990), 191:131-139.
- 15 DICK83:
Dickerson, RE, and I Geis, ..
Hemoglobin: Structure, Function, Evolution, and
Pathology,
The Benjamin/Cummings Publishing Co, Menlo Park, CA, 1983.
20
- DILL87:
Dill, KA,
"Protein Surgery",
Protein Engineering (1987), 1:369-371.
- 25 DOUG84:
Dogan, G, and P Morrissey,
"Molecular analysis of the virulence determinants of
enterotoxigenic Escherichia coli isolated from domestic
30 animals: applications for vaccine development",
Vet Microbiol (1984/5), 10:241-57.
- DON087
Donovan, W, Z Liangbiao, K Sandman, and R Losick,
35 "Genes Encoding Spore Coat Polypeptides from Bacillus
subtilis",
J Mol Biol (1987), 196:1-10.
- DUCH88:
40 Duchene, M, A Schweized, F Lottspeich, G Krauss, M Marget,
K Vogel, B-U von Specht, and H Domdey,
"Sequence and Transcriptional Start Site of the
Pseudomonas aeruginosa Outer Membrane Porin Protein F
Gene",
45 J Bacteriol (1987), 170:155-162.
- DUFT85:
Dufton, MJ,
"Proteinase inhibitors and dendrotoxins",
50 Eur J Biochem (1985), 153:647-654.

- DULB86:
Dulbecco, R,
"Viruses with Recombinant Surface Proteins",
US Patent 4,593,002, June 3, 1986.
- 5
DUPL88:
Duplay, P, and M Hofnung,
"Two Regions of Mature Periplasmic Maltose-Binding Protein
of Escherichia coli Involved in Secretion",
10 J Bacteriol (1988), 170(10)4445-50.
- DWAR89:
Dwarakanath, P, SS Viswiswariah, YVBK Subrahmanyam,
G Shanthi, HM Jagannatha, and TS Balganes,
15 "Cloning and hyperexpression of a gene encoding the heat-
stable toxin of Escherichia coli",
Gene (1989), 81:219-226.
- EHRM90:
20 Ehrmann, M, D Boyd, and J Beckwith,
"Genetic analysis of membrane protein topology by a
sandwich gene fusion approach",
Proc Natl Acad Sci USA, (Oct 1990), 87(19)7574-8.
- 25 EIGE90:
Eigenbrot, C, M Randal, and AA Kossiakoff,
"Structural effects induced by removal of a disulfide-
bridge: the X-ray structure of the C30A/C51A mutant of
basic pancreatic trypsin inhibitor at 1.6 Å",
30 Protein Engineering (1990), 3(7)591-598.
- EISE85:
Eisenbeis, SJ, MS Nasoff, SA Noble, LP Bracco, DR Dodds,
MH Caruthers,
35 "Altered Cro Repressors from engineered mutagenesis of a
synthetic cro gene",
Proc Natl Acad Sci USA (1985), 82:1084-1088.
- ELLE88:
40 Elleman, TC,
"Pilins of Bacteroides nodosus: molecular basis of
serotypic variation and relationships to other bacterial
pilins",
Microbiol Rev (1988), 52(2)233-47.
- 45
EMPI82:
Empie, MW, and M Laskowski, Jr,
"Thermodynamics and Kinetics fo Single Residue
Replacements in Avian Ovomucoid Third Domains: Effect on
50 Inhibitor Interactions with Serine Proteinases",
Biochemistry (1982), 21:2274-84.

FERE83:

Ferenci, T, and KS Lee,
 "Isolation by affinity chromatography, of mutant
Escherichia coli cells with novel regulation of lamB
 5 expression",
 J Bacteriol (1983), 154:984-987.

FERE84:

Ferenci, T,
 10 "Genetic manipulation of bacterial surfaces through
 affinity-chromatographic selection",
 Trends in Biological Science (1984) Vol. ?:44-48.

FERE86a:

15 Ferenci, T, and K-S Lee,
 "Temperature-Sensitive Binding of α -Glucans by Bacillus
stearothermophilus",
 J Bacteriol (1986), 166:95-99.

FERE86b:

Ferenci, T, M Muir, K-S Lee, and D Maris,
 "Substrate specificity of the Escherichia coli
 maltodextrin transport system and its component
 proteins.",
 25 Biochimica et Biophysica Acta (1986), 860:44-50.

FERE89a:

Ferenci, T, and KS Lee,
 "Channel architecture in maltoporin: dominance studies
 30 with lamB mutations influencing maltodextrin binding
 provide evidence for independent selectivity filters in
 each subunit",
 J Bacteriol (1989) 171(2)855-61.

FERE89b:

Ferenci, T, and S Stretton,
 "Cysteine-22 and cysteine-38 are not essential for the
 function of maltoporin (LamB protein)",
 35 FEMS Microbiol Lett (1989), 52(3)335-9.

FERR90:

Ferrer-Lopez, P, P Renesto, M Schattner, S Bassot, P
 Laurent, and M Chignard,
 "Activation of human platelets by C5a-stimulated
 45 neutrophils: a role for cathepsin G",
 American J Physiology (1990) 258:C1100-C1107.

T062330-5696360

FIOR85:

- Fioretti, E, G Iacopino, M Angeletti, D Barra, F Bossa,
and F Ascoli,
"Primary Structure and Antiproteolytic Activity of a
5 Kunitz-type Inhibitor from Bovine Spleen",
J Biol Chem (1985), 260:11451-11455.

FIOR88:

- Fioretti, E, M Angeletti, L Fiorucci, D Barra, F Bossa,
10 and F Ascoli,
"Aprotinin-Like Isoinhibitors in Bovine Organs",
Biol Chem Hoppe-Seyler (1988), 369(Suppl)37-42.

FRAN87:

- 15 Frankel, AD, JM Berg, and CO Pabo,
"Metal-dependent folding of a single zinc finger from
transcription factor IIIA",
Proc Natl Acad Sci USA (1987), 84:4841-45.

20 FRAN88:

Frankel, A, and CO Pabo,
"Fingering Too Many Proteins",
Cell (1988), 53:675.

25 FRAN89:

- Franconi, GM, PD Graf, SC Lazarus, JA Nadel, GH Caughey,
"Mast Cell Tryptase and Chymase Reverse Airway Smooth
Muscle Relaxation Induced by Vasoactive Intestinal Peptide
in the Ferret",
30 J Pharmacol and Exp Therap (1989), 248(3)947-51.

FREI90:

- Freimuth, PI, JW Taylor, and ET Kaiser,
"Introduction of Guest Peptides into Escherichia coli
35 Alkaline Phosphatase",
J Biol Chemistry, (15 January 1990), 265(2)896-901.

FREU89:

- Freudl, R, H Schwarz, M Degen, and U Henning,
40 "A lower size limit exists for export of fragments of an
outer membrane protein (OmpA) of Escherichia coli K-12",
J Mol Biol (1989), 205(4)771-5.

FRIT85:

- 45 Fritz, H-J,
"The Oligonucleotide-directed Construction of Mutations in
Recombinant Filamentous Phage",
DNA Cloning, Editor: DM Glover, IRL Press, Oxford, UK,
1985.

50

F05390"56055350

Gariepy, J, P O'Hanley, SA Waldman, F Murad, and
GK Schoolnik,

5

Gariepy, J, A Lane, F Frayman, D Wilbur, W Robien,
G Schoolnik, and O Jardetzky,

10

Gariepy, J, AK Judd, and GK Schoolnik,
"Importance of disulfide bridges in the structure and
activity of Escherichia coli enterotoxin ST1b",
Proc Natl Acad Sci USA (1987), 84:8907-11.

GAUS87:

Gauss, P, KB Krassa, DS McPheeters, MA Nelson, and L Gold,
"Zinc(II) and the single-stranded DNA binding protein of
bacteriophage T4",

25

Gebhard, W, and K Hochstrasser,
"Inter- α -trypsin inhibitor and its close relatives",
in Barret and Salvesen (eds.) Protease Inhibitors (1986)
Elsevier Science Publishers BV (Biomedical Division)
pp.389-401.

GEBH90:

35 Gebhard, W, K Hochstrasser, H Fritz, JJ Enghild, SV Pizzo,
and G Salvesen.

Structure of the inter- α -inhibitor (inter- α -trypsin inhibitor) and pre- α -inhibitor: current state and proposition of a new terminology",

40

Gehring, K, A Charbit, E Brissaud, and M Hofnung,
"Bacteriophage lambda receptor site on the Escherichia
coli K-12 LamB protein",
J Bacteriol (1987), 169(5)2103-6.

45

- GERD84:
Gerday, C, M Herman, J Olivy, N Gerardin-Otthiers, D Art,
E Jacquemin, A Kaeckenbeeck, and J van Beeumen,
"Isolation and characterization of the Heat Stable
5 enterotoxin for a pathogenic bovine strain of Escherichia
coli",
Vet Microbiol (1984), 2:399-414.
- GETZ88:
10 Getzoff, ED, HE Parge, DE McRee, and JA Tainer,
"Understanding the Structure and Antigenicity of
Gonococcal Pili",
Rev Infect Dis (1988), 10(Suppl 2)S296-299.
- 15 GIBS88:
Gibson, TJ, JPM Postma, RS Brown, and P Argos,
"A model for the tertiary structure of the 28 residue DNA-
binding motif ('Zinc finger') common to many eukaryotic
transcriptional regulatory proteins",
20 Protein Engineering (1988), 2(3)209-218.
- GIRA89:
Girard, TJ, LA Warren, WF Novotny, KM Likert, SG Brown, JP
Miletich, and GJ Broze Jr,
25 "Functional significance of the Kunitz-type inhibitory
domains of lipoprotein-associated coagulation inhibitor",
Nature (1989), 338:518-20.
- GOLD83:
30 Goldenberg, DP, and TE Creighton,
"Circular and circularly permuted forms of bovine
pancreatic trypsin inhibitor.",
J Mol Biol (1983), 165(2)407-13.
- 35 GOLD84:
Goldenberg, DP, and TE Creighton,
"Folding Pathway of a circular Form of Bovine Pancreatic
Trypsin Inhibitor",
J Mol Biol (1984), 179:527-45.
- 40 GOLD85:
Goldenberg, DP,
"Dissecting the Roles of Individual Interactions in
Protein Stability: Lessons From a Circularized Protein",
45 J Cellular Biochem (1985), 29:321-335.

T06290-3696850

Gold, L, and G Stormo, .
"Translation Initiation",
Volume 2, Chapter 78, p 1302-1307,

- 5 Escherichia coli and Salmonella typhimurium: Cellular and
Molecular Biology,
Neidhardt, FC, Editor-in-Chief,
Amer Soc for Microbiology, Washington, DC, 1987.
- 10 GOLD88:
Goldenberg, DP,
"Kinetic Analysis of the Folding and Unfolding of a Mutant
Form of Bovine Pancreatic Trypsin Inhibitor Lacking the
Cysteine-14 and -38 Thiols",
15 Biochem (1988), 27:2481-89.
- GOTT87:
Gottesman, S,
"Regulation by Proteolysis",
20 Volume 2, chapter 79, p 1308-1312.
Escherichia coli and Salmonella typhimurium: Cellular and
Molecular Biology,
Neidhardt, FC, Editor-in-Chief,
Amer Soc for Microbiology, Washington, DC, 1987.
- 25 GRAY81a:
Gray, WR, A Luque, BM Olivera, J Barrett, and LJ Cruz,
"Peptide Toxins from Conus geographicus Venom",
J Biol Chem (1981), 256:4734-40.
- 30 GRAY81b:
Gray, CW, RS Brown, and DA Marvin,
"Adsorption Complex of Filamentous Virus",
J Mol Biol (1981), 146:621-627.
- 35 GRAY83:
Gray, WR, JE Rivier, R Galyean, LJ Cruz, and BM Olivera,
"Conotoxin MI. Disulfide bonding and conformational
states",
40 J Biol Chem, (1983), 258(20)12247-51.
- GRAY84:
Gray, WR, FA Luque, R Galyean, E Atherton, and
RC Sheppard, BL Stone, A Reyes, J Alford, M McIntosh, BM
45 Olivera et al.
"Conotoxin GI: disulfide bridges, synthesis, and
preparation of iodinated derivatives",
Biochemistry, (1984), 23(12)2796-802.

5

10

15

20

25

30

35

40

45

5 neuromuscular junction",
Eur J Pharmacol (1985), 118(3)351-4.

10 "Synthesis of mu-conotoxin GIIIA: a chemical probe for sodium channels",
Chem Pharm Bull (Tokyo), (Jan 1990), 38:236-8.

15 Hecht, MH, JS Richardson, DC Richardson, and RC Ogden,
 "De Novo Design, Expression, and Characterization of
 Felix: A Four-Helix Bundle Protein of Native-Like
 Sequence",
 Science, (24 Aug 1990), 249:884-91.

25 Gene, (Dec 21 1989), 85(1)115-24.

Heijne, G von, and C Manoil,
"Review: Membrane proteins: from sequence to structure",
Protein Engineering (1990), 4(2)109-112.

Heine, HG, J Kyngdon, and T Ferenci,
"Sequence determinants in the lamB gene of Escherichia
coli influencing the binding and pore selectivity of
maltoporin.",
Gene (1987), 53:287-92.

40 Heine, HG, G Francis, KS Lee, and T Ferenci,
"Genetic analysis of sequences in maltoporin that
contribute to binding domains and pore structure",
J Bacteriol (April 1988), 170:1730-8.

50 Biochem (1989), 28:2392-98.

HENR87:

Henriksen, AZ, and JA Maeland,
 "The Porin Protein of the Outer Membrane of Escherichia coli: Reactivity in Immunoblotting, Antibody-binding by
 5 the Native Protein, and Cross-Reactivity with other
 Enteric Bacteria",
 Acta path microbiol immunol scand, Sect B (1987), 95:315-
 321.

10 HIDA90:

Hidaka, Y, K Sato, H Nakamura, J Kobayashi, Y Ohizumi, and
 Y SHimonishi,
 "Disulfide Pairings in geographutoxin I, a peptide
 neurotoxin from Conus geographus",

15 FEBS Lett (1990), 264(1)29-32.

HILL89:

Hillyard, DR, BM Olivera, S Woodward, GP Corpuz, WR Gray,
 CA Ramilo, LJ Cruz,
 20 "A Molluscivorus Conus Toxin: Conserved Framework in
 Conotoxins",
 Biochem (1989), 28:358-61.

HINE80:

25 Hines, JC, and DS Ray,
 "Construction and characterization of new coliphage M13
 cloning vectors.",
 Gene (1980), 11:(3-4)207-18.

30 HOCH84:

Hoschstrasser, K, and E Wachter,
 "Elastase inhibitors, a process for their preparation and
 medicaments containing these inhibitors",
 US Patent 4,485,100 (27 Nov 1984).

35

HOCJ85:

Ho, C, M Jasin, and P Schimmel,
 "Amino acid replacements that compensate for a large
 polypeptide deletion in an enzyme",
 40 Science (1985), 229:389-93.

HOJI82:

Hojima, Y, JV Pierce, and JJ Pisano,
 "Pumpkin Seed Inhibitor of Human Factor XII_a (activated
 45 Hageman Factor) and Bovine Trypsin",
 Biochem (1982), 21:3741-46.

T06290"56036360

HOLA89a:

Holak, TA, D Gondol, J Otlewski, and T Wilusz,
 "Determination of the Complete Three-Dimensional Structure
 of the Trypsin Inhibitor from Squash Seeds in Aqueous
 5 Solution by Nuclear Magnetic Resonance and a Combination
 of Distance Geometry and Dynamic Simulated Annealing",
 J Mol Biol (1989), 210:635-648.

HOLA89b:

10 Holak, TA, W Bode, R Huber, J Otlewski, and T Wilusz,
 "Nuclear magnetic resonance solution and X-ray structures
 of squash trypsin inhibitor exhibit the same conformation
 of the proteinase binding loop",
 J Mol Biol (Dec 5 1989), 210(3)649-54.

15

HORV89:

Horvat, S, B Grgas, N Raos, and VI Simeon,
 "Synthesis and acid ionization constants of cyclic cystine
 peptides H-Cys-(Gly)_n-Cys-OH (n=0-4)",

20

Int J Peptide Protein Res (1989), 34:346-51.

HOOP87:

Hoopes, BC, and WR McClure,
 25 "Strategies in Regulation of Transcription Initiation",
 Volume 2, Chapter 75, p 1231-1240,
Escherichia coli and Salmonella typhimurium: Cellular and
Molecular Biology,
 Neidhardt, FC, Editor-in-Chief,
 30 Amer Soc for Microbiology, Washington, DC, 1987.

HOUG84:

Houghten, RA, JM Ostresh, and FA Klipstein,
 "Chemical synthesis of an octadecapeptide with the
 35 biological and immunological properties of human heat-
 stable Escherichia coli enterotoxin",
 Eur J Biochem (1984), 145:157-162.

HUBB86:

40 Hubbard, RC, and RG Crystal,
 "Antiproteases and Antioxidants: Strategies for the
 Pharmacologic Prevention of Lung Destruction",
 Respiration (1986), 50(Suppl 1)56-73.

45

HUBB89:
 Hubbard, RC, MA Casolaro, M Mitchell, SE Sellers,
 F Arabia, MA Matthay, and RG Crystal,
 "Fate of aerosolized recombinant DNA-produced α -1-
 antitrypsin: Use of the epithelial surface of the lower
 50 respiratory tract to administer proteins of therapeutic
 importance",
 Proc Natl Acad Sci USA (1989), 86:680-4.

FOI 290 56096860

- HUBE74:
 Huber, R, D Kukla, W Bode, P Schwager, K Bartels,
 J Deisenhofer, and W Steigemann,
 5 "Structure of the Complex formed by Bovine Trypsin and
 Bovine Pancreatic Trypsin Inhibitor",
 J Mol Biol (1974), 89:73-101.
- HUBE75:
 10 Huber, R, W Bode, D Kukla, and U Kohl,
 "The Structure of the Complex Formed by Bovine Trypsin and
 Bovine Pancreatic Trypsin Inhibitor: III. Structure of the
 Anhydrotrypsin-Inhibitor Complex",
 Biophys Struct Mechan (1975), 1:189-201.
- 15 HUBE77:
 Huber, R, W Bode, D Kukla, U Kohl, CA Ryan,
 "The structure of the complex formed by bovine trypsin
 and bovine pancreatic trypsin inhibitor III. Structure of
 20 the anhydro-trypsin-inhibitor complex.",
 Biophys Struct Mech (1975), 1(3)189-201.
- HUTC87:
 Hutchinson, DCS,
 25 "The role of proteases and antiproteases in bronchial
 secretions",
 Eur J Respir Dis (1987), 71(Suppl.153)78-85.
- HYNE90:
 30 Hynes, TR, M Randal, LA Kenedy, C Eigenbrot, and
 AA Kossiakoff,
 "X-ray crystal structure of the protease inhibitor domain
 of Alzheimer's amyloid beta-protein precursor",
 Biochemistry (1990), 29:10018-10022.
- 35 ILIC89:
 Il'ichev, AA, OO Minenkova, SI Tat'kov, NN Karpyshev, AM
 Eroshkin, VA Petrenko, and LS Sandakhchiev,
 "[Production of a viable variant of the M13 phage with a
 40 foreign peptide inserted into the basic coat
 protein]<Original> Poluchenie zhiznesposobnogo varianta
 faga M13 so vstroennym chuzherodnym peptidom v osnovnoi
 belok obolochki",
 Dokl Akad Nauk SSSR, (1989), 307(2)481-3.
- 45 INOU82:
 Inouye, H, W Barnes, and J Beckwith,
 "Signal Sequence of Alkaline Phosphatase of Escherichia
coli",
 50 J Bacteriol (1982), 149(2)434-439.

T05230-5696850

Inouye, M, and R Sarma, Editors,
Protein Engineering: Applications in Science, Medicine,
and Industry.,

ITOK79:

10 Synthesis of procoat protein in Escherichia coli infected
with bacteriophage M13.",
Proc Natl Acad Sci USA (1979), 76:1199-1203.

15 Janatova, J, KBM Reid, and AC Willis,
"Disulfide Bonds Are Localized within the Short Consensus
Repeat Units of Complement Regulatory Proteins: C4b-
Binding Protein",
Biochem (1989), 28:4754-61.

Janin, J, and C Chothia,
"Domains in Proteins: Definitions, Location, and
Structural Principles",
25 *Methods in Enzymology* (1985), 115(28)420-430.

Jennings, PA, MM Bills, DO Irving, and JS Mattick,
"Fimbriae of *Bacteroides nodosus*: protein engineering of
30 the structural subunit for the production of an exogenous
peptide",
Protein Eng. (Jan 1989), 2(5)365-9.

35 Jering, H, and H Tschesche,
"Replacement of Lysine by Arginine, Phenylalanine, and
Tryptophan in the Reactive Site of the Trypsin-Kallikrein
Inhibitor (Kunitz)",
Angew Chem internat Edit (1974), 13:662-3.

Jering, H, and H Tschesche,
"Replacement of Lysine by Arginine, Phenylalanine, and
Tryptophan in the Reactive Site of the Bovine Trypsin-
Kallekrein Inhibitor (Kunitz) and Change of the Inhibitory
Properties",
Eur J Biochem (1976), 61:453-63.

50 Joubert, FJ,
"Trypsin Isoinhibitors from Momordica Repens Seeds",
Phytochemistry (1984), 23:1401-6.

JUDD85:

Judd, RC,

- "Structure and surface exposure of protein IIs of
 5 *Neisseria gonorrhoeae* JS3",
Infect Immun (1985), 48(2)452-7.

JUDD86:

Judd, RC,

- 10 "Evidence for N-terminal exposure of the protein IA
 subclass of *Neisseria gonorrhoeae* protein I",
Infect Immun (1986), 54(2)408-14.

KABS84:

- 15 Kabsch, W, and C Sander,

"On the use of sequence homologies to predict protein
 structure: identical pentapeptides can have completely
 different conformations",
Proc Natl Acad Sci USA (1984), 81(4)1075-8.

20

KAIS87a:

Kaiser, CA, D Preuss, P Grisafi, and D Botstein,

"Many Random Sequences Functionally Replace the Secretion
 Signal Sequence of Yeast Invertase",

- 25 *Science* (1987), 235:312-7.

KAOR88:

Kao, RC, NG Wehner, KM Skubitz, BH Gray, and JR Hoidal,

"Proteinase 3, A Distinct Human Polymorphonuclear

- 30 *Leukocyte Proteinase that Produces Emphysema in Hamsters*",
J Clin Invest (1988), 82:1963-73.

KAPL78:

Kaplan, DA, L Greenfield, and G Wilcox,

- 35 "Molecular Cloning of Segments of the M13 Genome.",
 in The Single-Stranded DNA Phages, Denhardt, DT,
 D Dressler, and DS Ray editors, Cold Spring Harbor
 Laboratory, 1978., p461-467.

- 40 KATZ86:

Katz, BA, and A Kossiakoff,

"The Crystallographically Determined Structures of
 Atypical Stained Disulfides Engineered into Subtilisin",
J Biol Chem (1986), 261(33)15480-85.

45

KATZ90:

Katz, B, and AA Kossiakoff,

"Crystal Structures of Subtilisin BPN' Variants Containing
 Disulfide Bonds and Cavities: Concerted Structural
 50 Rearrangements Induced by Mutagenesis",
Proteins, Struct, Funct, and Genet (1990), 7:343-57.

FOI 230 56096860

KAUM86:

- Kaumerer, JF, JO Polazzi, and MP Kotick,
 "The mRNA for a proteinase inhibitor related to the HI-30
 domain of inter- α -trypsin inhibitor also encodes α_1 -
 5 microglobulin (protein HC)",
 Nucleic Acids Res (1986), 14:7839-7850.

KIDO88:

- Kido, H, Y Yokogoshi, and N Katunuma,
 10 "Kunitz-type Protease Inhibitor Found in Rat Mast Cells",
 J Biol Chem (1988), 263:18104-7.

KIDO90:

- Kido, H, A Fukutomi, J Schelling, Y Wang, B Cordell, and N
 15 Katunuma,
 "Protease-Specificity of Kunitz Inhibitor Domain of
 Alzheimer's Disease Amyloid Protein Precursor",
 Biochem & Biophys Res Comm (16 Mar 1990), 167(2)716-21.

KING86:

- King, TC, R Sirdeskmukh, and D Schlessinger,
 "Nucleolytic processing of ribonucleic acid transcripts in
 procaryotes",
 20 Microbiol Rev (1986), 50(4)428-51.

KISH85:

- Kishore, R, and P Balaram,
 "Stablization of gamma-Turn Conformations in Peptides by
 Disulfide Bridges",
 30 Biopolymers (1985), 24:2041-43.

KOB89:

- Kobayashi, Y, T Ohkubo, Y Kyogoku, Y Nishiuchi, S
 Sakakibara, W Braun, nad N Go,
 35 "Solution Conformation of Conotoxin GI Determined by ^1H
 Nuclear Magnetic Resonance Spectroscopy and Distance
 Geometry Calculations",
 Biochemistry (1989), 28:4853-60.

KUB89:

- Kubota, H, Y Hidaka, H Ozaki, H Ito, T Hirayama, Y Takeda,
 and Y Shimonishi,
 "A Long-acting Heat-Stable Enterotoxin Analog of
 Enterotoxigenic *Escherichia coli* with a Single D-Amino
 45 Acid.",
 Biochem Biophys Res Comm (1989), 161:229-235.

KUHN85a:

- Kuhn, A, and W Wickner,
 50 "Conserved Residues of the Leader Peptide Are Essential
 for Cleavage by Leader Peptidase.",
 J Biol Chem (1985), 260:15914-15918.

- KUHN85b:
Kuhn, A, and W Wickner,
"Isolation of Mutants in M13 Coat Protein That Affect Its
5 Synthesis, Processing, and Assembly into Phage.",
J Biol Chem (1985), 260:15907-15913.
- KUHN87:
Kuhn, A,
10 "Bacteriophage M13 Procoat Protein Inserts into the Plasma
Membrane as a Loop Structure.",
Science (1987), 238:1413-1415.
- KUHN88:
15 Kuhn, A,
"Alterations in the extracellular domain of M13 procoat
protein make its membrane insertion dependent on secA and
secY",
Eur J Biochem (1988), 177(2)267-71.
20
- KUKS89:
Kuks, PFM, C Creminon, A-M Leseney, J Bourdais, A Morel,
and P Cohen,
"Xenopus laevis Skin Arg-Xaa-Val-Arg-Gly-endoprotease",
25 J Biol Chem (1989), 264(25)14609-12.
- KUOM90:
Kuo, MD, SS Huang, and JS Huang,
"Acidic fibroblast growth factor receptor purified from
30 bovine liver is a novel protein tyrosine kinase."
J Biol Chem (1990), 265(27)16455-63.
- KUPE90:
Kupersztach, YM, K Tachias, CR Moomaw, LA Dreyfus, R
35 Urban, C Slaughter, and S Whipp,
"Secretion of Methanol-Insoluble Heat-Stable Enterotoxin
(ST_B): Energy- and secA-Dependent Conversion of Pre-ST_B to
an Intermediate Indistinguishable from the Extracellular
Toxin",
40 J Bacteriol (1990), 172(5)2427-32.
- LAMB90:
Lambert, P, H Kuroda, N Chino, TX Watanabe, T Kimura, and
S Sakakibara,
45 "Solution Synthesis of Charybdotoxin (ChTX), A K⁺ Channel
Blocker",
Biochem Biophys Res Comm (1990), 170(2)684-690.

1990-05-05 15:00:00

Landick, R, and C Yanofsky,
"Transcription Attenuation",
Volume 2, Chapter 77, p 1276-1301,

- 5 Escherichia coli and Salmonella typhimurium: Cellular and
Molecular Biology,
Neidhardt, FC, Editor-in-Chief,
Amer Soc for Microbiology, Washington, DC, 1987.
- 10 LASK80:
Laskowski, M, Jr, and I Kato,
"Protein Inhibitors of Proteases",
Ann Rev Biochem (1980), 49:593-626.
- 15 LAZU83:
Lazure, C, NG Seidah, M Chretien, R Lallier, and
S St-Pierre,
"Primary structure determination of Escherichia coli heat-
stable enterotoxin of porcine origin",
20 Canadian J Biochem Cell Biol (1983), 61:287-92.
- LECO87:
Lecomte, J TJ, D Kaplan, M Llinas, E Thunberg, and
G Samuelsson,
25 "Proton Magnetic Resonance Characterization of Phoratoxins
and Homologous Proteins Related to Crambin",
Biochemistry (1987), 26:1187-94.
- LEEB71:
30 Lee, B, and FM Richards,
"The interpretation of protein structures: estimation of
static accessibility.",
J Mol Biol (1971), 55:(3)379-400,
- 35 LEEC83:
Lee, CH, SL Moseley, HW Moon, SC Whipp, CL Gyles, and
M So,
"Characterization of the Gene Encoding Heat-Stable Toxin
II and Preliminary Molecular Epidemiological Studies of
40 Enterotoxigenic Escherichia coli Heat-Stable Toxin II
Producers",
Infection and Immunity (1983), 42:264-268.
- LEEC86:
45 Lee, C, and J Beckwith,
"Cotranslational and Posttranslational Protein
Translocation in Prokaryotic Systems.",
Ann Rev Cell Biol (1986), 2:315-336.

- LENG89b:
 Le-Nguyen, D, D Nalis, and B Castro,
 "Solid phase synthesis of a trypsin inhibitor isolated
 from the Cucurbitaceae Ecballium elaterium",
 5 Int J Peptide Protein Res (1989), 34:492-97.
- LISS85:
 Liss, LR, BL Johnson, and DB Oliver,
 "Export defect adjacent to the processing site of
 10 staphylococcal nuclease is suppressed by a prlA
 mutation",
 J Bacteriol (1985), 164(2)925-8.
- LOPE85a:
 15 Lopez, J, and RE Webster,
 "Assembly site of bacteriophage f1 corresponds to adhesion
 zones between the inner and outer membranes of the host
 cell",
 J Bacteriol (1985), 163(3)1270-4.
- LOPE85b:
 Lopez, J, and RE Webster,
 "fipB and fipC: two bacterial loci required for
 morphogenesis of the filamentous bacteriophage f1",
 25 J Bacteriol (1985), 163(3)900-5.
- LOSI86:
 Losick, R, P Youngman, and PJ Piggot,
 "Genetics of Endospore formation in Bacillus subtilis",
 30 Ann Rev Genet (1986), 20:625-669.
- LUGT83:
 Lugtenberg, B, and L van Alphen,
 "Molecular Architecture and Function of the Outer Membrane
 35 of Escherichia coli and other Gram-Negative Bacteria",
 Biochim Biophys Acta (1983), 737:51-115.
- LUIT83:
 Luiten, RGM, JGG Schoenmakers, and RNH Konings,
 40 "The major coat protein gene of the filamentous
Pseudomonas aeruginosa phage Pf3: absence of an N-terminal
 leader signal sequence",
 Nucleic Acids Research (1983), 11(22)8073-85.
- LUIT85:
 45 Luiten, RGM, DG Putterman, JGG Schoenmakers, RNH Konings,
 and LA Day,
 "Nucleotide Sequence of the Genome of Pf3, an IncP-1
 Plasmid-Specific Filamentous Bacteriophage of Pseudomonas
 50 aeruginosa",
 J Virology, (1985), 56(1)268-276.

FO5230-56056860

5 DNA (1987), 6(2)129-37.

10 Beta Strand, Turn, and Other Conformations. Secondary Structures of Copper Proteins", J Inorgan Biochem (1986), 27:151-62.

25 within the *Escherichia coli* outer membrane protein OmpA",
J Biol Chem (1988), 263(35)19053-9.

J Mol Biol (1980), 140:149-181.

40

45

50 Science (1986), 233:1403-1408.

MANO88:

Manoil, C, D Boyd, and J Beckwith,
 "Molecular genetic analysis of membrane protein topology",
 Topics in Genetics (1988), 4(8)223-6.

5

MARK86:

Marks, CB, M Vasser, P Ng, W Henzel, and S Anderson,
 "Production of native, correctly folded bovine pancreatic
 trypsin inhibitor in Escherichia coli",
 J Biol Chem (1986), 261:7115-7118.

10

MARK87:

Marks, CB, H Naderi, PA Kosen, ID Kuntz, and S Anderson,
 "Mutants of Bovine Pancreatic Trypsin Inhibitor Lacking
 Cysteines 14 and 38 Can Fold Properly",
 Science (1987), 235:1370-1373.

15

MARQ83:

Marquart, M, J Walter, J Deisinhoffer, W Bode, and
 R Huber,
 "The geometry of the reactive site and of the peptide
 groups in trypsin, trypsinogen, and its complexes with
 inhibitors",
 Acta Cryst, B (1983), 39:480ff.

20

25

MARV75:

Marvin, DA and EJ Wachtel,
 "Structure and assembly of filamentous bacterial viruses",
 Nature (1975), 253:19-23.

30

MARV78:

Marvin, DA,
 "Structure of the Filamentous Phage Virion.",
 in The Single-Stranded DNA Phages, Denhardt, DT,
 D Dressler, and DS Ray editors, Cold Spring Harbor
 Laboratory, 1978., p583-603.

35

MARV80:

Marvin, D, and L Makowski,
 "Helical Viruses",
 Progr Clin Biol Res (1980), 40:347-48.

40

MASS90:

Massefski, W, Jr, AG Redfield, DR Hare, and C Miller,
 "Molecular Structure of Charybdotoxin, a Pore-Directed
 Inhibitor of Potassium Ion Channels",
 Science (3 Aug 1990), 249:521-524.

45

006230 56096860

Matsumura, M, WJ Becktel, M Levitt, and BW Matthews,
"Stabilization of phage T4 lysozyme by engineered
disulfide bonds",

- 40 MESS77:
Messing, J, B Gronenborn, B Muller-Hill, and
PH Hofschneider,
"Filamentous coliphage M13 as a cloning vehicle: insertion
of a HindII fragment of the lac regulatory region in M13
45 replicative form in vitro.",
Proc Natl Acad Sci USA (1977), 74:3642-6.

MESS78:

- Messing, J, and B Gronenborn,
 "The Filamentous Phage M13 as a Carrier DNA for Operon
 Fusions In Vitro.", in The Single-Stranded DNA Phages,
 5 Denhardt, DT, D Dressler, and DS Ray editors, Cold Spring
 Harbor Laboratory, 1978.,p449-453.

MILL87a:

- Miller, S, J Janin, AM Lesk, and C Chothia,
 10 "Interior and Surface Monomeric Proteins",
 J Mol Biol (1987), 196:641-656.

MILL87b:

- Miller, ES, J Karam, M Dawson, M Trojanowska, P Gauss,
 15 and L Gold,
 "Translational repression: biological activity of
 plasmid-encoded bacteriophage T4 RegA protein.",
 J Mol Biol (1987), 194:397-410.

MISR88a:

- Misra, R, and SA Benson,
 "Genetic identification of the pore domain of the OmpC
 porin of Escherichia coli K-12",
 J Bacteriol (1988), 170(8)3611-7.

25

MISR88b:

- Misra, R, and SA Benson,
 "Isolation and Characterization of OmpC Porin Mutants with
 Altered Pore Properties",
 30 J Bacteriol (1988), 170:528-33.

MOLL89:

- Molla, A, A Charbit, A Le Guern, A Ryter, and M Hofnung,
 "Antibodies against synthetic peptides and the topology of
 35 LamB, an outer membrane protein from Escherichia coli
 K12",
 Biochem (1989), 28(20)8234-41.

MORS87:

- Morse, SA, TA Mietzner, G Bolen, A Le Faou, and
 40 G Schoolnik,
 "Characterization of the major iron-regulated protein of
 Neisseria gonorrhoeae and Neisseria meningitidis",
 Antonie Van Leeuwenhoek (1987), 53(6)465-9.

45

MORS88:

- Morse, SA, C-Y Chen, A LeFaou, and TA Meitzner,
 "A Potential Role for the Major Iron-Regulated Protein
 Expressed by Pathogenic Neisseria Species",
 50 Rev Infect Dis (1988), 10(Suppl 2)S306-10.

Moses, PB, and K Horiuchi,
"Effects of Transposition and Deletion upon Coat Protein
Gene Expression in Bacteriophage f1",
Virology (1982), 119:231-244.

Moser, R, RM Thomas, and B Gutte,
"An Artificial Crystalline DDT-binding polypeptide",
FEBS Letters (1983), 157:247-251.

Moser, R, S Klauser, T Leist, H Langen, T Epprecht, and
B Gutte,
"Applications of Synthetic Peptides",
Angew Chemie, Int Edition English (1985), 24(9)719-27.

Moser, R, S Frey, K Muenger, T Hehlhans, S Klauser, H Langen, E-L Winnacker, R Mertz, and B Gutte, "Expression of the synthetic gene of an artificial DDT-binding polypeptide in Escherichia coli", Protein Engineering (1987), 1:339-343.

Nadel, JA, and B Borson,
"Secretion and ion transport in airways during
inflammation",
Biorheology (1987), 24:541-549.

Nadel, JA,
"Neutrophil Proteases and Mucus Secretion",
1990 Cystic Fibrosis Meeting, Arlington, Va., p156.

Nakashima, Y, B Frangione, RL Wiseman, WH Konigsberg,
"Primary Structure of the Major Coat Protein of the
Filamentous Bacterial Viruses, If1 and Ike",
J Biol Chem (1981), 256(11)5792-7.

Nakae, T, J Ishii, and T Ferenci,
"The Role of the Maltodextrin-binding Site in Determining
the Transport Properties of the LamB Protein",
J Biol Chem (1986), 261:622-26.

Nakae, T,
"Outer-Membrane Permeability of Bacteria",
CRC Crit Rev Microbiol (1986), 13:1-62.

NAKA87:

Nakamura, T, T Hirai, F Tokunaga, S Kawabata, and
S Iwanaga,

- "Purification and Amino Acid Sequence of Kunitz-type
5 Protease Inhibitor Found in the Hemocytes of Horseshoe
Crab (Tachypleus tridentatus)",
J Biochem (1987), 101:1297-1306.

NICH88:

- 10 Nicholson, H, WJ Bechtel, and BW Matthews,
"Enhanced protein thermostability from designed mutations
that interact with α -helix dipoles",
Nature (1988), 336:651-56.

NIKA84:

Nikaido, H, and HCP Wu,
"Amino acid sequence homology among the major outer
membrane proteins of Escherichia coli",
Proc Natl Acad Sci USA (1984), 81:1048-52.

20

NILE89:

Niles, JL, RT McCluskey, MF Ahmad, and MA Arnaout,
"Wgener's Granulomatosis Autoantigen Is a Novel Neutrophil
Serine Proteinase",

- 25 Blood (1989), 74(6)1888-93.

NISH82:

Nishiuchi, Y, and S Sakakibara,
"Primary and secondary structure of conotoxin GI, a
30 neurotoxic tridecapeptide from a marine snail",
FEBS Lett (1982), 148:260-2.

NISH86:

Nishiuchi, Y, K Kumagaye, Y Noda, TX Watanabe, and S
35 Sakakibara,
"Synthesis and secondary-structure determination of
omega-conotoxin GVIA: a 27-peptide with three
intramolecular disulfide bonds",
Biopolymers, (1986), 25:S61-8.

40

NORR89a:

Norris, K, and LC Petersen,
"Aprotinin analogues and process for the production
thereof",

- 45 European Patent Application 0 339 942 A2.

NORR89b:

Norris, K, F Norris, S BJorn,
"Aprotinin Homologues and Process for the Production of
50 Aprotinin and aprotinin homologues in Yeast",
PCT patent application W089/01968.

F05230-5056860

OAST88:

Oas, TG, and PS Kim,

"A peptide model of a protein folding intermediate",
Nature (1988), 336:42-48.

5

ODOM90:

Odom, L,

"Inter- α -trypsin inhibitor: a plasma proteinase inhibitor
with a unique chemical structure",10 Int J Biochem (1990), 22:925-930.

OHKA81:

Ohkawa, I, and RE Webster,

"The Orientation of the Major Coat Protein of

15 Bacteriophage f1 in the Cytoplasmic Membrane of Escherichia coli.",J Biol Chem (1981), 256:9951-9958.

OKAM87:

20 Okamoto, K, K Okamoto, J Yukitake, Y Kawamoto, and
A Miyama,"Substitutions of Cysteine Residues of Escherichia coli
Heat-Stable Enterotoxin by Oligonucleotide-Directed
Mutagenesis",25 Infection and Immunity (1987), 55:2121-2125.

OKAM88:

Okamoto, K, K Okamoto, J Yukitake, and A Miyama,

"Reduction of Enterotoxic Activity of Escherichia coli30 Heat-Stable Enterotoxin by Substitution for an Aspartate
Residue",Infection and Immunity (1988), 56:2144-8.

OKAM90:

35 Okamoto, K, and M Takahara,

"Synthesis of Escherichia coli Heat-Stable Enterotoxin STp
as a Pre-Pro Form and Role of the Pro Sequence in
Secretion",J Bacteriol (1990), 172(9)5260-65.

40

OLIP86:

Oliphant, AR, AL Nussbaum, and K Struhl,

"Cloning of random-sequence oligodeoxynucleotides",
Gene (1986), 44:177-183.

45

OLIP87:

Oliphant, AR, and K Struhl

"The Use of Random-Sequence Oligonucleotides for
Determining Consensus Sequences", in50 Methods in Enzymology 155 (1987)568-582.

Editor Wu, R; Academic Press, New York.

T06230"56036860

Oliver, D,
"Protein Secretion in Escherichia coli.",
Ann Rev Microbiol (1985), 39:615-648.

OLIV85b:

Olivera, BM, WR Gray, R Zeikus, JM McIntosh, J Varga,
J Riviera, V de Santos, and LJ Cruz,
"Peptide Neurotoxins from Fish Hunting Cone Snails",
Science (1985), 230:1338-43.

Olivera, BM, LJ Cruz, V de Santos, GW LeCheminant, D Griffin, R Zeikus, JM McIntosh, R Galyean, J Varga, WR Gray, et al.

"Neuronal calcium channel antagonists. Discrimination between calcium channel subtypes using omega-conotoxin from *Conus magus* venom",
Biochemistry, (1987), 26(8)2086-90.

OLIV90a:

Olivera, BM, J Rivier, C Clark, CA Ramilo, GP Corpuz,
FC Abogadie, EE Mena, SR Woodward, DR Hillyard, LJ Cruz,
"Diversity of Conus Neuropeptides",
Science, (20 July 1990), 249:257-263.

Olivera, BM, DR Hillyard, J Rivier, S Woodward, WR Gray,
G Corpuz, LJ Cruz,

"Conotoxins: Targeted Peptide Ligands from Snail Venoms",
Chapter 20 in Marine Topxins, American Chemical Society,
1990.

35 Oltersdorf, T, LC Fritz, DB Schenk, I Lieberburg,
KL Johnson-Wood, EC Beattie, PJ Ward, RW Blacher,
HF Dovey, and S Sinha,

"The Secreted form of the Alzheimer's amyloid precursor protein with the Kunitz domain is protease nexin-II", Nature (1989), 341:144-7.

Orndorff, PE, and S Falkow,
"Nucleotide Sequence of pilA, the Gene Encoding the
Structural Component of Type 1 Pili in Escherichia coli",
J Bacteriol (1985), 162:454-7.

45

PANT90:

Pantoliano, MW, and RC Ladner,
 "Computer Designed Stabilized Proteins and Method for
 Producing Same",

- 5 US Patent 4,908,773, 13 March 1990.

PAOL86:

Paoletti, E, and D Panicali,
 "Modified Vaccinia Virus",

- 10 US Patent 4,603,112, July 29, 1986.

PAPA82:

Papamokos, E, E Weber, W Bode, R Huber, M Empie, I Kato,
 and M Laskowski Jr,

- 15 "Crystallographic Refinement of Japanese Quail Ovomuroid,
 a Kazal-type Inhibitor, and Model Building Studies of
 Complexes with Serine Proteases",
 J Mol Biol (1982), 158:515-537.

20 PARD89:

Pardi, A, A Galdes, J Florance, and D Maniconte,
 "Solution Structures of α -Conotoxin G1 Determined by Two-
 Dimensional NMR Spectroscopy",
 Biochemistry (1989), 28:5494-5501.

25

PARG87:

Parge, HE, DE McRee, MA Capozza, SL Bernstein, ED Getzoff,
 and JA Tainer,

- 30 "Three dimensional structure of bacterial pili",
 Antonie Van Leeuwenhoek (1987), 53(6)447-53.

PARM88:

Parmley, SF, and GP Smith,
 "Antibody-selectable filamentous fd phage vectors:
 affinity purification of target genes",
 Gene (1988), 73:305-318.

35

PARR88:

Parraga, G, SJ Horvath, A Eisen, WE Taylor, L Hood,
 ET Young, RE Klevit,
 "Zinc-Dependent Structures of a Single-Finger Domain of
 Yeast ADR1",
 Science (1988), 241:1489-92.

40

45 PEAS88:

Pease, JHB, and DE Wemmer,
 Biochem (1988), 27:8491-99.

FO6290-5695860

PEAS90:

Pease, JHB, RW Storrs, and DE Wemmer,
 "Folding and activity of hybrid sequence, disulfide-
 stabilized peptides",

- 5 Proc Natl Acad Sci USA (1990), 87:5643-47.

PEET85:

Peeters, BPH, RM Peters, JGG Schoenmakers, and
 RNH Konings,

- 10 "Nucleotide Sequence and Genetic Organization of the
 Genome of the N-Specific Filamentous Bacteriophage Ike:
 Comparison with the Genome of the F-Specific Filamentous
 Phages M13, fd, and f1",
 J Mol Biol (1985), 181:27-39.

15

PEET87:

Peeters, BPH, JGG Schoenmakers, and RNH Konings,
 "Comparison of the DNA Sequences Involved in Replication
 and Packaging of the Filamentous Phages IKE, and Ff (M13,
 fd, and f1)",
 DNA (1987), 6(2)139-147.

20

PERR84:

Perry, LJ, and R Wetzel,
 "Disulfide Bond Engineered into T4 Lysozyme: Stabilization
 of the Protein Toward Thermal Inactivation",
 Science (1984), 226:555-7.

25

PERR86:

Perry, LJ, and R Wetzel,
 "Unpaired Cysteine-54 Interferes with the Ability of an
 Engineered Disulfide To Stabilize T4 Lysozyme",
 Biochem (1986), 25:733-39.

30

PETE89:

Peterson, MW,
 "Neutrophil cathepsin G increases transendothelial albumin
 flux",
 J Lab Clin Med (1989), 113(3)297-308.

35

40

PONT88:

Ponte, P, P Gonzalez-DeWhitt, J Schilling, J Miller,
 D Hsu, B Greenberg, K Davis, W Wallace, I Liederburg,
 F Fuller, and B Cordell,
 "A new A4 amyloid mRNA contains a domain homologous to
 serine proteinase inhibitors",
 Nature (1988), 331:525-7.

45

T06230 56096860

5 J Mol Biol (1983), 171:401-418.

10 Understanding of Protein-Ligand Interactions.",
in Crystallography in Molecular Biology, Moras, D. et
al., editors, Plenum Press, 1987.

15 Randall, LL, SJS Hardy, and JR Thom,
"Export of Protein: A Biochemical View",
Ann Rev Microbiol (1987), 41:507-41.

20 Rasched, I, and E Oberer,
"Ff Coliphages: Structural and Functional Relationships",
Microbiol Rev (1986) 50:401-427.

25 Rashin, A,
"Prediction of Stabilities of Thermolysin Fragments",
Biochemistry (1984), 23:5518.

30 Ray, C, KM Tatti, CH Jones, and CP Moran Jr,
"Genetic Analysis of RNA Polymerase-Promoter Interaction
during Sporulation in Bacillus subtilis",
J Bacteriol (1987), 169(5)1807-1811.

Reidhaar-Olson, JF, and RT Sauer,
"Combinatorial Cassette Mutagenesis as a Probe of the
Information Content of Protein Sequences",
Science (1988), 241:53-57.

Reid, J, H Fung, K Gehring, PE Klebba, and H Nikaido,
"Targeting of porin to the outer membrane of Escherichia
coli. Rate of trimer assembly and identification of a
45 dimer intermediate",
J Biol Chem (1988), 263(16)7753-9.

Rest, RF,
50 "Human Neutrophil and Mast Cell Proteases Implicated in
Inflammation",
Meth Enzymol (1988), 163:309-27.

- RICH81:
Richardson, JS,
"The Anatomy and Taxonomy of Protein Structure",
5 Adv Protein Chemistry (1981), 34:167-339.
- RICH86:
Richards, JH,
"Cassette mutagenesis shows its strength.",
10 Nature (1986), 323:187.
- RITO83:
Ritonja, A, B Meloun, and F Gubensek,
"The Primary Structure of Vipera ammodytes venom
15 chymotrypsin inhibitor",
Biochim Biophys Acta (1983), 746:138-145.
- RIVI87b:
Rivier, J, R Galyean, WR Gray, A Azimi-Zonooz,
20 JM McIntosh, LJ Cruz, and BM Olivera,
"Neuronal calcium channel inhibitors. Synthesis of
omega-conotoxin GVIA and effects on ⁴⁵Ca uptake by
synaptosomes",
J Biol Chem, (1987), 262(3)1194-8.
25
- ROBE86:
Roberts, S, and AR Rees
"The cloning and expression of an anti-peptide antibody: a
system for rapid analysis of the binding properties of
30 engineered antibodies.",
Protein Engineering (1986), 1:59-65.
- RONC90:
Ronco, J, A Charbit, and M Hofnung,
35 "Création of targets for proteolytic cleavage in the LamB
protein of E coli K12 by genetic insertion of foreign
sequences: implications for topological studies",
Biochimie (1990), 72(2-3)183-9.
- 40 ROSE85:
Rose, GD,
"Automatic Recognition of Domains in Globular Proteins",
Methods in Enzymology (1985), 115(29)430-440.
- 45 ROSS81:
Rossman, M, and P Argos,
"Protein Folding.",
Ann Rev Biochem (1981), 50:497-532.

RUEH73:

- Ruehlmann, A, D Kukla, P Schwager, K Bartels, and R Huber,
 "Structure of the Complex formed by Bovine Trypsin and
 Bovine Pancreatic Trypsin Inhibitor: Crystal Structure
 5 Determination and Stereochemistry of the Contact Region",
 J Mol Biol (1973), 77:417-436.

RUSS81:

- Russel, M, and P Model,
 10 "A mutation downstream from the signal peptidase cleavage
 site affects cleavage but not membrane insertion of phage
 coat protein.",
 Proc Natl Acad Sci USA (1981), 78:1717-1721.

SALI64:

- Salivar, WO, H Tzagoloff, and D Pratt,
 "Some physical, chemical, and biological properties of the
 rod-shaped coliphage M13",
 15 Virology (1964), 24:359-71.

20

SALI87:

- Salier, JP, M Diarra-Mehrpour, R Sesboue, J Bourguignon,
 R Benarous, I Ohkubo, S Kurachi, K Kurachi, and JP Martin,
 "Isolation and characterization of cDNAs encoding the
 25 heavy chain of human inter-alpha-trypsin inhibitor (IaTI):
 Unambiguous evidence for multipolypeptide chain structure
 of IaTI",
 Proc Nat Acad Sci USA (1987), 84:8271-8276.

SALI88:

- Sali, D, M Bycroft, and AR Fersht,
 "Stabilization of protein structure by interaction of α -
 helix dipole with a charged side chain",
 30 Nature (1988), 335:740-3.

35

SALI90:

- Salier, J-P,
 "Inter- α -trypsin inhibitor: emergence of a family within
 the Kunitz-type protease inhibitor superfamily",
 40 TIBS (1990), 15:435-439.

SALV87:

- Salvesen, G, D Farley, J Shuman, A Przybyla, C Reilly, and
 J Travis,
 45 "Molecular Cloning of Human Cathepsin G: Structural
 Similarity to Mast Cell and Cytotoxic T Lymphocyte
 Proteinases",
 Biochem (1987), 26:2289-93.

FO5230 56096860

SAMB89:

Sambrook, J, EF Fritsch, and T Maniatis,
Molecular Cloning, A Laboratory Manual, Second Edition,
 Cold Spring Harbor Laboratory, 1989.

5

SASA84:

Sasaki, T,

"Amino Acid Sequence of a Novel Kunitz-type chymotrypsin
 inhibitor from hemolymph of silkworm larvae, Bombyx mori",

10 FEBS Lett (1984), 168:227-230.

SAUE86:

Sauer, RT, K Hehir, RS Stearman, MA Weiss,
 A Jeitler-Nilsson, EG Suchanek, and CO Pabo,

15 "An Engineered Intersubunit Disulfide Enhances the
 Stability and DNA Binding of the N-Terminal Domain of λ
 Repressor",
 Biochem (1986), 25:5992-98.

20

SCHA78:

Schaller, H, E Beck, and M Takanami,

"Sequence and Regulatory Signals of the Filamentous Phage
 Genome.", in The Single-Stranded DNA Phages, Denhardt,
 D.T., D. Dressler, and D.S. Ray editors, Cold Spring

25 Harbor Laboratory, 1978., p139-163.

SCHN86:

Schnabel, E, W Schroeder, and G Reinhardt,
 "[Ala₂^{14,38}]Aprotinin: Preparation by Partial

30 Desulphurization of Aprotinin by Means of Raney Nickel and
 Comparison with other Aprotinin Derivatives",
 Biol Chem Hoppe-Seyler (1986), 367:1167-76.

SCHN88a:

35 Schnabel, E, G Reinhardt, W Schroeder, H Tschesche,
 HR Wenzel, and A Mehlich,

"Enzymatic Resynthesis of the 'Reactive Site' Bond in the
 Modified Aprotinin Derivatives [Seco-15/16]Aprotinin and
 [Di-seco-15/16,39/40]Aprotinin",

40 Biol Chem Hoppe-Seyler (1988), 369:461-8.

SCHU79:

Schulz, GE, and RH Schirmer,
Principles of Protein Structure,

45 Springer-Verlag, New York, 1979.

T06230"56096860

SCHW87:

- Schwarz, H, HJ Hinz, A Mehlich, H Tschesche, and
 HR Wenzel,
 "Stability studies on derivatives of the bovine
 5 pancreatic trypsin inhibitor.",
 Biochemistry (1987), 26:(12)p3544-51.

SCOT87a:

- Scott, MJ, CS Huckaby, I Kato, WJ Kohr, M Laskowski Jr.,
 10 M-J Tsai and BW O'Malley,
 "Ovoinhibitor Introns Specify Functional Domains as in the
 Related and Linked Ovomucoid Gene",
 J Biol Chem (1987), 262(12)5899-5907.

15 SCOT87b:

- Scott, CF, HR Wenzel, HR Tschesche, and RW Colman,
 "Kinetics of Inhibition of Human Plasma Kallikrein by a
 Site-Specific Modified Inhibitor Arg¹⁵-Aprotinin:
 Evaluation Using a Microplate System and Comparison With
 20 Other Proteases",
 Blood (1987), 69:1431-6.

SCOT90:

- Scott, JK, and GP Smith,
 25 "Searching for Peptide Ligands with an Epitope Library",
 Science, (27 July 1990), 249:386-390.

SEKI85:

- Sekizaki, T, H Akaski, and N Terakado,
 30 "Nucleotide sequences of the genes for Escherichia coli
 heat-stable enterotoxin I of bovine, avian, and porcine
 origins",
 Am J Vet Res (1985), 46:909-12.

35 SELL87:

- Selloum, L, M Davril, C Mizon, M Balduyck, and J Mizon,
 "The effect of the glycosaminoglycan chain removal on some
 properties of the human urinary trypsin inhibitor",
 Biol Chem Hoppe-Seyler (1987), 368:47-55.

40

SERW87:

- Serwer, P,
 "Review: Agarose Gel Electrophoresis of Bacteriophages and
 Related Particles",
 45 J Chromatography (1987), 418:345-357.

T06230 55096860

SHIM87:

Shimonishi, Y, Y Hidaka, M Koizumi, M Hane, S Aimoto, T
Takeda, T Miwatani, and Y Takeda,

- 5 "Mode of disulfide bond formation of a heat-stable
enterotoxin (ST_h) produced by a human strain of
enterotoxigenic Escherichia coli",
FEBS Lett (1987), 215:165-170.

SHOR81:

- 10 Shortle, D, D Koshland, GM Weinstock, and D Botstein,
"Segment-directed mutagenesis: Construction in vitro of
point mutations limited to a small predetermined region of
a circular DNA molecule",
Proc Natl Acad Sci USA (1980), 77:5375-79.

15

SHOR85:

Shortle, D, and B Lin,
"Genetic Analysis of Staphylococcal Nuclease:
Identification of Three Intragenic 'Global' Suppressors
20 of Nuclease-Minus Mutations.",
Genetics (1985), 110:539-555.

SIEK87:

- 25 Siekmann, J, HR Wenzel, W Schroeder, H Schutt, E
Truscheit, A Arens, E Rauenbusch, WH CHazin, K Wutrich,
and H Tschesche,
"Pyroglutamul-aprotinin, a new aprotinin homologue from
bovine lungs-isolation, properties, sequence analysis nad
characterization using ¹H nuclear magnetic resonance in
30 solution",
Biol Chem Hoppe-Seyler (1987), 368:1589-96.

SIEK88:

- 35 Siekmann, J, HR Wenzel, W Schroeder, and H Tschesche,
"Characterization and Sequence Determination of Six
Aprotinin homologues from bovine lungs",
Biol Chem Hoppe-Seyler (1988), 369:157-163.

SIEK89:

- 40 Siekmann, J, J Beckmann, A Mehlich, HR Wenzel,
H Tschesche, E Schnabel, W Mueller-Esterl,
"Immunological Characterization of Natural and
Semisynthetic Aprotinin Variants",
Biol Chem Hoppe-Seyler (1989), 370:677-81.

45

SILH77:

- Silhavy, TJ, HA Shuman, J Beckwith, and M Schwartz,
"Use of gene fusions to study outer membrane protein
localization in Escherichia coli",
50 Proc Natl Acad Sci USA (1977), 74(12)5411-5415.

T06230 55096260

SILH85:

Silhavy, TJ, and JR Beckwith,
 "Uses of lac Fusions for the Study of Biological Problems",

- 5 Microbiol Rev (1985), 49(4)398-418.

SINH90:

Sinha, S, HF Dovey, P Seubert, PJ Ward, RW Blacher,
 M Blaber, RA Bradshaw, M Arici, WC Mobley, and

- 10 I Lieberburg,
 "The Protease Inhibitory Properties of the Alzheimer's
 beta-amyloid Precursor Protein",
 J Biol Chem (1990), 265(16)8983-5.

15 SMIT85:

Smith GP,
 "Filamentous Fusion Phage: Novel Expression Vectors That
 Display Cloned Antigens on the Virion Surface",
 Science (1985), 228:1315-1317.

20

SMIT88a:

Smith, GP,
 "Filamentous Phage Assembly: Morphogenetically Defective
 Mutants That Do Not Kill the Host",

- 25 Virology (1988), 167:156-165.

SMIT88b:

Smith, GP,
 "Filamentous Phages as Cloning Vectors",
 Chapter 3 in Vectors: A Survey of Molecular Cloning
 Vectors and Their Uses, Editors: RL Rodriguez and DT
 Denhardt, Butterworth, Boston, 1988.

30

SODE85:

Sodergren, EJ, J Davidson, RK Taylor, and TJ Silhavy,
 "Selection for Mutants Altered in the Expression or Export
 of Outer Membrane Porin OmpF",
 J Bacteriol (1985), 162(3)1047-1053.

35

40 SOME85:

So, M, E Billyard, C Deal, E Getzoff, P Hagblom, TF Meyer,
 E Segal, and J Tainer,
 "Gonococcal Pilus: Genetics and Structure",
 Curr Top in Microbiol & Immunol (1985), 118:13-28.

45

SOMM89:

Sommerhoff, CP, GH Caughey, WE Finkbeiner, SC Lazarus,
 CB Basbaum, and JA Nadel,
 "A Potent Secretagogue for Airway Gland Serous Cells",
 J Immunol (1989), 142:2450-56.

50

105290 56096860

Sommerhoff, CP, JA Nadel, CB Basbaum, and GH Caughey,
"Neutrophil Elastase and Cathepsin G Stimulate Secretion
from Cultured Bovine Airway Gland Serous Cells",

- STAD86:
Stader, J, SA Benson, and TJ Silhavy ,
"Kinetic analysis of lamB mutants suggests the signal
10 sequence plays multiple roles in protein export",
J Biol Chem (1986), 261(32)15075-80.

Stader, J, LJ Gansheroff, and TJ Silhavy,
"New suppressors of signal-sequence mutations, prlG, are
linked tightly to the secE gene of Escherichia coli",
Genes & Develop (1989), 3:1045-1052.

20 States, DJ, TE Creighton, CM Dobson, and M Karplus,
"Conformations of intermediates in the folding of the
pancreatic trypsin inhibitor.",
J Mol Biol (1987), 195(3)731-9.

- STUB90:

- SUNX87:

- 45

Sutcliffe, MJ, I Haneef, D Carney, and TL Blundell,
"Knowledge based modelling of homologous proteins, part I:
three-dimensional frameworks derived from the simultaneous
superposition of multiple structures",
Protein Engineering (1987), 1:377-384.

Sutcliffe, MJ, FRF Hayes, and TL Blundell,
"Knowledge based modelling of homologous proteins, part
II: rules for the conformations of substituted
sidechains",
Protein Engineering (1987), 1:385-392.

Svendsen, IB,
"Amino Acid Sequence of Serine Protease Inhibitor CI-1
from Barley. Homology with Barley Inhibitor CI-2, Potato
Inhibitor I, and Leech Elgin",
Carlsberg Res Comm (1982), 47:45-53.

Swaim, MW, and SV Pizzo,
"Modification of the tandem reactive centres of human
inter- α -trypsin inhibitor with butanedione and cis-
dichlorodiammineplatinum(II)",
Biochem J (1988), 254:171-178.

Takahashi, H, S Iwanage, T Kitagawa, Y Hokama, and T Suzuki,

25 "Snake venom proteinase inhibitors. II. Chemical structure of inhibitor II isolated from the venom of Russell's viper (*Vipera russelli*).", J Biochem (1974), 76:721-733.

Takao, T, N Tominaga, S Yoshimura, Y Shimonishi, S Hara, T Inoue, and A Miyama,
"Isolation, primary structure and synthesis of heat-stable enterotoxin produced by Yersinia enterocolitica",
Eur J Biochem (1985), 152:199-206.

Takeda, T, GB Nair, K Suzuki, and Y Shimonishi,
"Production of a Monoclonal Antibody to Vibrio cholerae
Non-O1 Heat-Stable Enterotoxin (ST) Which is Cross-
Reactive with Yersinia enterocolitica ST",
Infection and Immunity (1990), 58(9)2755-9.

45 Tan, NH, and ET Kaiser,
"Synthesis and Characterization of a Pancreatic Trypsin
Inhibitor Homologue and a Model Inhibitor",
Biochemistry, (1977), 16:1531-41.

Theriault, NY, JB Carter, and SP Pulaski,
"Optimization of Ligation Reaction Conditions in Gene
Synthesis",

- THOR88:
Thornton, JM, BL Sibinda, MS Edwards, and DJ Barlow,
"Analysis, Design, and Modification of Loop Regions in
45 Proteins.",
BioEssays (?) SKG 3039 ??????

TOMM82:

Tomassen, J, P van der Ley, A van der Ende, H Bergmans,
and B Lugtenberg,

- "Cloning of ompF, the Structural Gene for an Outer
5 Membrane Pore Protein of E. coli K12: Physical
Localization and Homology with the phoE Gene",
Mol gen Genet (1982), 185:105-110.

TOMM85:

- 10 Tomassen, J, P van der Ley, M van Zeijl, and M Agterberg,
"Localization of functional domains in E. coli K-12 outer
membrane porins",
EMBO J (1985), 4(6)1583-7.

15 TRAB86:

Traboni, C, R Cortese,
"Sequence of a full length cDNA coding for human protein
HC (α_1 microglobulin)",
Nucleic Acids Res (1986), 14(15)6340.

20

TRIA88:

Trias, J, EY Rosenberg, and H Nikaido,
"Specificity of the glucose channel formed by protein D1
of Pseudomonas aeruginosa",

- 25 Biochim Biophys Acta (1988), 938:493-496.

TSCH86:

Tschesche, H, H Wenzel, R Schmuck, and E Schnabel,
"Homologues of Aprotinin with, in place of lysine, other
30 amino acids in position 15, process for their preparation
and their use as medicaments",
US Patent 4,595,674 (17 Jun 1986).

TSCH87:

- 35 Tschesch, H, J Beckmann, A Mehlich, E Schnabel,
E Truscheit, and HR Wenzel,
"Semisynthetic engineering of proteinase inhibitor
homologues",
Biochimica et Biophysica Acta (1987), 913:97-101.

40

VAND86:

van der Ley, P, M Struyve, and J Tomassen,
"Topology of outer membrane pore protein PhoE of
Escherichia coli. Identification of cell surface-exposed
45 amino acids with the aid of monoclonal antibodies",
J Biol Chem (1986), 261(26)12222-5.

VAND89:

- Vanderslacie, P, CS Craik, JA Nadel, GH Caughey,
 "Molecular Cloning of Dog Mast Cell Tryptase and a Related
 Protease: Structural Evidence of a Unique Mode of Serine
 5 Protease Activation",
 Biochem (1989), 28:4148-55.

VAND90:

- van der Werf, S, A Charbit, C Leclerc, V Mimic, J Ronco,
 10 M Girard, and M Hofnung,
 "Critical role of neighbouring sequences on the
 immunogenicity of the C3 poliovirus neutralization epitope
 expressed at the surface of recombinant bacteria",
 Vaccine (1990), 8(3)269-77.

VERS86a:

- Vershon, AK, K Blacker, and RT Sauer,
 "Mutagenesis of the Arc Repressor Using Synthetic Primers
 with Random Nucleotide Substitutions",
 20 pp243-256 in Protein Engineering. Applications in Science,
 Medicine, and Industry, Academic Press, 1986.

VERS86b:

- Vershon, AK, JU Bowie, TM Karplus, and RT Sauer,
 25 "Isolation and Analysis of Arc Repressor Mutants: Evidence
 for an Unusual Mechanism of DNA Binding",
 pp302-311 in Proteins: Structure, Function, and Genetics,
 Alan R. Liss, Inc., 1986.

VINC72:

- 30 Vincent &al,
 Biochem (1972), 11:2967ff.

VINC74:

- 35 Vincent &al.,
 Biochem (1974), 13:4205.

VITA84:

- Vita, C, D Dalzoppo, and A Fontana,
 40 "Independent Folding of the Carboxyl-Terminal Fragment
 228-316 of Thermolysin",
 Biochemistry (1984), 23:5512-5519.

VOGE86:

- 45 Vogel, H, and F Jahnig,
 "Models for the structure of outer membrane proteins of
E. coli derived from Raman spectroscopy and prediction
 methods",
 J Mol Biol (1986), 190:191-99.

50

FOI 56250-56096B50

VOND86:

Vonderviszt, F, GY Matrai, and I Simon,
 "Characteristic sequential residue environment of amino
 acids in proteins",

- 5 Int J Peptide Protein Res (1986), 27:483-92.

WACH79:

Wachter, E, K Hochstrasser, G Bretzel, and S Heindl,
 "Kunitz-Type Proteinase Inhibitors Derived by Limited
 10 Proteolysis of the Inter- α -trypsin Inhibitor, II.
 Characterization of a Second Inhibitory Inactive Domain by
 Amino Acid Sequence Determination",
 Hoppe-Seyler Z Physiol Chem (1979), 360:1297-1303.

15 WACH80:

Wachter, E, K Deppner, and K Hochstrasser,
 "A New Kunitz-type Inhibitor from Bovine Serum, Amino Acid
 Sequence Determination.",
 FEBS Letters (1980), 119:58-62.

20

WAGN78:

Wagner, G, K Wuthrich, and H Tschesche,
 "A H Nuclear-Magnetic-Resonance Study of the Solution
 Conformation of the Isoinhibitor K from Helix pomatia.",
 25 Eur J Biochem (1978), 89:367-377.

WAGN79:

Wanger, G, H Tschesche, and K Wuthrich,
 "The Influence of Localized Chemical Modifications of the
 30 Basic Pancreatic Trypsin Inhibitor on Static and Dynamic
 Aspects of the Molecular Conformation in Solution",
 Eur J Biochem (1979), 95:239-248.

WANG87:

Wagner, G, D Bruhwiler, and K Wuthrich,
 "Reinvestigation of the aromatic side-chains in the basic
 pancreatic trypsin inhibitor by heteronuclear
 two-dimensional nuclear magnetic resonance.",
 J Mol Biol (1987), 196(1)227-31.

40

WAIT83:

Waite, JH,
 "Evidence for a repeating 3,4-dihydroxyphenylalanine- and
 hydroxyproline-containing decapeptide in the adhesive
 45 protein of the mussel, Mytilus edulis L.",
 J Biol Chem (1983), 258(5)2911-5.

WAIT85:

Waite, JH, TJ Housley, and ML Tanzer,
 "Peptide repeats in a mussel glue protein: theme and
 variations.",

- 5 Biochemistry (1985), 24(19)5010-4.

WAIT86:

Waite, JH,
 "Mussel glue from Mytilus californianus Conrad: a
 comparative study. ",

- 10 J Comp Physiol [B] (1986), 156(4)491-6.

WATS87:

Molecular Biology of the Gene, Fourth Edition,

- 15 Watson, JD, NH Hopkins, JW Roberts, JA Steitz, and AM
 Weiner,
 Benjamin/Cummings Publishing Company, Inc., Menlo Park,
 CA., 1987.

20 WEBS78:

Webster, RE, and JS Cashman,
 "Morphogenesis of the Filamentous Single-stranded DNA
 Phages.", in The Single-Stranded DNA Phages, Denhardt, DT,
 D Dressler, and DS Ray editors, Cold Spring Harbor
 Laboratory, 1978., p557-569.

- 25

WEHM89:

Wehmeier, U, GA Sprenger, and JW Lengeler,
 "The use of lambda plac-Mu hybrid phages in Klebsiella
 pneumoniae and the isolation of stable Hfr strains",

- 30 Mol Gen Genet (1989), 215(3)529-36.

WEIN83:

Weinstock, GM, C ap Rhys, ML Berman, B Hampar, D Jackson,
 TJ Silhavy, J Weisemann, and M Zweig,
 "Open reading frame expression vectors: A general method
 for antigen production in Escherichia coli using protein
 fusions to beta-galactosidase",
 Proc Natl Acad Sci USA (1983), 80:4432-4436.

- 40

WELL86:

Wells, JA, and DB Powers,
 "In vivo Formation and Stability of Engineered Disulfide
 Bonds in Subtilisin",

- 45 J Biol Chem (1986), 261:6564-70.

WELL87a:

Wells, JA, BC Cunningham, TP Graycar, and DA Estell,
 "Recruitment of substrate-specificity properties from one
 enzyme into a related one by protein engineering",
 Proc Natl Acad Sci USA (1987), 84:5167-71.

- 50

- WELL87b:
Wells, JA, DB Powers, RR Bott, TP Graycar, and DA Estell,
"Designing substrate specificity by protein engineering of
electrostatic interactions",
5 Proc Natl Acad Sci USA (1987), 84:1219-23.
- WEMM83:
Wemmer, D, and NR Kallenbach,
10 Biochem (1983), 22:1901-6.
- WENZ80:
Wenzel, HR, and H Tschesche,
Hoppe-Seyler Z Physiol Chem (1980), 361:345.
- 15 WENZ81:
Wenzel, HR, and H Tschesche,
"'Chemical Mutation' by Amino Acid Exchange in the
Reactive Site of a Proteinase Inhibitor and Alteration of
Its Inhibitor Specificity",
20 Angew Chem Int Ed Engl (1981), 20(3)295-6.
- WETZ88:
Wetzel, R, et al.,
Proc Natl Acad Sci USA (1988), 85:401-5.
- 25 WEWE87:
Wewers, MD, MA Casolaro, SE Sellers, SC Swayze,
KM McPhaul, JT Wittes, and RG Crystal,
"Replacement therapy for α -1-antitrypsin deficiency
30 associated with emphysema",
New Engl J Med (1987), 316(17)1055-62.
- WHAR86:
Wharton, RP,
35 The Binding Specificity Determinants of 434 Repressor.,
Harvard U. PhD Thesis, 1986,
University Microfilms, Ann Arbor, Michigan.
- WIEC85:
40 Wieczorek, M, J Otlewski, J Cook, K Parks, J Leluk, A
Wilimowska-Pelc, A Polanowski, T Wilusz, and
L Laskowski, Jr,
"The Squash Family of Serine Protease Inhibitors. Amino
Acid Sequences and association equilibrium constants of
45 inhibitors from squash, summer squash, zucchini, and
cucumber seeds",
Biochem Biophys Res Comm (1985), 126(2)646-652.

WILK84:

- Wilkinson, AJ, AR Fersht, DM Blow, P Carter, and G Winter,
 "A large increase in enzyme-substrate affinity by protein
 engineering.",
 5 Nature (1984), 307:187-188.

WINT87b:

- Winter, AJ,
 "Outer membrane proteins of Brucella",
 10 Ann Inst Pasteur Microbiol (1987), 138(1)87-9.

WLOD84:

- Wlodawer, A, J Walter, R Huber, and L Sjolin,
 "Structure of bovine pancreatic trypsin inhibitor."
 15 Results of joint neutron and X-ray refinement of crystal
 form II.",
 J Mol Biol (1984), 180(2)301-29.

WLOD87a:

- Wlodawer, A, J Nachman, GL Gilliland, W Gallagher, and C
 20 Woodward,
 "Structure of form III crystals of bovine pancreatic
 trypsin inhibitor.",
 J Mol Biol (1987), 198(3)469-80.

25

WLOD87b:

- Wlodawer, A, J Deisenhofer, and R Huber,
 "Comparison of two highly refined structures of bovine
 pancreatic trypsin inhibitor.",
 30 J Mol Biol (1987), 193(1)145-56.

WOOD90:

- Woodward, SR, LJ Cruz, BM Olivera, and DR Hillyard,
 "Constant and hypervariable regions in conotoxin
 35 propeptides",
 EMBO J (1990), 9:1015-1020.

WUNT88:

- Wun, T-C, KK Kretzmer, TJ Girard, JP Miletich, and
 40 GJ Broze, Jr,
 "Cloning and Characterization of a cDNA Coding for the
 Lipoprotein-associated Coagulation Inhibitor Shows That It
 Consists of Three Tandem Kunitz-type Inhibitory Domains",
 J Biol Chem (1988), 263:6001-4.

45

T06290 56095860

Yager, TD, and PH von Hippel,
"Transcription Elongation and Termination in E. coli",
Volume 2, Chapter 76, p 1241-1275,
Escherichia coli and Salmonella typhimurium: Cellular and
Molecular Biology,
Neidhardt, FC, Editor-in-Chief,
Amer Soc for Microbiology, Washington, DC, 1987.

- 10 YANI85:
Yanisch-Perron, C, J Vieira, and J Messing,
"Improved M13 phage cloning vectors and host strains:
nucleotide sequences of the M13mp18 and pUC19 vectors",
Gene, (1985), 33:103-119.
- 15 YOKO77:
Yokosawa, H, and S-I Ishii,
"Anhydrotrypsin: New Features in Ligand Interactions
Revealed by Affinity Chromatography and Thionine
20 Replacement",
J Biochem (1977), 81:647-56.
- YOSH85:
Yoshimura, S, H Ikemura, H Watanabe, S Aimoto,
25 Y Shimonishi, S Hara, T Takeda, T Miwatani, and Y Takeda,
"Essential structure for full enterotoxigenic activity of
heat-stable enterotoxin produced by enterotoxigenic
Escherichia coli",
FEBS Lett (1985), 181:138-42.
- 30 ZAFA88:
Zafaralla, GC, C Ramilo, WR Gray, R Karlstrom, BM Olivera,
and LJ Cruz,
"Phylogenetic specificity of cholinergic ligands:
35 α -conotoxin SI",
Biochemistry, (1988), 27(18)7102-5.
- ZIMM82:
Zimmermann, R, C Watts, and W Wickner,
40 "The Biosynthesis of Membrane-bound M13 Coat Protein:
Energetics and Assembly Intermediates.",
J Biol Chem (1982), 257:6529-6536.
- ZOLL84:
Zoller, MJ, and M Smith,
45 "Oligonucleotide-Directed Mutagenesis: A Simple Method
Using Two Oligonucleotide Primers and a Single-Stranded
DNA Template.",
DNA (1984), 3(6)479-488.